

299-176

Spray or wash animals thoroughly. Do not apply within 28 days of slaughter. Do not use on animals under 2 months old. For Mange control, repeat treatment at 7-10 day intervals for two or more applications, as needed. In cases of light infestation one application may be adequate. Repeat as necessary.

**REPLENISHMENT OF DIPPING VATS** For Cattle: Add 1 gallon of Stock Tox 6115 to 100 gallons of water added to vat. For sheep: Add 1/2 gallon of Stock Tox 6115 to 100 gallons of water added to vat.

**WARNING**

**MAY BE FATAL IF SWALLOWED.** Do not breathe spray mist. Do not get in eyes, on skin or clothing. Wash thoroughly after using. Do not contaminate feed or foodstuffs.

**DO NOT USE, POUR, SPILL, OR STORE NEAR HEAT OR OPEN FLAME.**

This product is toxic to fish, birds, and other wildlife. Birds feeding on treated areas may be killed. Keep out of lakes, streams, or ponds. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

Do not reuse empty container. Destroy by crushing or perforating. Bury containers and wastes in a safe place away from water supplies.

**ANTIDOTE:** Give one tablespoonful of salt in a glass of warm water to induce vomiting. Repeat until vomit fluid is clear. Call a physician immediately.

Avoid spraying and dipping animals during cold, stormy weather. Avoid treating sick or overheated animals. Feed and water animals before treating. Spraying is recommended for young animals. If young animals must be dipped, individual hand immersion is essential to avoid swallowing of dip solution.

Do not use on dogs, cats, or other household pets or on poultry or in poultry houses. Use only as directed on this label.

Stock Tox forms a stable emulsion with water, but some settling may occur, especially in hard water. Thoroughly agitate mixture before dipping or spraying.

E.P.A. Reg. No. 299-176

27528



# STOCK TOX 6115

TOXAPHENE EMULSIFIABLE CONCENTRATE

FOR USE AS A LIVESTOCK SPRAY OR DIP TO CONTROL

SCABIES (PSOROPTIC, CHORIOPTIC AND SARCOPTIC MANGE), HORNFLIES, LICE, TICKS

**CONTENTS: 1 GALLON**

**WARNING: Keep Out of Reach of Children**

**SEE BACK PANEL FOR OTHER WARNINGS.**

**ACTIVE INGREDIENTS:**

- \*\* Toxaphene . . . . . 61.0%
- Petroleum Distillate . . . . . 23.8%

**INERT INGREDIENTS:**

. . . . . 15.2%

\*\* Technical Chlorinated Camphene (Chlorine content 67% to 69%)

606 W. Main St.

*C.J. Martin Company*

• Nac

*Martin's*  
ESTABLISHED 1883

# TOX 6115

MULSIFIABLE CONCENTRATE

LIVESTOCK SPRAY OR DIP  
TO CONTROL

PSOROPTIC AND SARCOPTIC  
MANGE, HORNFLIES, LICE, TICKS

CONTENTS: 1 GALLON

Keep out of Reach of Children.

SEE REVERSE PANEL FOR OTHER WARNINGS.

..... 61.0%  
..... 23.8%  
..... 15.2%

..... DDT (Chlorine content 67% to 69%)

*Martin Company*

**Nacogdoches, Texas 75961**

## DIRECTIONS

### BEEF CATTLE AND HORSES

To control Scabies (Psoroptic, Sarcoptic and Chorioptic Mange), Lone Star Ticks, Cattle Ticks, Winter Ticks, Gulf Coast Ticks, Hornflies, Cattle Lice, Ear Ticks: Mix 1 gallon with 150 gallons of water. Thoroughly wet animals by spraying or dipping. For ear ticks, use mist sprayer and apply into each ear and to head and neck. Repeat treatment at 7 to 10 day intervals for two or more applications as needed for scabies control. Repeat as necessary for control of other insects. Do not apply within 28 days of slaughter. Do not use on dairy cattle. Do not use on animals under 3 months old. Do not dip animals under 6 months old. If calves between 3 and 6 months old must be treated, apply only as a spray.

### BACK RUBBERS

As an aid in the control of Hornflies and Lice on beef cattle. Mix 1 quart with 5 gallons of kerosene or diesel fuel. Fill tank of backrubber device or soak burlap roll with solution so it is thoroughly wet but not to point of dripping. Recharge as necessary. Do not permit animals access to treatment within 28 days of slaughter. Do not permit access to backrubbers by dairy animals or calves under 6 months old.

### SHEEP AND GOATS

To control Psoroptic Sheep Scab, Fleeceworms, Hornflies, Lone Star Ticks, Ear Ticks: Mix 1 gallon with 150 gallons of water. Thoroughly wet animals by spraying or dipping. For ear ticks, use a mist sprayer and apply into each ear and to head and neck. Repeat treatment at 7 to 10 day intervals for two or more applications as needed for scabies control. Repeat as necessary for other insects.

To control Lice and Sheep Ticks (Keds): As a dip, use 1 gallon with 800 gallons of water. For spraying, 1 gallon with 200 gallons of water will give best results. Thoroughly wet animals. Repeat as necessary.

Do not apply within 28 days of slaughter. Do not use on dairy goats. Do not use on animals under 3 months old.

### HOGS

To control Lice, Sarcoptic Mange: Mix 1 gallon with 150 gallons of water.