

spoon of salt in a glass of warm water and repeat until vomit fluid is clear. Call a physician immediately.

Treated baits should be placed in locations not accessible to children, pets, wildlife and domestic animals or in tamper-proof bait boxes.

Do not contaminate water by cleaning of equipment or disposal of wastes.

Apply this product only as specified on this label. Do not reuse empty container. Destroy it by burying with waste or burning. Stay away from smoke or fumes.

**NOTE TO PHYSICIANS:** When humans have been known to have ingested RAT BAIT "D", blood transfusions, combined with intravenous injections and oral doses of Vitamin K are indicated as in the case of hemorrhage caused by overdoses of bishydroxycoumarin. This package contains enough bait to kill one or two animals.



# RAT BAIT "D" THROW PACK

WITH DIPHACINONE

Ready To Use Rat And Mouse Bait

**ACTIVE INGREDIENT:**

Diphacinone (2-Diphenylacetyl-1, 3-Indandione) . . . .005%

**INERT INGREDIENTS:** . . . . . 99.995%

**NET WEIGHT: 4 OUNCES**

**CAUTION: KEEP OUT OF REACH OF CHILDREN.**  
See Back Panel for Other Cautions.

Manufactured For

606 W. Main St. • C. J. Martin Co. • Nacogdoches, Texas 75961

E.P.A. Reg. No. 299.177

## MARTIN'S RAT BAIT "D" THROW PACK

**DIRECTIONS FOR USE:**

**RATS:** Place 1 or 2 RAT BAIT "D" THROW PACKS in protected spot next to burrows in runways or where rats are accustomed to feeding.

**MICE:** Place 1 ounce of RAT BAIT "D" pellets in runways where mice are accustomed to feeding at intervals of 8 to 12 feet.

Check bait stations daily to assure uninterrupted supply of bait for a period of at least 10 days for rats or 15 days for mice. Continue until all signs of feeding have stopped.

**LONG TERM CONTROL:** Bait continuously with MARTIN'S RAT BAIT "D". Stale, damp or moldy bait should be replaced with fresh bait.

**CAUTION:** Keep away from humans, domestic animals and pets. If swallowed, this material may reduce the clotting ability of the blood and serious hemorrhage may result. If baits are eaten, give a table-

**ACCEPTED**

Dec. 14, 1972

UNDER THE FEDERAL INSECTICIDE  
FUNGICIDE AND FIDENICING ACT  
FOR ECONOMIC POISON REGISTERED  
UNDER NO. 299-177


spoon of salt in a glass of warm water and repeat until vomit fluid is clear. Call a physician immediately.

Treated baits should be placed in locations not accessible to children, pets, wildlife and domestic animals or in tamper-proof bait boxes.

Do not contaminate water by cleaning of equipment or disposal of wastes.

Apply this product only as specified on this label. Do not reuse empty container. Destroy it by burying with waste or burning. Stay away from smoke or fumes.

**NOTE TO PHYSICIANS:** When humans have been known to have ingested RAT BAIT "D", blood transfusions, combined with intravenous injections and oral doses of Vitamin K are indicated as in the case of hemorrhage caused by overdoses of bishydroxycoumarin. This package contains enough bait to kill one or two animals.



# RAT BAIT "D" THROW PACK

WITH DIPHACINONE

Ready To Use Rat And Mouse Bait

**ACTIVE INGREDIENT:**

Diphacinone (2-Diphenylacetyl-1, 3-Indandione) . . . . .005%

**INERT INGREDIENTS:** . . . . . 99.995%

NET WEIGHT: 4 OUNCES

**CAUTION:** KEEP OUT OF REACH OF CHILDREN.  
See Back Panel for Other Cautions.

Manufactured For

606 W. Main St. • C. J. Martin Co. • Nacogdoches, Texas 75961

E.P.A. Reg. No. 299-177

## MARTIN'S RAT BAIT "D" THROW PACK

### DIRECTIONS FOR USE:

**RATS:** Place 1 or 2 RAT BAIT "D" THROW PACKS in protected spot next to burrows in runways or where rats are accustomed to feeding.

**MICE:** Place 1 ounce of RAT BAIT "D" pellets in runways where mice are accustomed to feeding at intervals of 8 to 12 feet.

Check bait stations daily to assure uninterrupted supply of bait for a period of at least 10 days for rats or 15 days for mice. Continue until all signs of feeding have stopped.

**LONG TERM CONTROL:** Bait continuously with MARTIN'S RAT BAIT "D". Stale, damp or moldy bait should be replaced with fresh bait.

**CAUTION:** Keep away from humans, domestic animals and pets. If swallowed, this material may reduce the clotting ability of the blood and serious hemorrhage may result. If baits are eaten, give a table-