



# COW & CALF DUST



**READY TO USE ON:**  
ADULT AND YOUNG ANIMALS

Kills

Hornflies, Lice, Ticks

on

Beef Cattle, Dairy Cattle, Sheep, Goats

Swine, Horses, Dogs

NET WEIGHT 1 POUND

**CAUTION: Keep Out of Reach of Children**

SEE BACK PANEL FOR OTHER CAUTIONS.

**CAUTION** Harmful if swallowed. Avoid breathing of dust. Avoid contact with skin. Wash thoroughly after using. Do not contaminate feed, water, or food-stuffs. Do not reuse empty container. Destroy container by burying with waste or burning. Stay away from smoke or fumes. Apply this product only as specified on this label.

This product contains Malathion, a cholinesterase inhibitor. Do not use this product simultaneously or within a few days of treatment with other drugs, insecticides, or chemicals having cholinesterase inhibiting activity.

This product is toxic to fish. Keep out of lakes, streams, or ponds. Do not apply where runoff is likely to occur. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

**ACTIVE INGREDIENTS**

Malathion 100%

Inert Ingredients

**INERT INGREDIENTS**

**DIRECTIONS**

**TO CONTROL HORNFLIES, LICE, AND TICKS ON BEEF CATTLE, NON-LACTATING DAIRY CATTLE, CALVES, SHEEP, GOATS, SWINE, AND HORSES:** Apply 1-2 ounces of dust to animal. Repeat at 14 day intervals if needed. Do not apply to lactating dairy animals for at least 2 weeks after treatment.

**TO CONTROL HORNFLIES ON LACTATING DAIRY CATTLE:**

Apply 1-2 ounces (2 rounded tablespoonsful) per animal to the back and neck. Rub powder into hair. Do not apply within 5 hours of milking or during milking. Repeat at 2 week intervals if needed.

**TO CONTROL TICKS AND FLEAS ON DOGS:**

Apply 1-2 ounces of dust to the back and neck of the dog. Rub powder into hair. Do not apply within 5 hours of bathing or during bathing. Repeat at 2 week intervals if needed.

Do not treat pregnant or nursing animals.

EPA Reg. No. 299-180

37101