

apply within 3 days of harvest on carrots, beets, head lettuce, turnips. Do not apply within 14 days of harvest on beet tops, leaf lettuce, spinach, turnip tops.

CABBAGE - BROCCOLI - CAULIFLOWER -- For aphids, armyworms, corn earworm, imported cabbage worms, diamondback moth, cabbage looper, flea beetles, Harlequin bugs. Do not apply within 3 days of harvest.

LAWNS -- To control chinch bugs, leafhoppers, fall armyworms, sod webworms, ants, fleas, mosquitoes, earwigs. Apply at the rate of 1 pound to 250 square feet. For best results, apply after rain or watering and do not water for at least two days after application. Repeat in two or three weeks if necessary.

CAUTION -- HARMFUL IF SWALLOWED OR INHALED. Avoid breathing dust. Avoid contact with skin. Wash thoroughly after handling. Do not reuse empty containers. Destroy by burying with waste or burning. Stay away from smoke or flame. Change contaminated clothing daily. Avoid contamination of feed and foodstuffs.

CAUTION -- THIS PRODUCT IS TOXIC TO BEES. Any use of the product when honey bees are visiting the crop will seriously reduce bee population. Bees are important pollinators and are essential to the production of many crops. Before using this product during such a period consult your State Agricultural Extension Service.

THIS PRODUCT IS TOXIC TO FISH. Keep out of lakes, streams, or ponds. Do not apply where runoff is likely to occur. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.


NOTE FOR PHYSICIAN: Carbaryl is a moderate reversible cholinesterase inhibitor. Atropine is an antidote.

E.P.A. Reg. No. 299-182

47421



VEGETABLE DUST



CONTAINS SEVIN® AND ROTENONE

For Control Of
MANY DESTRUCTIVE INSECT PESTS
IN GARDENS

NET WEIGHT: 1 POUND

CAUTION: Keep out of reach of children
SEE BACK PANEL FOR OTHER CAUTIONS.

® Trademark of Union Carbide Corporation for 1-naphthyl N-methylcarbamate

ACTIVE INGREDIENTS:

Carbaryl (1-naphthyl N-methylcarbamate) 5%
Rotenone 1%
Other Cube Resins 2%

INERT INGREDIENTS 92%

DIRECTIONS

Begin application when insect damage is first observed and repeat on a regular 7 to 10 day schedule or as necessary for good insect control. Dust upper and lower leaf surfaces thoroughly.

BEANS -- For aphids, armyworms, bean leaf beetles, cucumber beetles, flea beetles, leafhoppers, Mexican bean beetles, stink bugs, tarnished plant bugs. Do not apply within 1 day of harvest.

PEAS -- For aphids, armyworms, leafhoppers, pea weevils, grasshoppers. Do not apply within 1 day of harvest.

CORN -- For aphids, fall armyworms, adult corn rootworms, European corn borer, flea beetles, leafhoppers. Do not apply within 1 day of harvest. For corn earworms, apply to silks as soon as they appear. Make 3 to 4 applications at 2 to 3 day intervals as infestation warrants. Reapply immediately after rain. Application of this product to the tassel region of corn during pollen shed period will seriously reduce bee population. For European corn borer, apply thoroughly into the whorls when plants are about knee high. Repeat at weekly intervals until tassels emerge.

POTATOES - TOMATOES - PEPPER - EGGPLANT -- For aphids, flea beetles, Colorado potato beetles, leafhoppers, stink bugs, tarnished plant bug, tomato fruitworm, tomato hornworm. Do not apply within 1 day of harvest. Apply to foliage only on potatoes.

CUCUMBERS - SQUASH - PUMPKINS - MELONS -- For aphids, cucumber beetles, flea beetles, melonworm, leafhoppers, pickleworm, squash bug. Do not apply within 1 day of harvest.

OKRA -- For aphids, corn earworm, stink bugs. Do not apply within 1 day of harvest.

CARROTS - BEETS - LETTUCE - SPINACH - TURNIPS -- For aphids, armyworms, flea beetles, leafhoppers, stink bugs. Do not

606 W. Main St.

C. J. Martin Company

● Nacogdoches, Texas 75961