

5399456

JUL 18 1991

Mr. Mike Compton
C.J. Martin Company
606 West Main Street
Nacogdoches, TX 75961

Dear Mr. Compton:

**Subject: Martins Vegetable Dust
EPA Registration No. 299-182
Your Submission of a Confidential Statement of Formula and
Finished Labels in Response to the Registration Standard
for Rotenone**

The label you submitted is acceptable to satisfy the requirements of the Registration Standard. A stamped copy of your label is enclosed.

The Confidential Statement of Formula you submitted will be placed in the file for this product. Further revisions may be required if changes are made in labels for Manufacturing-Use Products supplying the active ingredients for this product.

Sincerely yours,



Phil Hutton
Product Manager (17)
Insecticide-Rodenticide Branch
Registration Division (H7505C)

Enclosure

52343:I:Schroeder:C.Disk:KEVRIC:06/14/91:CL:EK:CL

CONCURRENCES

SYMBOL							
SURNAME							
DATE							

Begin application when insect damage is first observed and repeat on a regular 7 to 10 day schedule or as necessary for good insect control. Dust upper and lower leaf surfaces thoroughly.

BEANS: For aphids, armyworms, bean leaf beetles, cucumber beetles, flea beetles, leafhoppers, Mexican bean beetles, stink bugs, tarnished plant bugs. Do not apply within 1 day of harvest.

PEAS: For aphids, armyworms, leafhoppers, pea weevils, grasshoppers. Do not apply within 1 day of harvest.

CORN: For aphids, fall armyworms, adult corn rootworms, European corn borers, flea beetles, leafhoppers. Do not apply within 1 day of harvest. For corn earworms, apply to silks as soon as they appear. Make 3 to 4 applications at 2 to 3 day intervals as infestation warrants. Reapply immediately after rain. Application of this product to the tassel region of corn during pollen shed period will seriously reduce bee population. For European corn borers, apply thoroughly into the whorls when plants are about knee high. Repeat at weekly intervals until tassels emerge.

POTATOES, TOMATOES, PEPPERS, EGGPLANTS: For aphids, flea beetles, Colorado potato beetles, leafhoppers, stink bugs, tarnished plant bugs, tomato fruitworms, tomato hornworms. Do not apply within 1 day of harvest. Apply to foliage only on potatoes.

CUCUMBERS, SQUASH, PUMPKINS, MELONS: For aphids, cucumber beetles, flea beetles, melonworms, leafhoppers, pickleworms, squash bugs. Do not apply within 1 day of harvest.

OKRA: For aphids, corn earworms, stink bugs. Do not apply within 1 day of harvest.

CARROTS, BEETS, LETTUCE, SPINACH, TURNIPS: For aphids, armyworms, flea beetles, leafhoppers, stink bugs. Do not apply within 3 days of harvest on carrots, beets, head lettuce, turnips. Do not apply within 14 days of harvest on beet tops, leaf lettuce, spinach, turnip tops.

CABBAGE, BROCCOLI, CAULIFLOWER: For aphids, armyworms, corn earworms, imported cabbage worms, diamondback moths, cabbage loopers, flea beetles, Harlequin bugs. Do not apply within 3 days of harvest.

SUBSEQUENT PLANTINGS OF FOOD AND FEED CROPS IN TREATED SOIL MUST BE MADE WITH CROPS ON THIS OR OTHER CARBARYL LABELS.

LAWNS: To control chinch bugs, leafhoppers, fall armyworms, sod webworms, ants, fleas, mosquitoes, earwigs. Apply at the rate of 1 pound to 250 square feet. For best results, apply after rain or watering and do not water for at least two days after application. Repeat in two or three weeks if necessary.

E.P.A. Reg. No. 299-182
E.P.A. Est. No. 299-182

1001

Manufactured and
packaged for the

299-182




Martin's

ESTABLISHED 1883

4765

VEGETABLE DUST



Contains SEVIN® brand carbaryl insecticide and ROTENONE for control of many destructive pests in gardens.

NET WEIGHT:
1 POUND

KEEP OUT OF REACH
OF CHILDREN
CAUTION

See Back Panel For Other Cautions

ACTIVE INGREDIENTS:

Carbaryl
(1-naphthyl N-methylcarbamate)..... 5%
Rotenone 1%
Other Cube Resins 2%
INERT INGREDIENTS: 92%

© Registered Trademark of Rhone Poulenc for carbaryl insecticide

PRECAUTIONS

HAZARD AND DANGER:
CAUTION: May be harmful if inhaled. Avoid contact with eyes. Wash thoroughly after handling.

ENVIRONMENT:
This product is highly toxic to fish and aquatic life. Do not apply this product to crops or weeds if bees are present.

This pesticide is extremely toxic to water and wetland organisms. Runoff and drainage are hazardous to aquatic organisms. Do not contaminate water or washwaters.

NOTE TO PHYSICIAN:
sible cholinesterase inhibitor.

DIRECTIONS

It is a violation of Federal law to use this product in a manner inconsistent with the label directions.

RE-ENTRY

Do not apply this product to crops or through drift exposure. The area being treated must be restricted to authorized persons. Do not enter treated areas until dusts have settled. Clothing worn in treated areas may require more restrictive decontamination procedures. Written or oral warnings must be given to persons who are expected to be in treated areas. Warnings must be given if there is a possibility of drift. Warnings cannot be under any circumstances.

STORAGE

STORAGE: Do not store in areas where children, pets, and animals are present. Store in a cool, dry place until used. Store in original container and keep tightly closed. Do not reuse empty containers. Dispose of empty containers by burning. If burned, do not breathe fumes. Sulfiting from the use of this product may occur on site or at an approved disposal site.

C. J. M...
808 W...
Nacogdoches

BEST AVAILABLE