



LINDANE CONCENTRATE

Emulsifiable

To be diluted with water

For the control of certain insects on beef cattle, sheep, goats, hogs and for the control of certain insects on fruit trees, vegetables, flowers, and, ornamentals.

CONTENTS: 1 QUART

CAUTION:
KEEP OUT OF REACH OF CHILDREN.
SEE BACK PANEL FOR OTHER CAUTIONS.

ACTIVE INGREDIENTS:

Lindane (Gamma isomer of benzene hexachloride) 10.60%
Aromatic Petroleum Distillate 85.32%
INERT INGREDIENTS: 4.00%

ACCEPTED
AUG 12 1975
Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No.

Cabbage, Celery, Collards, Cucumbers, Lettuce, Okra, Peppers, Squash, Melons, Tomatoes: White grubs, Cutworms, Wireworms. For preplant soil application only. Use 4 gallons for each 250 square feet. Work into soil thoroughly before or during planting. Do not treat land to be planted in root crops.

Bulbs (Daffodil, Gladiolus, Hyacinth, Iris, Lilies, Narcissus, Tulips): Aphids, Bulb fly, European corn borer, Iris borer, Thrips. Apply when insects appear. Repeat as necessary.

Flowers (Carnations, Chrysanthemums, Coleus, Daisies, Gardenia, Geraniums, Hibiscus, Nasturtiums, Pansies, Petunias, Sweet Peas, Zinnias): Aphids, blister beetles, Leafhoppers, Thrips. Apply when insects appear. Repeat as necessary.

Ornamentals (Azaleas, Camellias, Heather, Lilacs, Rhododendrons, Roses): Aphids, Blackvine weevil, Cucumber beetles, Thrips, White fly. Apply when insects appear. Repeat as necessary.

CAUTION: Harmful if swallowed. May be absorbed through skin. Avoid breathing spray mist. Avoid contact with skin. In case of contact, wash immediately with soap and water. In case of contact with eyes, flush with water and get medical attention.

This product is toxic to fish, birds, and other wildlife. Keep out of lakes, streams, or ponds. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

Do not reuse empty container. Destroy it by perforating or crushing. Bury or discard in a safe place.

WARNING:

FLAMMABLE - KEEP AWAY FROM HEAT OR OPEN FLAME.

CAUTION: Use only as a spray. Do not treat animals under 3 months of age. Do not treat sows within two weeks of farrowing or for at least 3 weeks thereafter. Do not use on dairy animals. Do not use within 30 days of slaughter. Avoid contamination of feed and foodstuffs. Do not use this product in poultry houses, dairy barns, or milk rooms. Avoid treating animals during cold, stormy weather. Avoid treating sick or overheated animals. Do not use on household pets or humans.

This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

E.P.A. Reg. No. 299-184

E.P.A. Est. No. 299-TX-1

DIRECTIONS FOR USE

Measure accurately - Stir thoroughly before

HORNFLIES ON BEEF CATTLE: Spray animals thoroughly with Lindane solution prepared by adding 2 tablespoonfuls Lindane Concentrate to 2 gallons of water or 1 pint to 40 gallons. Repeat as necessary.

LICE AND TICKS ON BEEF CATTLE, HORSES, SHEEP, animals with 0.03 Lindane solution prepared by adding Lindane Concentrate to 2 gallons of water or 1 pint to 40 gallons. Repeat as necessary.

LICE ON HOGS: Spray animals with 0.06 Lindane solution prepared by adding 4 tablespoonfuls Lindane Concentrate to 2 gallons of water or 1 pint to 20 gallons. Repeat as necessary. Do not apply to young animals.

SARCOPTIC MANGE ON BEEF CATTLE AND HOGS AND MANGE (SHEEP SCABIES) ON SHEEP AND GOATS: Spray animals with 0.03 Lindane solution prepared by adding 2 tablespoonfuls Lindane Concentrate to 2 gallons of water. Wet animals thoroughly. Use 2 or more treatments at 10 day intervals as needed. Pens and shelters should also be treated to prevent reinfestation.

FLIES, ROACHES, SPIDERS, ANTS, MOSQUITOES IN LIVING AND BUILDINGS: Spray with a 0.3 Lindane solution prepared by adding 3 tablespoonfuls Lindane Concentrate to 1 gallon of water or 1 pint to 20 gallons. Apply 1 gallon of mix for each 500 square feet of surface. Thoroughly wet, about to run off. For crawling insects apply thorough treatment to crevices, baseboards and other hiding places. Remove animals from area. Do not contaminate milk or milking equipment, feed or manure.

INSECTICIDAL PAINT FOR FLIES, ROACHES, SPIDERS, AND TOES: Use 1.25 Lindane solution prepared by adding 1 1/2 tablespoonfuls Lindane Concentrate to 1 gallon of water. Apply with a brush on and around doors, windows, beams, baseboards, cracks, crevices, and other areas as a spot treatment. This concentration may blister paint. Do not contaminate milk, milking equipment, feed or manure.

FRUIT TREES, VEGETABLES, FLOWERS, AND ORNAMENTALS: Use 0.03 Lindane solution as directed below. Prepare 0.03 Lindane solution by adding 2 tablespoonfuls Lindane Concentrate to 2 gallons of water or 1 pint to 40 gallons of water.

Peaches, Plums: Aphids, Leafhoppers, Thrips. Spray to control insects. Do not apply within 60 days of harvest.

Tomatoes: Aphids. Use 1 gallon per 150 square feet. Do not spray during fruiting.

606 W. Main St.

C. J. Martin Company

Nacogdoches, Texas 75961

ry, Collards, Cucumbers, Lettuce, Okra, Peppers, Squash,
 es: White grubs, Cutworms, Wireworms: For preplant soil
 y. Use 4 gallons for each 250 square feet. Work into soil
 ore or during planting. Do not treat land to be planted in
 l, Gladiolus, Hyacinth, Iris, Lilies, Narcissus, Tulips):
 fly, European corn borer, Iris borer, Thrips. Apply when
 . Repeat as necessary.
 ions, Chrysanthemums, Coleus, Daisies, Gardenia, Gerani-
 , Nasturtiums, Pansies, Petunias, Sweet Peas, Zinnias):
 r beetles, Leafhoppers, Thrips: Apply when insects appear.
 ssary.
 azaleas, Camellias, Heather, Lilacs, Rhododendrons, Roses):
 vine weevil, Cucumber beetles, Thrips, White fly: Apply
 appear. Repeat as necessary.

Harmful if swallowed. May be absorbed through skin.
 ng spray mist. Avoid contact with skin. In case of con-
 mediately with soap and water. In case of contact with
 ith water and get medical attention.

is toxic to fish, birds, and other wildlife. Keep out of
 , or ponds. Do not apply where runoff is likely to occur.
 when weather conditions favor drift from areas treated.
 minate water by cleaning of equipment or disposal of
 y this product only as specified on this label.*

empty container. Destroy it by perforating or crushing.
 rd in a safe place.

WARNING:

FLAME - KEEP AWAY FROM HEAT OR OPEN FLAME.

Use only as a spray. Do not treat animals under 3 months
 ot treat sows within two weeks of farrowing or for at least
 eafter. Do not use on dairy animals. Do not use within
 slaughter. Avoid contamination of feed and foodstuffs.
 is product in poultry houses, dairy barns, or milk rooms.
 g animals during cold, stormy weather. Avoid treating
 eated animals. Do not use on household pets or humans.

is highly toxic to bees exposed to direct treatment or
 crops. Protective information may be obtained from your
 Agricultural Extension Service.

E.P.A. Reg. No. 299-184
 E.P.A. Est. No. 299-TX-1



LINDANE CONCENTRATE

Emulsifiable

To be diluted with water

For the control of certain insects on beef cattle, sheep,
 goats, hogs and for the control of certain insects on
 fruit trees, vegetables, flowers, and ornamentals.

CONTENTS: 1 QUART

CAUTION:
 KEEP OUT OF REACH OF CHILDREN.
 SEE BACK PANEL FOR OTHER CAUTIONS.

ACTIVE INGREDIENTS:
 Lindane (Gamma isomer of benzene hexachloride) . . . 10.68%
 Aromatic Petroleum Distillate 85.32%
INERT INGREDIENTS: 4.00%

ACCEPTED
 AUG 12 1975
 Under the Federal Insecticide,
 Fungicide, and Rodenticide Act,
 as amended, for the pesticide
 registered under
 EPA Reg. No. 299-184

DIRECTIONS FOR USE

Measure accurately - Stir thoroughly before use.

HORNFLIES ON BEEF CATTLE: Spray animals thoroughly with 0.07 Lindane solution prepared by adding 2 tablespoonfuls Lindane Concentrate to 2 1/2 gallons of water or 1 pint to 40 gallons. Repeat as necessary.

LICE AND TICKS ON BEEF CATTLE, HORSES, SHEEP, GOATS: Spray animals with 0.03 Lindane solution prepared by adding 2 tablespoonfuls Lindane Concentrate to 2 1/2 gallons of water or 1 pint to 40 gallons. Wet skin thoroughly. Repeat as necessary.

LICE ON HOGS: Spray animals with 0.06 Lindane Solution prepared by adding 4 tablespoonfuls Lindane Concentrate to 2 1/2 gallons of water or 1 pint to 20 gallons. Repeat as necessary. Do not apply to young animals.

SARCOPTIC MANGE ON BEEF CATTLE AND HOGS AND PSOROPTIC MANGE (SHEEP SCABIES) ON SHEEP AND GOATS: Spray animals with 0.03 spray prepared by adding 2 tablespoonfuls Lindane Concentrate to 2 1/2 gallons of water. Wet animals thoroughly. Use 2 or more treatments at 7 to 10 day intervals as needed. Pens and shelters should also be sprayed to prevent reinfestation.

FLIES, ROACHES, SPIDERS, ANTS, MOSQUITOES IN LIVESTOCK PENS AND BUILDINGS: Spray with a 0.3 solution prepared by adding 6 tablespoonfuls Lindane Concentrate to 1 gallon of water or 1 pint to 4 gallons. Apply 1 gallon of mix for each 500 square feet of surface area or until thoroughly wet, about to run off. For crawling insects apply thoroughly to cracks, crevices, baseboards and other hiding places. Remove animals before spraying. Do not contaminate milk or milking equipment, feed or drinking water.

INSECTICIDAL PAINT FOR FLIES, ROACHES, SPIDERS, ANTS, MOSQUITOES: Use 1.25 Lindane solution prepared by adding 1 pint Lindane Concentrate to 1 gallon of water. Apply with a brush on and around sills, doors, windows, beams, baseboards, cracks, crevices, and other hiding places as a spot treatment. This concentration may blister paint. Do not use in the home. Do not contaminate milk, milking equipment, feed or drinking water.

FRUIT TREES, VEGETABLES, FLOWERS, AND ORNAMENTALS: Apply 0.07 Lindane solution as directed below. Prepare 0.07 solution by adding 2 tablespoonfuls Lindane Concentrate to 2 1/2 gallons of water or 1 pint to 40 gallons of water.

Peaches, Plums: Aphids, Leafrollers, Thrips. Spray trees thoroughly. Do not apply within 60 days of harvest.
Tomatoes: Aphids. Use 1 gallon per 150 square feet. Do not apply after fruit starts to form.

606 W. Main St.

C. J. Martin Company

Nacogdoches, Texas 75961