



# RABON POULTRY DUST


**CAUTION:** Harmful if swallowed. Avoid breathing dust. Do not get in eyes, on skin, or on clothing. If the material gets into the eyes, wash with plenty of water for 15 minutes; if irritation persists, see a physician. Wash thoroughly with soap and water after handling and before eating or smoking. Avoid contamination of feed and foodstuffs. Keep away from heat and open flame. Do not apply in dwellings.

This product is toxic to fish. Keep out of lakes, streams or ponds. Do not contaminate water by cleaning of equipment or disposal of wastes. Do not apply where runoff is likely to occur. Apply this product only as specified on this label.

Do not reuse empty container. Destroy it by perforating or crushing. Bury or discard in a safe place.

E.P.A. Reg. No. 299-187  
E.P.A. Est. No. 299-TX-1

Controls Northern Fowl mites, chicken mites, and lice on or around chickens.



NET WEIGHT: 1 POUND

**CAUTION:**  
**Keep Out of Reach of Children**  
SEE BACK PANEL FOR OTHER CAUTIONS

ACTIVE INGREDIENT:  
2-chloro-1-(2,4,5-trichlorophenyl) vinyl dimethyl phosphate . . . 3.00%  
INERT INGREDIENTS: . . . . . 97.00%

### DIRECTIONS

**BIRDS HOUSED IN WIRE CAGES:** For control of Northern Fowl mites, chicken mites, and lice, use 1 lb. of dust per 300 birds. Apply with a plunger or rotary type duster or shaker-can. For individual treatment, direct dust to vent and fluff area. Group treatment may be preferred. Dust should reach skin. Do not repeat more often than every 14 days. Wire rungs and corners should also be treated.

For northern fowl mites on roosters, thorough individual application of dust will assure long-lasting control and reduce reinfestation of breeding flocks.

**FLOOR MANAGEMENT:** For the control of Northern Fowl mites, chicken mites, and lice.

**LITTER TREATMENT:** Use 1 pound of dust to 100 sq. ft. Apply with a plunger or rotary type duster. Treat evenly and thoroughly. Also treat roosts, cracks, and crevices where pests may hide.

**DUST BOX:** Use 2 pounds of dust per 100 birds. Use a box about 2 ft. by 2 ft. by 1 ft. deep.

**ROOST PAINT:** Use 1 pound of dust per 100 ft. Make a thick slurry by mixing 1 pound of dust with 1 pint of water. Continually stir. Treat roosts thoroughly, particularly the cracks and crevices.

606 W. Main St.

*C. J. Martin Company*

● Nacogdoches, Texas 75961