HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Avoid breathing of dust. Avoid contact with skin, eyes, and clothing. Avoid contamination of. feed and foodstuffs. Wash thoroughly after handling. Harmful if swallowed. May be absorbed through skin. Treat lawns in late afternoon. Keep children and pets off treated areas until the following afternoon.

ENVIRONMENTAL HAZARDS

This product is toxic to fish, birds. and other wildlife. Birds feeding on treated areas may be killed. Keep out of lakes, streams, or ponds. Do: not contaminate water by cleaning? of equipment or disposal of wastes.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

STORAGE & DISPOSAL

STORAGE: Store in original container until used. Do not store, use, or spill near heat or open flame. Store out of reach of children and domestic animals. Do not contaminate water. 52 food, or feed by storage or disposal. 2 DISPOSAL: Do not reuse empty bag. Discard bag in trash.

Overall Lawn Application: For control of Chinch Bugs, Earwigs, Ticks, Ants, Lawn Moths (Sod Webworms), & Bermuda Mites, Crickets, Army- & worms, Brown Dog Ticks, Chiggers, Clover Mites, Cutworms, Fleas, Leafhoppers, Sowbugs, Millipedes, Springtails (Collembola) en grass

ACDGDOCHES, TX. 75963-1089



THIS PRODUCT MUST NOT BE USED ON GOLF COURSES OR SOD FARMS

DIAZINON° GRANULAR LAWN INSECT CONTROL

KILLS:

CHINCH BUGS ARMYWORMS BERMUDA MITES BILLBUGS **BROWN DOG TICKS** CHIGGERS **CLOVER MITES** CRICKETS **CUTWORMS** DIGGER WASPS **EARWIGS**



LAWN MOTHS LEAFHOPPERS SOD WEBWORMS WHITE GRUBS OF:

JAPANESE BEETLES

• EUROPEAN CHAFERS

•SOUTHERN CHAFERS

KEEP OUT OF REACH OF CHILDREN CAUTION

SEE OTHER PRECAUTIONARY STATEMENTS ON SIDE

NET WEIGHT: 25 POLINDS

ACTIVE INGREDIENT:

0, 0, diethyl 0-(2-isopropyl-6-methyl-4-pyrimidinyl) phosphorothicate 2.0% INERT INGREDIENTS: 98.0%

Diazinon ® - Registered Trademark of Ciba-Geigy Corporation

DIRECTIONS FOR USE (Continued)

and dichondra lawns, apply Diazinon @ Granular Lawn Insect Control at a rate of 8 oz, per 100 sq. ft. or 5 lbs. per 1000 sq. ft. For heavy lawns such as St. Accustine lawns application should be increased to 12 oz. per 100 sq. ft. or 7 1/2 lbs. per 1000 sq. ft. Make even application over entire lawn when grass is dry. Water lawn lightly after application Repeat as necessary.

For control of Billbugs, and White Grubs of Japanese Beetle, European Chafer and Souther Chafer, apply 6 lbs. per 1000 sq. ft. of lawn. Water grass thoroughly after application. For grub control make application any time between late July and early October. For Billbug control, apply when activity is first observed. Repeat as necessary.

Spot Treatment: For Ants apply one tablespoon of Diazinon® Granular Lawn insect Control over and around each individual ant hill.

For Digger Wasps, apply one tablespoon in and around each individual ground nest opening. Make application about dusk when the wasps are not active.

Band Treatment Around Foundation: For Ants, Brown Dog Ticks, Clover Mites, Crickets, Earwigs, Fleas, Millipedes, Sowbugs, and Springtails found around or near the foundation of houses, apply a five foot band of Diazinon® Granular Lawn insect Control next to the foundation of the house. Distribute the granules uniformly using a rate of 8 oz. for each 5 ft. by 20 ft. area to be treated.

> E.P.A. Reg. No. 299-194 E.P.A. Est. No. 299-TX-1

> > 606 W. MAIN ST.

