
PM 13. 299-194



PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Harmful if swallowed or absorbed through skin. Causes moderate eye injury. Avoid contact with skin, eyes, or clothing. Wash thoroughly with soap and water after handling and before eating or smoking. Do not store near food or feed products. Food utensils such as spoons or measuring cups should not be used for food purposes after use in measuring pesticides. STATEMENT OF PRACTICAL TREATMENT

If Swallowed: Call a physician immediately. Drink one or two glasses of water and induce vomiting by touching the back of the throat with finger. Repeat until vomit fluid is clear. Do not induce vomiting or give anything by mouth if person is unconscious or convulsing.

If Inhaled: Remove victim to fresh air and apply artifical respiration if indicated.

If On The Skin: Wash promptly with soap and water. Rinse thoroughly. Get medical attention.

If In Eyes: Rinse eyes with plenty of water and call a physician immediately.

NOTE TO PHYSICIAN: This product is an organophosphate insecticide. If symptoms of cholinesterase inhibition are present, atropine sulfate by injection is antidotal. 2-PAM is also antidotal and may be administered, but only in conjunction with atropine.

ENVIRONMENTAL HAZARDS

This pesticide is highly toxic to birds, fish and other wildlife. Birds, especially waterfowl, feeding or drinking on treated areas may be killed. Because of the migratory habits of certain Atlantic Coast waterfowl, do not apply this product to lawns in Nassau County, New York, between November 1 and May 20. Do not exceed maximum permitted label rates. Application rates above those recommended significantly increase potential hazards to birds and waterfowl. Avoid overlapping of granules. For lawns, where irrigation (watering) is recommended in the directions, or if waterfowl, i.e., ducks or geese, can be expected in the treated area after treatment (except spot treatment), water lawn with at least ¼ inch of water immediately after applying this product, however, stop watering before puddling occurs. Keep out of lakes, streams, ponds, tidal marshes, and estuaries. Do not apply directly to water, to areas where surface water is present, or to intertidat areas below the mean high water mark. Runolf may be hazardous to aquatic organisms in neighboring areas. Shrimp and crab may be killed at application rates recommended on this label. Do not apply where fish, shrimp, crab, and other aquatic tile are important resources. Do not contaminate water by cleaning of equipment or disposal of equipment wash waters.

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

١

STORAGE AND DISPOSAL

STORAGE: Do not contaminate water, food, or feed by storage or disposal. Store in original container in a cool, dry, wellventilated area inaccessible to children and domestic animals. Do not store near fertilizers, herbicides, seeds, food, or feed. DISPOSAL: Do not reuse empty can. Wrap in several layers of newspaper and discard in trash.

SPOT TREATMENT

ANTS: Apply one tablespoon over and around each individual ant hill.

FIRE ANTS: To aid in the control of fire ants in home lawns and other outdoor home recreational areas, apply 1% cups over, and 2 ft. around, each fire ant mound. Immediately after application, water in thoroughly. A minimum of 1 gallon of water should be applied to each mound area. Apply water gently to avoid disturbing ants. Use of high pressure watering equipment may disturb the ants and cause migration, reducing product effectiveness. Treat as necessary.

NOTE: For best results, apply in cool weather, 65-80°F or in the early morning or late evenings hours. Treat new mounds as they appear.

DIGGER WASPS: Apply one tablespoon in and around each individual ground nest opening. Make the application about dusk when the wasps are not active.

BAND TREATMENT AROUND FOUNDATION. The following insects are commonly found around or near the foundation of houses as well as in the lawn. Ants, Brown Dog Ticks, Clover Mites, Crickets, Earwigs, Fleas, Millipedes, Sowbugs, Springtails (Collembola). For control of the above insects, apply a live-foot band of Diazinon Granular Lawn Insect Control next to the foundation of the house. Distribute the granules uniformly using a rate of 8 ozs. for each 5' by 20' area to be treated.

DIRECTIONS FOR USE AND CONDITIONS OF SALE AND WARRANTY

IMPORTANT: Read the entire Directions for Use and the Conditions of Sale and Warranty before using this product. If terms are not acceptable, return the unopened product container at once.

CONDITIONS OF SALE AND WARRANTY

The Directions for Use of this product reflect the opinion of experts based on field use and tests. The directions are believed to be reliable and should be followed carefully. However, it is impossible to eliminate all risks inherently associated with use of this product. Crop injury, ineffectiveness, or other unintended consequences may result because of such factors as iven ther conditions, presence of other materials, or the manner of use or application all of which are beyond the control of C.J. Martin Company or the Siller. All such risks shall be assumed by the buyer. C.J. Martin Company warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes referred to in the Directions for Use subject to the inherent risks referred to above. C.J. Martin Company makes no other express or implied warranty of Fitness or Merchar.t ability or any other express or implied warranty. In no case shall C.J. Martin Company or the Seller be liable for consequentfal, special, or indirect damages resulting from the use or handling of this product. C.J. Martin Company and the Seller offer that product. and the Buyer and user accept it, subject to the foregoing Conditions of Sale and Warranty, which may be varied bely by agreement in writing signed by a duly authorized representative of C.J. Martin Company.

E.P.A. Reg. No. 299-194 E.P.A. Est. No. 299-TX-1

C.J. MARTIN COMPANY 606 W. Main Street Nacogdoches, TX 75961

BEST AVAILABLE COTY

. •

3013

CONTROLS: Brown Dog Ticks, Chinch Bugs, Sod Webworms, Lawn Moths, Earwigs, Fleas, Fire Ants*, White Grubs of Japanese Beetle, European Chater, Southern Chater, Billbugs and many others. *Aids In Control.

DIRECTIONS FOR USE

OVERALL LAWN APPLICATION

Diazinon Granular Lawn Insect Control Iwan pests on both grass and dichondra lawns. Apply to your lawn with a fertilizer spreader at the rate shown on chart. For application with other types of equipment, apply at the rate of 5 lbs. per 1000 sq. ft. (an area 20' by 50') or 8 oz. per 100 sq. ft. (an area 10' by 10').

PRECAUTIONS: Birds, especially waterlowl, feeding or drinking on treated areas may be killed. Apply this product only as specified on this label. Because of the migratory habits of certain waterlowl, do not apply in Nassau County, New York between November 1 and May 20. Do not exceed maximum permitted label rates, since rates above those recommended significantly increase potential hazards to birds and waterlowl. Avoid overlapping treated areas. Where thorough watering is recommended in the directions, or if waterlowl, i.e., ducks or geese, can be expected to enter the treated area after treatment (except spot treatment), water lawn with at least ¼ inch of water immediately after applying this product, however, stop watering before puddling occurs.

Chinchbugs, Earwigs, Ants, Lawn Moths, (Sod Webworms), Bermudagrass Mites, Crickets, Armyworms, Brown Dog Ticks, Chiggers, Clover Mites, Cutworms, Fleas, Leathoppers, Sowbugs, Millipedes, Springtails (Collembola). Application should be made when grass is dry. Water lawn lightly after application. Repeat application in 7-10 days if necessary.

Diffibugs, White grubs of Japanese beetle, European chafer, Southern Chafer, Ataenius spretulus (dung beetle): Water grass thoroughly after application. For billbug control, apply when activity is first observed or when chewed or brown grass indicates damage from these insects. Repeat as necessary.

For grub control, except dung beetle, treat any time between late July and early October. For dung beetle grub control, apply in Spring or early Summer when damage is first noticed. Water grass thoroughly with ¼-½ inch of water after application. Treat again in late Summer or early Fall if a second generation occurs.

HYPERODES WEEVILS

Vater grass thoroughly after application. Treat problem areas in mid-April and again in Mid-May.

NOTE: The spreader settings listed below are approximate. Accurately weigh a few pounds into your spreader and apply to a measured area. Then adjust the spreader setting if necessary in order to assure correct rate of application.

Spreader Make and Model

- -----

For Insects Listed Grass and Dichondra Lawns 5 lbs./1,000 sg. ft. of lawn

BEST AVAILABLE CON.

Central Imperial Mark No. 1662-G	716
Central Suburbia Lawn Spreader No. 1622-0	71/2
Cyclone Model B1 Rotary	6
Greenfield Model 105	8
J.C. Penney Model 6005	81/2
J.C. Penney Rotary Spreader	8
Scott's Precision Flow Spreader PF-1	61/2
Sears Craftsman Model 671 19198	7%