STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal. Open dumping is prohibited. Pesticides that cannot be used or chamically reprocessed should be disposed of in a landfill approved for pesticides or buried in a safe place away from water supplies. Dispose of container in an incinerator or landfill approved for pesticide containers, or bury in a safe place. Consult federal, state or local disposal authorities for approved alternative procedures.

1 🏚

ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish and wildlife. Use with care when applying in areas adjacent to any body of water. Keepout of lakes, streams, and ponds. De not apply when weather conditions favor run-off or drift from target area. Do not contaminate water by cleaning of equipment or disposal of wastes.

CAUTION

Harmful if swallowed. Do not take internally. Avoid eye contact. Flush contaminated eyes with plenty of water and get medical attention it irritation persists. Wash thoroughly after handling. Wash contaminated clothing before re-use. Keep away from food, feedstuffs, and domestic water supplies.

NOTE TO PHYSICIAN: Active Ingredient is a cholinesterase inhibitor. Treat symptomatically. Atropine is an antidote.

Notice of Warranty

C.J. Martin Company makes no warranty of merchantability, filness for any purpose, or otherwise, expressed or implied, concerning this product or its uses which extend beyond the use of the product under normal conditions in accord with the statements made on this label

> E.P.A. Reg. No. 299-200 E.P.A. Est. No. 299-TX-1



Under the Federal Insecticide. Fungicide, and Rodenticide Act. as amended, for the pesuside registered under 299-200 Reg. No. 299-200

Controls: Ants, Chinch Bugs, Sod Webworms (Lawn Moths), Brown Dog Ticks, Crickets, Cutworms, Earwigs, Pillbugs, Certain White Grubs, and Hyperodes (Turfgrass) Weevils Infesting Home Lawns and Other Ornamental Turf Areas.

NET WEIGHT: 50 Pounds

CAUTION: Keep Out Of Reach Of Children. SEE BACK PANEL FOR OTHER CAUTIONS.

ACTIVE INGREDIENT:

Chlorpyrifos [0,0-diethyl 0-(3,5,6-trichloro-2-pyridyl) phosphorothioate]

MarlinCompa

*DURSBAN® - Trademark of The Dow Chemical Company: U.S. Patent No. 3,244,586

606 W.Main St.

Nacogdoches, Texas 75961

的问题,我们就是你们的,我们们还是这些"你们,我们就是**我的**,

99.50%

DIRECTIONS FOR USE

General Classification

シャークの

 $\dot{\mathbf{v}}$

1

It is a violation of federal law to use this product in a manner inconsistent with its labeling,

TO CONTROL: ANTS, CHINCH BUGS, SOD WEB-WORMS (LAWN MOTHS), BROWN DOG TICKS, CRICKETS, CUTWORMS, EARWIGS, GRASSHOPPERS, AND PILLBUGS INFESTING HOME LAWNS AND OTHER ORNAMENTAL TURF AREAS:

Uniformly apply Dursban* Insecticide Granules using any hand or power operated gravity or rotary type spreader designed for application of granular materials. Treat when insects first appear using Dursban* Insecticide Granules at the rate of 200 lbs. per acre. Immediately after application, sprinkle the turf with 1300 gallons of water per acre so as to wash the granules into the zone of insect infestation. Consult the following table to determine amounts of Dursban* insecticide Granules and water needed for smaller sized areas. Re-treat as necessary.

RATES TO USE

Area to be Treated	Pounds of Dursban* Insecticide Granules	Gailons of Water to Sprinkle on Turf After Treatment
1000 Sq. Ft.	5	30
5000 Sq. Ft.	25	150
10000 Sq. Ft.	50	325

TO CONTROL CERTAIN WHITE GRUBS (LARVAE OF EUROPEAN CHAFER AND JAPANESE BEETLE): Use two to four times the amount of Oursban⁺ insecticide Granules shown in the table above. Apply the granules uniformly over the area treated followed immediately by thorough watering to wash the Insecticide into the turf. For best results, treat when grubs are young and actively feeding near the soil surface, usually during late July and August. Consult your local Agricultural Extension Service specialist for best time to treat in your area.

TO CONTROL HYPERODES (TURFGRASS) WEEVIL: Use twice the rate of Dursban* Insecticide Granules shown in the above table. Treat suspected problem areas in mid-April and again in mid-May. Consult your local Agricultural Extension Service specialist for best time to treat in your area.

Kcsp children Sifu polson treated areas until this material is washed into the soll and grass is dry. Keep out of fishpands and other bodies of water. Do not apply where rug-off is likely thooccure +