

SHEEP AND GOATS — LICE, SHEEP TICKS (KEYS) Mix 1 gallon with 150 gallons of water. Apply as a spray or dip, thoroughly wetting animals. Repeat as necessary.

FLEECEWORMS, PSOROPTIC SHEEP SCAB Mix 1 gallon with 150 gallons of water. Apply as a spray or dip, thoroughly wetting animals. Repeat as necessary. For scabies repeat at 7 to 10 day intervals for 2 or more applications as needed.

Use in many plants. Do not apply within 28 days of slaughter.

HOGS — LICE, SARCOPTIC MANGE Mix 1 gallon with 150 gallons of water. Apply as a spray or dip, thoroughly wetting animals. Repeat as necessary. For mange control, repeat at 7 to 10 day intervals for 2 or more applications as needed. Do not apply within 28 days of slaughter.

Apply to spraying and dipping animals during stormy weather. Avoid treating sick or over-treated animals. Feed and water animals before treating. Spraying is recommended for young animals. For young animals must be dipped. Individual hand immersion is essential to avoid sealowing dip solution.

Do not use on dogs, cats or other household pets or on poultry, or in poultry houses. Use only as directed on this label.

Malathion is a cholinesterase inhibitor. Do not use this product simultaneously with other insecticides, pesticides or chemicals having cholinesterase-inhibiting activity nor within a few days before or after treating with any cholinesterase inhibitor.

AGRICULTURAL PREMISES — FLIES, MOSQUITOES Mix 1 quart in 10 gallons of water. Spray walls of barns, sheds, animal shelters, fences, and other farm buildings. Wet surfaces thoroughly but not to the point of excessive runoff. Repeat as needed. Do not use inside dairy barns, milk rooms, or poultry houses. Do not contaminate milk, milking equipment, feed or drinking water.

REPLENISHMENT OF DIPPING VATS Add Stock Tox Plus in accordance with the recommended directions. Add water added to the vat or control replenishment by vats de test.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

WARNING

May be fatal if swallowed. Do not breathe spray mist. Do not get in eyes or on skin or clothing. Wash thoroughly after using.

ANTIDOTE Give one tablespoonful of salt in a glass of warm water to induce vomiting. Flush with fluid. Seek medical attention immediately.

ENVIRONMENTAL HAZARDS

This product is toxic to fish and wildlife. Do not contaminate streams, lakes or ponds. Do not apply where runoff is likely to occur. Do not contaminate water by cleaning of equipment or disposal of wastes.

PHYSICAL OR CHEMICAL HAZARDS

DO NOT USE, POUR, SPILL OR STORE NEAR HEAT OR OPEN FLAME

EPA REG NO 299-209

EPA EST NO 299-TX-1



STOCK TOX PLUS

LIVESTOCK SPRAY OR DIP

Contains Toxaphene and Malathion

CONTROLS: Hornflies, Lice, Ticks, and Scabies (Psoroptic, Sarcoptic, and Chorioptic Mange).

WARNING

KEEP OUT OF REACH OF CHILDREN
SEE BACK PANEL FOR OTHER WARNINGS.

ACTIVE INGREDIENTS:

Toxaphene (Technical chlorinated camphene containing 67-69% Chlorine)	43.00%
Malathion (0,0-dimethyl dithiophosphate of diethyl mercaptosuccinate)	4.30%
Petroleum Distillates	46.27%
INACTIVE INGREDIENTS:	6.43%

CONTENTS: 1 GALLON

ACCEPTED
SEP 1 1981
Fungicide, Insecticide, and Rodenticide Act
as amended, for the pesticide registered under EPA Reg. No. 299-209

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

STORAGE & DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

PESTICIDE DISPOSAL

Spray mixture or rinseate that is left over must be disposed of according to label instructions. Do not use for any other purpose. Must be disposed of according to Federal, State or Local regulations under the Resource Conservation and Recovery Act.

CONTAINER DISPOSAL

Rinse (or equivalent) and dispose of in a sanitary landfill or by other means in accordance with State and Local procedures.

BEEF CATTLE — HORNFLIES, LICE, TICKS, SCABIES (PSOROPTIC, SARCOPTIC, AND CHORIOPTIC MANGE), EAR TICKS Accurately mix 1 gallon with 150 gallons of water. Apply as a dip or spray, thoroughly wetting animals. For ear ticks, apply spray into each ear and to head and neck. Repeat as necessary at 7 to 10 day intervals for 2 or more applications as needed. Do not use on lactating dairy animals. Do not apply to lactating dairy animals.

BACK RUBBERS To aid in controlling hornflies and lice, mix 1 gallon with 150 gallons of water. Charge burrap roll with solution so it is wet with solution. Repeat charging as necessary. Do not apply to dairy animals within 28 days of slaughter.

606 W. Main St.

C. J. Martin Company

Nacogdoches, Texas 75



STOCK TOX PLUS

LIVESTOCK SPRAY OR DIP

Contains Toxaphene and Malathion

CONTROLS: Hornflies, Lice, Ticks, and Scabies (Psoroptic, Sarcoptic, and Chorioptic Mange).

WARNING

KEEP OUT OF REACH OF CHILDREN
SEE BACK PANEL FOR OTHER WARNINGS.

ACTIVE INGREDIENTS:

Toxaphene (Technical chlorinated camphene containing 67-69% Chlorine)	43.00%
Malathion (0,0-dimethyl dithiophosphate of diethyl mercaptosuccinate)	4.30%
Petroleum Distillates	46.27%
INACTIVE INGREDIENTS:	6.43%

CONTENTS: 1 GALLON

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

STORAGE & DISPOSAL


Do not contaminate water, food, or feed by storage or disposal.

PESTICIDE DISPOSAL — Pesticide spray mixture or rinsate that cannot be used according to label instructions must be disposed of according to Federal, State or Local procedures under the Resource Conservation and Recovery Act.

CONTAINER DISPOSAL — Triple rinse (or equivalent) and dispose of in a sanitary landfill or by other approved State and Local procedures.

BEEF CATTLE — HORNFLEES, LICE, TICKS, SCABIES (PSOROPTIC, SARCOPTIC AND CHORIOPTIC MANGE), EAR TICKS: Accurately mix 1 gallon with 150 gallons of water. Apply as a dip or spray, thoroughly wetting animals. For ear ticks, use a mist sprayer and apply 3, 4, into each ear and to head and neck. Repeat as necessary. For scabies, repeat at 7 to 10 day intervals for 2 or more applications as needed. Do not apply within 28 days of slaughter. Do not apply to lactating dairy animals.

BACK RUBBERS: To aid in controlling horn flies and lice, mix 1 quart with 3 gallons of spray or dip solution. Charge butler roll with solution so it is wet but not to the point of dripping. Repeat charging as necessary. Do not apply to dairy animals. Do not apply within 28 days of slaughter.



ation with 150 gallons of water as necessary.
gallons of water. Apply For scabies repeat at 7 to 10 day intervals for 2 or more applications as needed. Do not apply within 28 days of slaughter.
one of water. Apply as a range control, repeat at 7 to 10 day intervals for 2 or more applications as needed. Do not apply within 28 days of slaughter.
Avoid treating sick or overworked animals. Not recommended for young calves. Caution is essential to avoid injury to animals.
Do not use on poultry houses. Use only on clean, dry surfaces.
Do not mix with other pesticides. Do not use within 28 days of slaughter.
Do not use on animals with 10 gallons of water. Wet surfaces. Do not use inside dairy buildings. Do not use on milking equipment. Do not use on animals with open wounds.
Do not use on animals with open wounds.
Do not use on animals with open wounds.
Do not use on animals with open wounds.

W. Main St. *C. J. Martin Company* Nacogdoches, Texas 75961