OF PESTICIDES PROGRAM STRATION DRYISION (78-767) WASHINGTON, DC 20460

NOTICE OF PESTICIDE: REGISTRATION REPESISTRATION

(Under the Federal Insecticide, Fundicide, and Rodenticide Act, as amended)

TUNE 27, 1990

Until

NAME OF PESTICIDE PRODUCT

Martin's Termite & Soil Insect Control

NAME AND ADDRESS OF REGISTRANT (Include EIP code)

C.J. Martin Company P.O. Box 631090 606 West Main Street Nacogdoches, TX 75963

NOTE: Changes in labeling formula differing in aubstance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In at a correspondence on this product always refer to the above U.S. EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby Registered/Reregistered under the Federal Insecticide, Fungicide, and Rodenticide Act.

A copy of the labeling accepted in connection with this Registration/Reregistration is returned herewith.

Registration is in no way to be construed as an indorsement or approval of this product by this Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is conditionally registered in accordance with FIFRA section 3(c)(7)(A) provided that you:

- 1. Submit/cite all data required for registration/reregistration of your product under FIFRA section 3(c)(5) when the Agency requires all registrants of similar products to submit such data.
- Add the phrase "EPA Registration No. 299-223" to your label before you release the product for shipment.
- Submit five (5) copies of your final printed labeling store you release the product for shipment. Refer to the A-79 enclosure for a further description of final printed labeling.
- If these conditions are not complied with, the requestration will be subject to cancellation in accordance with FIFRA section o(e). Your release for amplified of the product constitutes acceptance of these conditions. Astamped copy of the label is enclosed for your records.

BEST AVAILABLE COPY

dennis d. Edwards, Jr. Product Manager (12) Insecticide-Rodenticide Branch Registration Division (H7505C)

Enclosures

ATTACHMENT IS APPLICABLE

SIGNATURE OF APPROVING OFFICIAL

6/27/90

EPA Form \$570-6 (Rev. 5-76)

MARTIN'S

TERMITE & soil insect control

with DURSBAN Insecticide

CONTROLS:

Termites

Wood Destroving Pests

Carpenter Ants

Armyworms

Fire Ants

Chinch Buds

Sed Webwerms

Ticks

KEEP OUT OF THE REACH OF CHILDREN WARNING (See Fack famel for Other Cautions)

ACTIVE INGREDIENT:

Chlorovrifos [0,0-diethyl 0-(3,5,6-trichloro-2-pyridinyl)

INERT INGREDIENTS:....

87.4%

TOTAL

100.00%

* DURSBAN - Trademark of The Dow Chemical Company

EPA Reg. No. 299-___

EPA Est. No. 299-TX-1

Net Weight _____

Manufactured by:

U.J. Martin Company 606 W. Main St.

Nacoudoches, IX 75963-1089

ACCEPTET/ with CONNEY

JUN 2 7 1990

Under the Poderal Inc Pungicide, and Redenticide Act a amended, for the pa red under EPA Rog. No.

3.112

「我選問会の主に、 STATEMENTS 「我ならばない」、 かい ディーのはない。 データ Lindual ディー・ディー 一手的ながみによって

which the causes eve innitation, manuful it swallowed. Do not det in eves, on skin or clothing. Avoid breathing vapors and solay mist, mandle concentrate in ventilated area. Wash thoroughly after handling and before eating or smoking. Do not wear contaminated clothing or shoes. Reep away from foud, readstorfs, and domestic water supplies. Reep children and bets out or freated areas until these areas are dry.

STATEMENT OF PRACTICAL TREATMENT- IF SWALLOWED: Do NOT make person vomit. Call a doctor immediately. If ON SKIN: In case of contact, remove contaminated clothing and immediately wash skin with soap and water. If IN EYES: Flush eyes with plenty of water. Call physician and bet prompt medical attention.

NOTE TO PHYSICIANS: This product contains an aromatic petroleum solvent and may present an aspiration problem. Chlorpyrifos is a cholinesterase inhibitor. If signs and symptoms of cholinesterase inhibition develop, atropine by injection is antidotal; 2-PAM may also be given in conjunction with atropine, but should never be used alone.

ENVIRONMENTAL HAZARDS: This product is toxic to fish, birds and other wildlife. Bo not apply directly to water. Do not contaminate water when disposing of equipment washwaters.

PHYBICAL OR CHEMICAL HAZARDS: Do not use or stone near heat or open flame.

DIRECTIONS FOR USE

BEST AVAILABLE COPY

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WATH LABEL PRECAUTIONARY STATEMENTS AND DIRECTIONS.

DIRECTIONS:

DENERAL: Nantin's TERMITE & soil insect control is particularly useful for prevention and control of ants, termites, and certain wood destroying insects. For termites, the chemical solution should be adequately dispersed in the soil to provide a uniform barrier between the wood in the structure and termite colonies in the soil. It can also by used for treatment of localized areas of wooden structures, the soil under firewood piles, and stored lumber to control wood intesting borers. Carpenter Ants and Carpenter Bees, in addition, this product controls many common bests in lawns and other outdoor areas around the home. Make new dilution for each use. NOTE 2 TABLESPOONS EQUALS 1 FL. 02.

JK ~

SHAKE WELL BEFORE USING

FOR TERMITES (Outdoors): Remove all wood debris and wood forms. Determine the extent of termite infestation. For prevention and light infestation, treat soil adjacent to both sides of infested foundations of poured concrete, pipes or piers by digding a trench 6 inches wide and 4 inches deep (or down to the top of the footing whichever is less). For brick and block foundations dig the trench about 12 inches deep. It tooting is More than 12inches deep, make holes with a crowbar, pipe or a rod about one foot apart that extend from the trench bottom to the top of the Make holes closer in hard-packed clay soils. permit trenching, surface application will not conditions: adjacent to interior foundation walls may be made but the treated strip should not exceed 18 inches in width. solution with a sprinkling can. Mix 5 1/3 fl. oz. per gallon of water and treat 2 1/2 linear feet of trench or foundation. Mix solution into soil as it is replaced in the trench. Be sure to treat the interiors or accessible hollow block foundations at same rate as above.

USE PRECAUTIONS

NEVER DIG BELOW THE TOP OF THE FOOTING. Avoid contamination of public and private water supplies by following these precautions. Do not allow a hose or any type of faucet extension to reach into the termite chemical solution while filling or mixing this material with water. Use antibackflow or anti-siphonage equipment on all filling equipment. These devices at hardware or plumbing supply stores. perenda antibackflow equipment will help insure that the user of this product will not contaminate domestic water supplies. Refer to FHA (Federal Housing Administration), state, and local specifications for safe distances of treatment areas from wells. Soil in the vicinity of wells, cisterns or ponds should not be Do not treat soil under or adjacent to structures that cisterns or wells. Be careful to have' adequate contain ventilation when treating areas under structure. 'For' severa infestations or treatment under slabs, in basements and buildings with insufficient crawl space, etc., contact a licensed post control ageretor or termite control service.

RETREATMENT RESTRICTIONS

BEST AVAILABLE COPY

Retreatment for subterranean termites should only be made when there is evidence of reinfestations subsequent to the initial treatment, or there has been a disruption of the chemical barrier in the soil due to construction, excavation, landscaping, etc. Reapplication should be made as a spot treatment to those areas.

Annual retreatment of the entire premises must be avoided.

FENCE POSTS-TERMITE PREVENTION: Mix 5.3 azs per dailon of water or 2 1/2 tablespoons per quart. Apply one gallon of diluted Martin's TERMITE & soil insect control per post hole. Pour one

BEST AVAILABLE COPY

half of solution to the soil when filling the hole and add the remaining 1/2 gallon diluted solution to the soil when filling the hole so that all the fill soil is treated to form a protective barrier.

NOTE: Do not treat soi: that is water saturated or frozen.

TREATMENT FOR WOOD DESTROYING PESTS (Outdoors): (Powder Post Beetles, Furniture Beetles, Deathwatch Beetles, Round-headed House Borers, Carpenter Ants, Carpenter Bees). Treat localized areas of infested wood. With a paint brush, use 2 1/2 TBS (Tablesboons) per duant of water. Apply emulsion evenly on wood surfaces. Allow to dry and repeat treatment to ensure penetration.

Before stacking firewood or lumber, apply 5 1/3 fl. oz. per qallon of water as a localized spray to surfaces immediately below such materials. Treating soil around firewood may also be helpful in reducing pests immigrating from surrounding areas. Do not make direct application to firewood. Do not use in structures occupied by animals, to be used for food purposes or which produce products for human consumption.

HOME INVADING PESTS (Outside Dwellings): Ants, Fleas, Ticks, Earwigs, Crickets, Sowbugs (Pillbugs), Millipedes — For best results, apply with a sprayer. Use 1-1/2 fl. oz. in 15 gallons to treat 500 sq. ft. NOTE: One half Taplespoon in 2 gallons of water covers 66 sq. ft. To help prevent infestations of buildings, treat a band of soil 6-10 feet wide around and adjacent to the building, also the building's foundation to a height of 2-3 feet when pests are active and may find entrance. Do not treat bets with this product. Do not use inside dwellings.

IMPORTED FIRE ANT MOUNDS: Mix 2 TBS (Tablespoonfuls) per gallon of water to each individual mound, using a sprinkler can. Thoroughly wet mound and treat a 4 foot diameter circle around the mound. For best results, apply in cool weather or in early morning or late afternoon. Treat new mounds as they appear.

LAWN PESTS: Armyworm, Lawn Moth Larvae (Sod Webworms), Dichondra Flea Beatles. Chinchbuds, Curworms, European Crane Fly (Larvae), Unasshuppers, Unickets, Chiqqens, Ulover Mites, Earwigs, Fleas-For best results apply with a sprayer, mix 1-1/2 fl. oz. in 15 dallons of water to treat 500 sq. ft. of lawn.

Mow lawn and water well 1-2 days before treatment. Refrain from mowing or watering treated surface for two days following application. Spray when pests first appear. Treat entire lawn. Do not attempt to control lawn pests by spot spraying. Repeat treatments may be necessary at 4 to 6 week intervals.

WHISE GRUBS IN LAWNS: (Larvae of European Chafer and Japanese Beetle) Mix 6-1/2 fl. oz. in 15 dallons of water to treat 500 sq. ft. For best results use a sprayer. Treat entire lawn. Do not attempt to control lawn pests by spot spraying. For best

results, water lawn thoroughly (1 to 2 inches) immediately following application to wash the insecticide into the soil. Treat lawn when grubs are young and actively feeding near the soil surface, usually during late duly and August. Consult your local Agricultural Extension behavior Specialist for proper time to treat in your area.

NOTE: Mosquitoes coming to rest on areas treated for the control of lawn and home invading pests will be killed for varying periods of time after treatment, depending on exposure of treated areas to weathering conditions. NOTE: do not use on vegetables or food crops. Do not use on azaleas, camellias, poinsettias, rose bushes or varigated ivy because of possible injury to these plants.

STORAGE AND DISPOSAL

STORAGE: Do not contaminate water, food, or feed by storage or disposal. Keep pesticide in original container. Do not put concentrate or dilute into food or drink containers. Avoid contamination of feed, foodstuffs and domestic water supplies. Store in a cool, dry place, preferably in a locked storage area. Do not store diluted spray. Store above freezing. Handle concentrate in a ventilated area. Keep container closed. DISPOSAL: FRODUCT- Partially filled bottle may be disposed of by securely wrapping container in several layers of newspaper and discard in trash. CONTAINER- Do not reuse empty bottle. Rinse thoroughly before discarding in trash.

NOTICE: Buyer assumes all responsibility for safety and use not in accordance with directions.

