

PM 31 303-209

10/20

# ASCEND

## GERMICIDAL DETERGENT

Disinfectant · Deodorizer · Fungicide  
\*Virucide · Detergent

Active Ingredients:  
Didecyl dimethyl ammonium chloride . . . . . 9.22%  
n-Alkyl (C<sub>14</sub> 50%, C<sub>17</sub> 40%, C<sub>18</sub> 10%)  
dimethyl benzyl ammonium chloride . . . . . 6.14%  
Inert Ingredients: . . . . . 84.64%

**KEEP OUT OF REACH OF CHILDREN  
DANGER**

### STATEMENT OF PRACTICAL TREATMENT

In case of contact, immediately flush eyes or skin with plenty of water for at least 15 minutes. For eyes, call a physician. If swallowed, drink promptly a large quantity of water. Avoid alcohol. Get medical attention.

**NOTE TO PHYSICIAN:** Probable mucosal damage may contraindicate the use of gastric lavage. Measures against circulatory shock, respiratory depression and convulsion may be needed.

### PRECAUTIONARY STATEMENTS

#### HAZARDS TO HUMANS AND DOMESTIC ANIMALS

##### DANGER

**CORROSIVE.** Causes eye damage and severe skin irritation. Do not get in eyes, on skin, or on clothing. To protect eyes wear goggles or face shield and to protect skin, wear rubber gloves when handling. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse. Harmful if swallowed.

##### ENVIRONMENTAL HAZARDS

Do not discharge effluent containing this product into lakes, streams, ponds, estuaries, oceans, or other waters unless in accordance with the requirements of a National Pollutant Discharge Elimination System (NPDES) permit and the permitting authority has been notified in writing prior to discharge. Do not discharge effluent containing this product to sewer systems without previously notifying the local sewage treatment plant authority. For guidance contact your State Water Board or Regional Office of the EPA

#### Net Contents:

EPA Reg. No. 303-209 EPA Est. No. 303-IN-1

Huntington Laboratories, Inc. · Huntington, IN 46750 · (219) 356-8100  
Lansdale, PA 19446 · Denver, CO 80205 · Dallas TX, 75227  
Oakland, CA 94621 · Bramalea, Ontario, Canada L6T 1E3

#### (Directions for Use continued)

\*Kills HIV-1 (AIDS virus) on pre-cleaned, environmental surfaces/objects previously soiled with blood/body fluids in healthcare settings or other settings in which there is an expected likelihood of soiling of inanimate surfaces/objects with blood/body fluids, and in which the surfaces/objects likely to be soiled with blood/body fluids can be associated with the potential for transmission of Human Immunodeficiency Virus Type 1 (HIV-1) (associated with AIDS)

#### SPECIAL INSTRUCTIONS FOR CLEANING AND DECONTAMINATION AGAINST HIV-1 (AIDS VIRUS) OF SURFACES/OBJECTS SOILED WITH BLOOD/BODY FLUIDS:

**Personal Protection:** Disposable latex or vinyl gloves, gowns, face masks, or eye coverings as appropriate must be worn during all cleaning of blood/body fluids and during decontamination procedures

**Cleaning Procedures:** Blood/body fluids must be thoroughly cleaned from surfaces/objects before application of disinfectant

**Contact Time:** HIV-1 (AIDS virus) is inactivated after a contact time of 4 minutes at 25°C (room temperature). Use a 10-minute contact time for other viruses, fungi, and bacteria listed

**Disposal of Infectious Materials:** Blood/body fluids should be autoclaved and disposed of according to federal, state, and local regulations for infectious waste disposal

### STORAGE AND DISPOSAL

#### PROHIBITIONS

Do not contaminate water, food, or feed by storage or disposal. Open dumping is prohibited. Do not reuse empty container.

#### PESTICIDE DISPOSAL

Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

#### CONTAINER DISPOSAL

Triple rinse (or equivalent), then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or incinerate, or, if allowed by state and local authorities, burn. If burned, stay out of smoke.

Ascend is a multi-purpose germicidal detergent and deodorant effective when diluted 1:256 in up to 400 ppm hard water (calculated as CaCO<sub>3</sub>) plus 5% organic serum according to the AOAC Use-dilution and virucidal qualification tests. Disinfects, cleans, and deodorizes in one labor saving step. Recommended for use in hospitals, nursing homes, schools and colleges, veterinary clinics and animal life science laboratories, federally inspected meat and poultry establishments, poultry farms, turkey farms, dairy farms, hog farms, and breeding establishments.

Effective against the following pathogens on hard, non-porous, inanimate surfaces when diluted 1:256 (1/2 oz. to 1 gallon):

#### Bactericidal against:

- Pseudomonas aeruginosa*<sup>1</sup>
- Staphylococcus aureus*<sup>1</sup>
- Salmonella choleraesuis*
- Acinetobacter calcoaceticus*
- Bordetella bronchiseptica*
- Chlamydia psittaci*
- Enterobacter aerogenes*
- Enterobacter cloacae*
- Escherichia coli*<sup>1</sup>
- Fusobacterium necrophorum*
- Klebsiella pneumoniae*<sup>1</sup>
- Listeria monocytogenes*
- Pasteurella multocida*
- Proteus mirabilis*
- Proteus vulgaris*
- Salmonella enteritidis*
- Salmonella typhi*
- Salmonella typhimurium*
- Serratia marcescens*
- Shigella flexneri*
- Shigella sonnei*
- Staphylococcus aureus*,  
(methicillin resistant)
- Staphylococcus epidermidis*<sup>1</sup>
- Streptococcus faecalis*<sup>1</sup>
- Streptococcus pyogenes*

#### Virucidal against:

- \*Adenovirus type 4
- \*Canine distemper
- \*Canine parvovirus
- \*Feline leukemia
- \*Feline panleukopenia
- \*Feline picornavirus
- \*Herpes simplex type 1
- \*Herpes simplex type 2
- \*HIV-1 (AIDS virus)
- \*Infectious bovine rhinotracheitis
- \*Infectious bronchitis (Avian IBV)
- \*Influenza A/Hong Kong
- \*Porcine parvovirus
- \*Pseudorabies
- \*Rabies
- \*Respiratory syncytial virus (RSV)
- \*Rubella (German Measles)
- \*Transmissible gastroenteritis virus (TGE)
- \*Vaccinia

#### Fungicidal against:

- Aspergillus niger*
- Candida albicans*
- Trichophyton mentagrophytes*

<sup>1</sup> ATCC & antibiotic-resistant strain  
<sup>2</sup> antibiotic-resistant strain only

## BEST AVAILABLE COPY

DILUTION: 1 to 256

1/2 OUNCE PER GALLON

#### DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

This product is not to be used as a terminal sterilant/high level disinfectant on any surface or instrument that (1) is introduced directly into the human body, either into or in contact with the bloodstream or normally sterile areas of the body, or (2) contacts intact mucous membranes but which does not ordinarily penetrate the blood barrier or otherwise enter normally sterile areas of the body. This product may be used to pre-clean or decontaminate critical or semi-critical devices prior to sterilization or high-level disinfection.

#### GENERAL USE DIRECTIONS

For use on hard, non porous surfaces such as floors, walls, metal surfaces, stainless steel surfaces, glazed ceramic tile, glazed porcelain, rubber and plastic surfaces (such as polypropylene, polystyrene, etc.) and other hard, non-porous surfaces.

Remove gross filth and heavy soil deposits then thoroughly wet surfaces. Use 1/2 ounce per gallon of water for a minimum contact time of 10 minutes. Can be applied with a cloth, mop, sponge, coarse spray or by soaking. The recommended use solution is prepared fresh for each use then discarded. Rinsing is not necessary unless used on floors that are to be waxed or polished.

USDA: For use in federally inspected meat and poultry plants on all hard, non-porous surfaces in inedible product processing areas, nonprocessing areas, and/or exterior areas.

#### MILDEWSTATIC INSTRUCTIONS:

Will effectively control the growth of mold and mildew plus the odors caused by them, when applied to hard, non porous surfaces such as walls, floors, and table tops. Apply solution (1/2 ounce per gallon of water) with a cloth, mop, sponge, or coarse spray. Make sure to wet all surfaces completely. Let air dry. Repeat application weekly or when growth reappears.

DEC 15 1988  
303-209

(label continued)

**ASCEND  
GERMICIDAL DETERGENT**

**FARM PREMISE, LIVESTOCK, POULTRY & TURKEY HOUSE DISINFECTANT**

DILUTION: 1:256 ..... 1/2 OUNCE PER GALLON OF WATER

**DIRECTION FOR USE**

1. Remove all animals and feeds from premises, trucks, coops, crates, and enclosures.
2. Remove all litter and manure from floors, walls, and surfaces of barns, pens, stalls, chutes, vehicles, and other facilities and fixtures occupied or traversed by animals.
3. Empty all troughs, racks, and other feeding and watering appliances.
4. Thoroughly clean all surfaces with soap or detergent, and rinse with water.
5. Saturate all surfaces with the recommended disinfecting solution for a period of 10 minutes.
6. Immerse all halters, ropes, and other types of equipment used in handling and restraining animals, as well as forks, shovels, and scrapers used for removing litter and manure.
7. Ventilate buildings, coops, cars, boats, and other closed spaces. Do not house animals or employ equipment until treatment has been absorbed, set, or dried.
8. After treatment with disinfectant, thoroughly scrub feed racks, troughs, automatic feeders, fountains, and waterers with soap or detergent, and rinse with potable water before reuse.

[May be used as a box insert and/or hang tag.]

010495

**ACCEPTED**

FEB 15 1995

Under the Federal Insecticide, Fungicide, and  
Rodenticide Act, as amended, for the  
pesticide registered under  
EPA Reg. No. 303-267