

**DIRECTIONS FOR USE
GENERAL INFORMATION**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Do not apply when weather conditions favor runoff or drift from the target area.

is a pre-emergence herbicide which is applied to established turfgrasses for the control of annual bluegrass (Poa annual), smooth and hairy crabgrass, goosegrass, annual crabgrass or crowfoot barnyardgrass (watergrass) and green and yellow foxtail. Controls these annual weeds by killing their seeds as they germinate. Apply before these annual weeds begin to germinate since it does not control established weeds.

Application Directions

Rate, Frequency and Timing of Application
Successful pre-emergence control of the annual grass weeds listed on this label requires proper timing of application.

Apply approximately one to two weeks prior to the germination of annual grass weeds. At summer annual grasses do not germinate at the same time. If application timing does not coincide with the normal germination period of any of the annual grass weeds listed on this label, weed control results may be erratic or poor. Optimum weed control performance will be obtained if treated areas are irrigated soon after application.

Any cultural practices that disturb the soil such as aeration or verticuring should be done prior to application.

1. Spring germinating summer annual grasses—crabgrasses, foxtails and barnyardgrass.

A. Northern (cool season) turfgrasses
For pre-emergence control, apply **1 1/2 gallons per acre or 4oz per 1,000 sq. ft.** in the spring one to two weeks prior to the onset of conditions favorable for annual weed grass seed germination.

B. Southern (warm season) turfgrasses
For pre-emergence control, apply **2 gallons per acre or 6oz per 1,000 sq. ft.** in the late winter or early spring one to two weeks prior to the onset of conditions favorable for annual weed grass seed germination.

In areas of the United States where germination of these annual weed grass species continues for extended periods of time, a second application at the recommended rate may be made 8 to 10 weeks after the first treatment for continued control.

2. Late spring/early summer germinating annual grasses—goosegrass. Although classified as a summer annual grass, goosegrass normally germinates 4 to 6 weeks later than crabgrass, foxtails and other summer annual grass species. Applications should therefore be timed accordingly.

A. Northern (cool season) turfgrasses
For pre-emergence control, apply **1 1/2 gallons per acre or 4oz per 1,000 sq. ft.** in the late spring or early summer one to two weeks prior to the expected germination of goosegrass.

When applied as directed the following established turfgrasses are tolerant:
Kentucky bluegrass, Zoysia grass, Perennial ryegrass, Bermudagrass, Centipedegrass, St. Augustine grass, Fescue, Bahiagrass.

and will not harm most nearby established ornamental trees and shrubs.

B. Southern (warm season) turfgrasses
For pre-emergence control, apply **2 gallons per acre or 6oz per 1,000 sq. ft.** in the late spring or early summer one to two weeks prior to the expected germination of goosegrass.

3. Northern and southern turfgrasses
In areas of the United States where pre-emergence control of both early germinating annual grasses (crabgrass, foxtails and barnyardgrass) and the late germinating goosegrass is desired, two applications at the recommended rate spaced 8 to 10 weeks apart may be made to provide for season control of these annual grass species.


4. Poa annual (annual bluegrass)
In areas of heavy infestation of annual bluegrass, its germination will temporarily result in thin turfgrass areas. Proper fertilization, irrigation and soil aeration and reseeding will encourage existing desirable turfgrasses and newly planted seedling turfgrasses to fill in these thin areas previously occupied by annual bluegrass.

A. Northern (cool season) turfgrasses
For pre-emergence control, apply **1 1/2 gallons per acre or 4oz per 1,000 sq. ft.** in the late spring or early summer one to two weeks prior to the expected germination of annual bluegrass.

B. Southern (warm season) turfgrasses
For pre-emergence control, apply **2 gallons per acre or 6oz per 1,000 sq. ft.** in the late spring or early summer one to two weeks prior to the expected germination of annual bluegrass. In southern areas of the United States where germination of annual bluegrass continues for extended periods of time, a second application at **2 gallons per acre or 6oz per 1,000 sq. ft.** can be made 10 to 12 weeks after the first application.

TURF LEADER CRABGRASS HERBICIDE CONCENTRATE

Huntington




ACTIVE INGREDIENT
N-butyl-N-ethyl-a, a-trifluoro-2, 6-dinitro-p-toluidine 16.32
INERT INGREDIENTS 83.72
TOTAL 100%

Contains 0.5 lbs. Benflin per gallon. Benflin is a product of Elanco Products Company, a division of Eli Lilly and Company.

- Kills Crabgrass and other weeds listed as they germinate
- Feeds the lawn
- Contains Balan®

**KEEP OUT OF REACH OF CHILDREN
DANGER**

(SEE SIDE PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS)



BEST DOCUMENT AVAILABLE

When applying this product to established turfgrasses, it is essential that proper irrigation and fertilization be followed. For satisfactory results, irrigation should be increased and equipment designed to place seed in contact with the soil (such as the Baggers Aero Seeder) should be employed. When using the highest recommended rate, irrigation should be delayed 12 to 16 weeks after application.

Precautions
Apply to the turf prior to the germination of annual bluegrass (Poa annual) turfs and fine leaved fescues. Do not apply in the spring to turfgrasses planted the previous fall in bermudagrass areas that have been overseeded with winter grasses. Because of varying cultural practices, should not be used on putting greens. Should not be used on dicyandra dactyloctenium grass or centropogon grass. These turfs are well established.

**HAZARD TO HUMANS & DOMESTIC ANIMALS
DANGER**

Corrosive. Causes eye damage. Do not get in eyes. Wear goggles or face shield when handling concentrate. If splashed in eyes, flush thoroughly with water. Harmful if swallowed or absorbed through the skin. Avoid contact with skin, eyes or clothing. Wash thoroughly after handling.

IF SWALLOWED: Call a physician or Poison Control Center. Drink 1 or 2 glasses of water and induce vomiting by touching back of throat with finger. Do not induce vomiting or give anything by mouth to an unconscious person.

IF IN EYES: Immediately flush with large amounts of water and contact a physician.

IF ON SKIN: Wash exposed areas with soap and water. Wash all contaminated clothes before reuse. Contact a physician if irritation develops.

IF INHALED: If discomfort occurs, move to fresh air. If breathing difficulty occurs, start mouth-to-mouth artificial respiration and contact a physician.

ENVIRONMENTAL HAZARD

This pesticide is toxic to fish. Do not apply directly to water. Do not contaminate water by cleaning of equipment or disposal of wastes.

STORAGE AND DISPOSAL

STORAGE: Do not contaminate water, food or feed by storage, disposal or cleaning of equipment. Open dumping is prohibited. Keep in a dry place. **AVOID FREEZING. STORE ABOVE 40°F. IF FROZEN, POUR MELT. CONTROL MAY RESULT. DO NOT STORE NEAR HEAT OR FLAME.**

PESTICIDE DISPOSAL: Pesticide, spray mixture or residue that cannot be used according to label instructions must be disposed of according to Federal, State or local procedures under the Resource Conservation and Recovery Act.

CONTAINER DISPOSAL: Triple rinse (or equivalent). Then offer for recycling or other reuse, if appropriate, and dispose of in a sanitary land fill or other controlled facility approved by state and local authorities.

NOTICE: The seller makes no warranty, expressed or implied, concerning the use of this product other than that called on the label. Buyer assumes all risks of use and/or handling of this material when such use and/or handling is contrary to label instructions.

EPA Reg. No. 303-

EPA Est. No. 303-IN-1

Huntington Laboratories, Inc., Huntington, IN 46780
Lansdale, PA 19448 • Dallas, TX 75227
Tulsa, OK 74106 • Birmingham, AL 35202