ACCEPTED

Jun 5, 1970

THE FEDERAL INSECTIONS
THE FEDERAL INSECTIONS
FOR AND RODENTH INF. FOR
FOR AND RODENTH INF. FOR
FOR AND SON AFCISTERFOR AND SON AFCISTERS
THE FEDERAL INSECTIONS
THE FEDERAL INSECTIO

NET WEIGHT: 20 LBS.

Covers 10,000 Square Feet



### GRANULAR

## LAWN FUNGICIDE

For control of BROWN PATCH, COPPER, DOLLAR & LEAF 570T, SNOW MOLD.

Apply with Spreader

### ACTIVE INGREDIENTS

Thiram		5.00 %
(Tetramethylthiuram	Disulfide)	
*Cadmium Chloride		.75%
INERT INGREDIENTS		94.25 %
*Total Cadmium		.48 %

For the prevention and control of Copper Spot, Dollar Spot, Leaf Spot (Helminthosporium-Curvalarial), Brown Patch and Snow Mold on Lawns containing Kentucky Blue including Merion and other strains, the various Fescues, Bents and most other gasses.

### SPECIAL INSTRUCTIONS

Do not water or mow lawn area for 24 hours after application. Application will be more effective if made in the early morning hours when dew is on the grass. Repeat treatments are necessary until fungus diseases disappear. Apply contents of this bag to 10,000 square feet for prevention program and at double rate (5000 square feet) if fungus diseases have already infested lawn area.

### SPRING PROGRAM

Application should be made immediately at a double rate (contents of this bag to 5000 square feet) if no disease treatments have been given and Dollar or Leaf Spot have invaded lawn area. For a prevention program against these diseases apply RUSSELL'S GRANULAR LAWN FUNGICIDE soon after the second mowing in the Spring and repeat every 7 to 10 days for a minimum of five times. This program will prevent Fungus activity of Dollar Spot and Leaf Spot which may kill good grasses.

### SUMMER APPLICATION

Apply immediately at double rate if Brown Patch or Curculvaria Disease is active in lawn area. Prevention of these diseases may be obtained by an application of RUSSELL'S GRANULAR LAWN FUNGICIDE every 7 to 10 days during the hot humid months.

### EARLY FALL APPLICATION

Prevent infestation of Dollar Spot and Copper Spot with treatments at 7 to 10 day intervals during the period of cool nights.

#### SNOW MOLD

Apply at double rate in the late Fall, before the first snowfall and repeat again at double rate in January.

### **CAUTION**

to 10,000 square feet for prevention program and at double rate (5000 square feet) if fungus diseases have already infested lawn area.

### SPRING PROGRAM

Application should be made immediately at a double rate (contents of this bag to 5000 square feet) if no disease treatments have been given and Dollar or Leaf Spot have invaded lawn area. For a prevention program against these diseases apply RUSSELL'S GRANULAR LAWN FUNGICIDE soon after the second mowing in the Spring and repeat every 7 to 10 days for a minimum of five times. This program will prevent Fungus activity of Dollar Spot and Leaf Spot which may kill good grasses.

### SUMMER APPLICATION

Apply immediately at double rate if Brown Patch or Curculvaria Disease is active in lawn area. Prevention of these diseases may be obtained by an application of RUSSELL'S GRANULAR LAWN FUNGICIDE every 7 to 10 days during the hot humid months.

### EARLY FALL APPLICATION

Prevent infestation of Dollar Spot and Copper Spot with treatments at 7 to 10 day intervals during the period of cool nights.

### SNOW MOLD

Apply at double rate in the late Fall, before the first snowfall and repeat again at double rate in January.

# CAUTION KEEP OUT OF REACH OF CHILDREN

Avoid contact with eyes, skin or clothing. In case of contact wash exposed areas thoroughly with water to avoid irritation. Avoid breathing dust since inhalation may prove harmful. Store away from feed and foodstuffs.

This product is toxic to fish and wildlife. Birds feeding on treated areas may be killed. Keep out of any body of water. Do not contaminate water by cleaning of equipment, or disposal of wastes.

### INTERNAL

If swallowed give raw white of egg. Induce vomiting with mixture of salt, soap or mustard in warm water. Call physician at once.

NOTE: We make no warranty of any kind, expressed or implied, concerning the suitability of this product. Buyer, user or handler assumes all risks whether or not directions are followed.

Spreader Settings for Proper Application—
Normal Rate of 20 pounds per 10,000 square feet.
Scotts 5
Lawn Beauty 5
Jackson — 20" 30
Buch #2 — 20" 3/4
Cyclone 3.4

Others 3/16 of an inch opening
Where double rate application is wanted, go over area twice at above settings.

Distributed by

Gerald S. Russell

25-16 50th STREET

WOODSIDE, LONG ISLAND, N. Y. 11377

USDA Reg. No. 318-39