

PRECAUTIONARY STATEMENTS

HAZARD TO HUMANS (& DOMESTIC ANIMALS)

CAUTION

Harmful if swallowed. May cause irritation of eyes, nose, throat and skin. Avoid contact with skin, eyes or clothing. Wash thoroughly after handling.

ENVIRONMENTAL HAZARD

Keep out of lakes, streams or ponds. Do not contaminate water by cleaning of equipment or disposal of waste.

DIRECTIONS FOR USE

"GENERAL CLASSIFICATION"

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

STORAGE AND DISPOSAL

1. PROHIBITIONS

Do not contaminate water, food, or feed by storage, disposal or cleaning of equipment. Open dumping is prohibited. Avoid contamination of feed and food-stuff.

2. PESTICIDE DISPOSAL

Pesticide, spray mixture, or rinsate that cannot be used or chemically reprocessed should be disposed of in a landfill approved for pesticides or buried in a safe place away from water supplies.

CONTAINER DISPOSAL

Dispose of in an incinerator or landfill approved for pesticide containers, or bury in a safe place.

4. GENERAL

Consult Federal, State or Local disposal authorities for approved alternative procedures such as limited open burning.

SPRAKIL B-10 PELLETS WEED KILLER

ACTIVE INGREDIENTS:

Bromacil (5-bromo-3-methyl-2-butyl-6-methyluracil)..... 10%

INERT INGREDIENTS:..... 90%

Total..... 100%

KEEP OUT OF REACH OF CHILDREN

CAUTION

AVOID BREATHING DUST

Harmful if swallowed. In case of contact, wash skin with soap and water; for eyes, flush with water for 15 minutes and get medical attention.

See back panel for additional Precautionary Statements.

MANUFACTURED FOR

SSI INDUSTRIES, INC.
4711 MEDMONT ROAD
HUNTINGTON, W. VA. 25704

EPA Registration No. 34913-14
EPA Establishment No. 33560-TN-1

NET CONTENTS: 50 Pounds

NON-CROP BRUSH & WEED CONTROL INDUSTRIAL USE ONLY

Sprakil B-10 Pellets is a pelleted product for dry applications to the soil for the control of brush and many annual and perennial weeds. Effects are slow to appear and will not become apparent until the chemical has been activated by moisture. The degree of control and duration of effect will vary with the amount of chemical applied, soil type, rainfall and other conditions.

Sprakil B-10 Pellets is non-corrosive to equipment, non-volatile and non-flammable.

BRUSH CONTROL

To control undesirable woody plants on non-cropland areas such as RAILROAD RIGHT OF WAYS, STORAGE AREAS, INDUSTRIAL PLANT SITES and DRAINAGE DITCHES: Apply pellets in spring or early summer as a broadcast or basal (spot) treatment; for use on drainage ditches, apply as a basal (spot) treatment only. Growth of herbaceous vegetation will be prevented in the treated area for an extended period of time. Note: For effective brush control and prevention of damage to desirable vegetation: do not apply to brush standing in water; do not use water from treated ditches for irrigation; do not use in irrigation ditches nor on right of ways or other sites where marketable timber or other desirable trees or shrubs are immediately adjacent to the treated area.

BASAL (SPOT) TREATMENT

To control undesirable woody plants such as cottonwood, hackberry, maple, oak, poplar, red-bud, sweet gum, wild cherry, willow and winged elm: apply pellets in spring or early summer by spreading 1 to 2 tablespoonfuls on the ground (cover an area 1/2 to 1 sq. ft.) at the base of each brush stem or clump 2 to 4 inches in basal diameter; on larger clumps apply 1 tablespoonful to each of 3 to 4 spots around the base of the brush. Apply to uphill side of plant if ground is not level.

BROADCAST TREATMENT

Apply in spring or early summer: use 60 to 96 lbs. per acre to control oak, willow, sweet gum and pine; use 120 to 240 lbs. per acre to control American elm, winged elm, hackberry and cottonwood.

WEED CONTROL

To control most weeds for an extended period of time on non-cropland areas such as EQUIPMENT, STORAGE SITES, DITCH BANKS, AROUND FARM BUILDINGS, or other non-crop areas that are not immediately adjacent to cropland or desirable trees and shrubs: Apply preemergence or early postemergence to weed growth by spreading Sprakil B-10 Pellets uniformly over the surface of the ground to be treated. Apply at the following rates; use the higher levels of the dosage ranges on absorptive soils (those high in organic matter or carbon).

ANNUAL WEEDS AND GRASSES, such as foxtail, ryegrass, wild oat, crabgrass, cheatgrass, bromegrass, ragweed, lambquarters, puncturevine and turk, mullen: use 30 to 60 lbs. per acre.

PERENNIAL WEEDS AND GRASSES, such as smooth brome, Bahiagrass, bluegrass, redtop, purpletop, quackgrass, broomsedge, aster, dandelion, dog fennel, goldenrod, plantain and wild carrot: use 60 to 120 lbs. per acre. Higher rates of 120 to 240 lbs. per acre are required for Johnsongrass and other hard-to-kill perennials such as Bermudagrass, Vaseygrass, Dallisgrass, nutsedge, saltgrass, fountaingrass, dogbane, bracken fern and horsetail; apply these high rates only as spot treatments to localized infestations of weeds.

RETREATMENT

Apply 30 to 60 lbs. per acre when annual weeds and grasses reappear on sites where weed growth has been controlled.

FOR SMALL AREAS - one-half lb. per 435 square feet is approximately 50 lbs. per acre.