

327-4P
 ACCEPTED
 2-13-70
 By Order of
 County Clerk
 H. H. ...

Dr. Rogers'

74% Chlordane Emulsifiable Concentrate

KILLS

Grasshoppers, ants, and flies.

Active Ingredients:	
*Technical Chlordane	74.00%
Refined Petroleum Oils	15.00%
Inert Ingredients	11.00%
	<hr/>
	100.00%

*Equivalent to 44.4% Octachloro-4,7-methanotetrahydroindane and 29.6% related compounds.

WARNING: Keep out of reach of children.
 See side panel for additional precautions.

NET CONTENTS: 1 GALLON

MANUFACTURED BY
TEXAS PHENOTHIAZINE COMPANY
 FORT WORTH, TEXAS 76106

DIRECTIONS:

For best results for grasshopper control, use at least one pint Dr. Rogers' 74% Chlordane Emulsifiable Concentrate in 50 to 100 gallons of water for each acre of area to be sprayed. Each pint of Dr. Rogers' 74% Chlordane Emulsifiable Concentrate contains approximately one pound of technical Chlordane. The amount of water in which the concentrate is mixed is not so important as long as it is ample to give good coverage of area sprayed. Be sure to spray young or growing vegetation upon which the grasshoppers are feeding. When grasshoppers are young and feeding only in fence rows or at the edges of fields, spraying of such strips will often give the desired control.

CAUTION: Do not use on crops to be used for feed, food, forage, or oil purposes.

Certain species of ants and flies, may be controlled by mixing one and one-half pints to two pints of this concentrate to 100 gallons of water for an area spray for each acre of area to be sprayed. For an area control of flies, spray thoroughly all manure piles, refuse piles, and especially garbage which may be around the area. Apply spray liberally and repeat at intervals as necessary.

For control of Red Harvester Ant use at the rate of one pint of concentrate to 35 gallons of water. Pour as much of this mixture as possible into the ant holes and saturate a small area around each hill or colony. For control of houseflies in

barns, use at the rate of two quarts concentrate to 35 gallons of water. Wet thoroughly all surfaces where these insects are likely to alight or crawl, applying at least one gallon of finished spray for each 1000 square feet of surface to be treated. Repeat at intervals as necessary. For specific recommendations, consult your local Agricultural Agent.

NOTICE: Do not use this product on dairy animals nor as a spray in dairy barns.

CAUTION: Do not use on crops to be used for feed, food, forage, or oil purposes. Do not use in poultry houses.

Warning: May be fatal if swallowed. Harmful vapors. Concentrate, vapor or spray may be absorbed through skin. Do not get in eyes, on skin or on clothing. Do not breathe spray mist and vapors. Wash with soap and water. Avoid contamination of feed and foodstuffs. Keep away from children or pet animals.

Warning: Toxic to fish and wildlife. Birds feeding on treated areas may be killed. Do not contaminate streams, lakes or ponds with spray or washing from spray tanks. Rinse equipment and containers and dispose of waste by burying in non-crop lands away from water supplies. Empty containers should be disposed of by punching holes in them and burying with wastes or by burning. Keep out of sight. Do not use the concentrate material without diluting with water.

USDA Reg. No. 327-48

1-9-70