

APPROVED



**WARNINGS:**

May be fatal if swallowed. Harmful vapors. Vapor, or spray may be absorbed through skin. Do not get in eyes, on skin or on clothing. May produce irritation of skin and eyes. Respirator or goggles may be used if desired. Do not breathe spray mist and vapors. Wash thoroughly with soap and water after using. Avoid contamination of feed and foodstuffs.

**POISON** Internal: Give table salt (sodium chloride) 1 tablespoon in warm water. Repeat until vomit is clear. Call a physician.

**Directions for Use as a Spray:**

FOR CATTLE: TICKS, HORNS, EYES, and LICE use at rate of 1 gallon TPC TOX-A-DANE to 150 gallons of water, or 1 pint TOX-A-DANE to 20 gallons of water. Mix thoroughly. Spray animals all over. Thorough wetting of animals is essential for best results. Repeat as needed for control.

For use on Back-Rubbing Devices for the control of Hornflies and an aid in controlling lice on Beef Cattle. Dilute 1 pint TOX-A-DANE with 1/2 gallon diesel oil. Repeat application on devices at two to three week intervals or as needed.

FOR SHEEP AND GOATS: Lice and Sheep Ticks (Ked), use at the rate of 1 gallon TPC TOX-A-DANE to 450 gallons of water, or 1 pint TOX-A-DANE to 50 gallons of water. Spray animals thoroughly until completely wet. One application will usually give good control of these pests. However, repeat treatment as necessary for control.

FOR FLEECE WORMS AND TICKS: Use TPC TOX-A-DANE at the rate of 1 gallon to 150 gallons of water or 1 pint TOX-A-DANE to 20 gallons of water. Wet sheep thoroughly with spray. Repeat as needed.

FOR DOGS: LICE: Use TOX-A-DANE at the rate of 1 gallon to 150 gallons of water or 1 pint to 20 gallons of water. Usually one treatment will control the pest. Repeat as needed.

# TOX-A-DANE

A Combination Of  
**TOXAPHENE AND LINDANE**  
(Containing 2% Lindane and 45% Toxaphene)

Active Ingredients:

Toxaphene

Technical Chlorinated Camphene, Chrysene, Cumene, Gamma

Gamma Isomer of Benzene Hexachloride

Fresh Lindane

Refined Petroleum Oil

Xylene

Inert Ingredients:

**WARNING:**

See back panel for additional information.

**NET CONTENT: 128 FL. OZ. (1 GALLON)**

MANUFACTURED BY

**TEXAS PHENOTHIAZINE COMPANY**

P. O. BOX 4186 • FORT WORTH, TEXAS 76106

Do not treat sows within two weeks of farrow three weeks thereafter. Avoid treating anir weather. Avoid treating overheated or sick anim

Do not treat young animals under three months.

**NOTICE:** Do not use this material on dairy an

in dairy barns nor other animals being finished

Do not spray within 30 days of slaughter or

to backrubbers within 30 days of slaughter.

within 60 days of slaughter.

FOR RESIDUAL SURFACE SPRAY: FLIES AND TPC TOX-A-DANE at the rate of 1 gallon to 40 1 pint to 5 gallons of water. Spray surfaces thoroughly wet, but not to the point of exce outside walls of dairy barns, fences, inside hog cans. Repeat as needed for control.

Do not use in poultry houses. Do not use on ca hold pets.

Toxic to fish and wildlife. Do not contaminate ponds with spray or washing from spray tanks.

Rinse equipment and containers and dispose o in appropriate manner away from water supplies.

Empty containers shall be disposed of by pur an burying with wastes or by burning. Keep out of reach of children. Do not eat, drink, or smoke near containers. Do NOT eat, pour, spill or store near flame.

**NOTICE:** The use of this product is beyond the scope of the Texas Phenothiazine Company and is not represented or implied concerning th effectiveness or safety of the label. Buyer assume responsibility for the material when such use or han

EPA Reg. No. 327-62

LOT NO.



ACCEPTED



**WARNINGS:**

If swallowed. Harmful vapors, vapor, or spray may be absorbed through skin. Do not get in eyes, on skin or on clothing. Irritation of skin and eyes. Respirator or goggles may be desired. Do not breathe spray mist and vapors. Wash with soap and water after using. Avoid contamination of foodstuffs.

Oral: Give table salt (sodium chloride) 1 tablespoon. Repeat until vomit is clear. Call a physician.

**for Use as a Spray:**

TICKS, HORNFIES, and LICE use at rate of 1 gallon per 150 gallons of water, or 1 pint TOX-A-DANE to water. Mix thoroughly. Spray animals all over. Thorough treatment of animals is essential for best results. Repeat as needed.

Tick-Rubbing Devices for the control of Hornflies and other lice on Beef Cattle. Dilute 1 pint TOX-A-DANE with 100 gallons of water. Repeat application on devices at two to three week intervals as needed.

GOATS: Lice and Mange (As Keds). Use at the rate of 1 pint TOX-A-DANE to 450 gallons of water, or 1 pint to 60 gallons of water. Spray animals thoroughly until wet. One application usually gives good control; however repeat treatment as necessary for control.

FLIES AND TICKS: Use TPC TOX-A-DANE at the rate of 1 gallon to 150 gallons of water or 1 pint TOX-A-DANE to 20 gallons of water. Wet sheep thoroughly with spray. Repeat as needed.

WEEVILS: Use TOX-A-DANE at the rate of 1 gallon to 150 gallons of water, or 1 pint to 20 gallons of water. Usually one application is sufficient to control this pest. Repeat as needed.

# TOX-A-DANE

A Combination Of

## TOXAPHENE AND LINDANE

(Containing 2% Lindane and 45% Toxaphene)

Active Ingredients

Toxaphene	45.0%
Technical Chlorinated Camphene - Chlorine Content 67.69%	2.0%
Gamma Isomer of Benzene Hexachloride From Lindane	27.0%
Refined Petroleum Oil	15.0%
Xylene	11.0%
Total	100.0%

Inert Ingredients

**WARNING** keep out of reach of children. See side panel for additional cautions.

**NET CONTENT: 128 FL. OZ. (1 GALLON)**

MANUFACTURED BY

**TEXAS PHENOTHIAZINE COMPANY**  
P. O. BOX 4186 • FORT WORTH, TEXAS 76106

Do not treat sows within two weeks of farrowing nor for at least three weeks thereafter. Avoid treating animals in cold, stormy weather. Avoid treating overheated or sick animals.

Do not treat young animals under three months of age.

NOTICE: Do not use this material on dairy animals nor as a spray in dairy barns nor other animals being finished for slaughter.

Do not spray within 30 days of slaughter or permit animal access to backrubbers within 30 days of slaughter. Do not dip animals within 60 days of slaughter.

FOR RESIDUAL SURFACE SPRAY: FLIES AND MOSQUITOES: Use TPC TOX-A-DANE at the rate of 1 gallon to 40 gallons of water or 1 pint to 5 gallons of water. Spray surfaces to be treated until thoroughly wet, but not to the point of excessive run off. Spray outside walls of dairy barns, fences, inside hogs pens, and garbage cans. Repeat as needed for control.

Do not use in poultry houses. Do not use on cats, dogs, and household pets.

Toxic to fish and wildlife. Do not contaminate streams, lakes or ponds with spray or washing from spray tanks.

Rinse equipment and containers and dispose of waste by burying in non-crop lands away from water supplies.

Empty containers should be disposed of by puncturing holes in them and burying with wastes or by burning. Keep out of smoke.

DO NOT USE POUR, SPILL OR STORE NEAR HEAT OR OPEN FLAME.

NOTICE: Since the use of this product is beyond the manufacturer's control, neither The Texas Phenothiazine Company nor the seller make any warranty expressed or implied concerning the use of this product other than indicated on the label. Buyer assumes all risk of use or handling of this material when such use or handling is contrary to label instructions.

EPA Reg. No. 327-62

LOT NO.

1-10-72