

Dosage:

This 3 Oz. medication made expressly for animals of 500 to 700 lbs. body weight.

Directions:

Reduce amount of feed at evening to a very minimum then the following morning mix the entire contents of this package with the normal amount of feed. Allow horse to consume all of medicated feed before additional feed is given. Remove and destroy any medicated feed not consumed within 12 hours after being offered.

Caution: Do not treat pregnant mares within thirty days prior to foaling. Do not treat horses weighing less than 500 lbs. with this formulation. Do not use on horses intended for use as food. Do not exceed the recommended dosage. Do not retreat horses at intervals of less than 30 days.

Notice: Do not treat sick or emaciated horses except on the advice of a veterinarian. Should symptoms of overdosing such as muscular incoordination, colic, or abdominal pain appear, consult your veterinarian. Should this material cause cholinesterase inhibition Atropine is antidotal. Give 100 mg. or 1 1/2 grains per 1000 lbs. body weight.

EQUI-VERM

500-700

An Anthelmintic for Horses containing Trichlorfon and Phenothiazine. To be used in the Control and Removal of Large and Small Strongyles and Bots in Horses.

Active Ingredients:

Trichlorfon† (0,0-Dimethyl (2,2,2-trichloro-1-hydroxyethyl) phosphonate)	11.205%
Phenothiazine‡	23.517%

Inert Ingredients:

Dextrose, Cane Sugar, and Colloidal Silica	65.278%
Total	100.000%

CAUTION: Keep out of reach of children.

See side panel for additional precautions.

NET WEIGHT: 3 OZ. AVOIR.

MANUFACTURED BY

TEXAS PHENOTHIAZINE COMPANY

FORT WORTH, TEXAS 76106

Warning: For humans handling Equi-Verm 500-700: Harmful if swallowed, inhaled, or absorbed through skin. Avoid contact with the skin. Do not breathe dust. Wash thoroughly with soap and warm water after using. Do not contaminate food. Keep out of reach of children. Should poisoning occur, obtain prompt medical aid. Note to physician: Atropine is antidotal.

Contraindication: Equi-Verm is a cholinesterase inhibitor. Do not use any drug, insecticide, pesticide or chemical having cholinesterase inhibiting activity simultaneously or for a few days before or after treatment with Equi-Verm.

Do not reuse container. Destroy when empty.

‡The Phenothiazine contained in this product is from a purified technical grade of Phenothiazine.

†U. S. Patent 2,701,225

EPA REG. NO. 327-88

Lot No.

2-18-72

ACCEPTED WITH COMMENTS

ACCEPTED

JUN 19 1972

U.S. DEPARTMENT OF AGRICULTURE

OFFICE OF THE FEDERAL REGISTER

FUNGICIDE AND RELATED PRODUCTS