

[327-95]

REGISTERED  
Dec 11, 1968  
327-95

**DIRECTIONS:** Blue Death Fly Killer can be used dry "as is" or dissolved in water and used as described below. Good sanitation is essential to fly control. Remove wastes promptly and apply Blue Death daily until flies are controlled. Repeat as needed for control.

**As a Dry Bait:** Scatter lightly in strips on floors, window sills, or other clean horizontal areas in and around dairy barns, stables and animal hospitals where flies are found. Use about 4 ozs. per 1000 square feet, scattering lightly so bait is barely visible. Apply directly to garbage and manure area where flies breed and congregate. Blue Death may be applied to moistened surfaces for faster kill.

**As a Water Solution:** Mix 4 ozs. Blue Death in one gallon of water, and two cups of corn syrup or one pound of sugar and stir until dissolved. Sprinkle on floors, window sills of farm building and outside buildings where fly population is intense.

**Active Ingredient:**

Diethyl (2,2,2-trichloro-1-hydroxyethyl) phosphonate**	1%
Inert Ingredients	99%
	100%

\*\*U.S. Patent No. 2,701,225 Canadian Patent No. 529,379



**CAUTION:** Keep out of the reach of children. See back panel for additional precautions.

Net Weight — 1 Pound

Manufactured by

**TEXAS PHENOTHIAZINE COMPANY**  
FORT WORTH, TEXAS 76106

**CAUTION:** May be harmful if swallowed, inhaled or absorbed through skin. Avoid contact with skin, eyes or clothing. Wash thoroughly with soap and warm water after handling. Wash contaminated clothing with soap and hot water before reuse. Do not contaminate feed or food. Do not contaminate milk, milking equipment, feed or drinking water. If illness occurs call a physician immediately. Atropine is antidotal.

**WARNING:** Do not use in human dwellings. Do not apply directly to animals. Do not treat portion of buildings where poultry feed or which animals can lick. Direct application to surfaces previously treated with lime, whitewash or other alkali may be ineffective.

**NOTICE:** Since the use of this product is beyond the manufacturer's control, neither The Texas Phenothiazine Company nor the seller make any warranty expressed or implied concerning the use of this product other than indicated on the label. Buyer assumes all risk of use or handling of this material when such use or handling is contrary to label instructions.

U.S.D.A. REG. NO. 327-95

11-1-68

[32]


**DIRECTIONS:** Blue Death Fly Killer can be used dry "as is" or dissolved in water and used as described below. Good sanitation is essential to fly control. Remove wastes promptly and apply Blue Death daily until flies are controlled. Repeat as needed for control.

**Sugar Bait:** Scatter lightly in strips on floors, window sills, or other clean horizontal areas in and around dairy barns, stables and animal hospitals where flies are found. Use about 4 ozs. per 1000 square feet, scattering lightly so bait is barely visible. Apply directly to garbage and manure area where flies breed and congregate. Blue Death may be applied to moistened surfaces for faster kill.

**Water Solution:** Mix 4 ozs. Blue Death in one gallon of water, and two cups of corn syrup or one pound of sugar and stir until dissolved. Sprinkle on floors, window sills of farm building and outside buildings where fly population is intense.

Diethyl (2,2,2-trichloro-1-hydroxyethyl) phosphonate**	1%
Inert Ingredients	99%
	100%

\*\*U.S. Patent No. 2,701,225 Canadian Patent No. 529,379



# BLUE-DEATH SUGAR BAIT FLY-KILLER

**CONTAINING DIPTEREX\***

\*Reg. U.S. and Canadian Pat. Off. by  
Farbenfabriken Bayer, A. G. Chemagro

**CAUTION:** Keep out of the reach of children.  
See back panel for additional precautions.

Net Weight - 1 Pound

Manufactured by

## TEXAS PHENOTHIAZINE COMPANY

FORT WORTH, TEXAS 76106

**CAUTION:** May be harmful if swallowed, inhaled or absorbed through skin. Avoid contact with skin, eyes or clothing. Wash thoroughly with soap and warm water after handling. Wash contaminated clothing with soap and hot water before reuse. Do not contaminate feed or food. Do not contaminate milk, milking equipment, feed or drinking water. If illness occurs call a physician immediately. Atropine is antidotal.

**WARNING:** Do not use in human dwellings. Do not apply directly to animals. Do not treat portion of buildings where poultry feed or which animals can lick. Direct application to surfaces previously treated with lime, whitewash or other alkali may be ineffective.

**NOTICE:** Since the use of this product is beyond the manufacturer's control, neither The Texas Phenothiazine Company nor the seller make any warranty expressed or implied concerning the use of this product other than indicated on the label. Buyer assumes all risk of use or handling of this material when such use or handling is contrary to label instructions.

**U.S.D.A. REG. NO. 327-95**

11-1-68