

TPC SEVIN[®] 5% DUST

for
Fruit, Vegetables, Poultry
Lice, and Fleas on Dogs
and Cats.

DIRECTIONS: Read the entire label. Use strictly in accordance with label cautions, warnings, and directions. Use in conformity with Federal and State Regulations. Apply TPC Sevin 5% Dust when insects or their damage appear. Make applications in the morning or evening when it is cool. Treat upper and lower surface of all foliage for complete coverage. Repeat as necessary, generally at 7-day intervals. Apply with a dust gun for best results. Do not apply in excess.

BEANS: For use in the control of MEXICAN BEAN BEETLE — Use at the rate of 1/4 lb. TPC Sevin 5% Dust per 1,000 sq. ft. LEAFHOPPERS, BEAN LEAF BEETLE, CUCUMBER BEETLES, WESTERN BEAN CUTWORMS, FLEA BEETLES, JAPANESE BEETLES — 1/2 lb. per 1,000 sq. ft. CORN EARWORM, ARMYWORMS, VELVET BEAN CATERPILLAR — 1/2 to 3/4 lb. per 1,000 sq. ft. LIMA BEAN POD BORER, LYGUS and STINK-BUGS — 1 lb. per 1,000 sq. ft. May be used up to harvest. Apply thoroughly at first signs of insect infestation. When necessary, repeat every 7 to 10 days.

CUCUMBER, SUMMER SQUASH, PICKLEWORM, MELONWORM — 1/2 to 3/4 lb. per 1,000 sq. ft. CUCUMBER BEETLES, SQUASH BUGS, FLEA BEETLES — 1/2 lb. per 1,000 sq. ft. May be used up to harvest.

GRAPES, GRAPE LEAF FOLDER — 1st brood, 1/2 lb. per 1,000 sq. ft. just before first brood larvae emerge from first rolls. 2nd and 3rd Broods — 1/2 to 1 lb. per 1,000 sq. ft. May be applied up to harvest. **GRAPE LEAFHOPPER** — 1/2 lb. per 1,000 sq. ft. Thorough coverage is essential. This application may be expected to control DDT resistant leafhoppers.

ACTIVE INGREDIENT:	
Carbaryl (1-Naphthyl N-Methylcarbamate)	5.00%
INERT INGREDIENTS	95.00%
Total	100.00%

Trade name of Union Carbide Corp. for the active ingredient, 1-Naphthyl N-Methylcarbamate.

CAUTION: Harmful if swallowed or inhaled. Do not breathe dust. Do not get in eyes or on skin. Wash thoroughly with soap and water after handling. Change to clean clothing after handling. Avoid contamination of feed and foodstuffs. During application, avoid drift to adjacent food and forage crops that may be damaged or rendered unfit for sale, use or consumption.

WARNING: Keep out of reach of children and animals. Honeybees should be kept at a safe distance from treated areas.

Note to Physicians: Carbaryl is a moderate, reversible, cholinesterase inhibitor. Atropine is antidotal.

CAUTION: Keep out of reach of children. See side panel for additional precautions.

LETTUCE, MUSTARD GREENS, SPINACH, TURNIPS for Flea Beetle, Harlequin bug, Leafhopper apply 1/2 to 1 lb. per 1,000 sq. ft. **IMPORTANT:** Do not apply to head lettuce within 3 days of harvest. Do not apply to leaf lettuce or Spinach, Turnip or Mustard Greens within 14 days of harvest.

DOGS, CATS, BROWN DOG TICKS, FLEAS — Dust liberally over the animal and rub thoroughly into skin. Apply also to the feet, between the toes and on legs. Dust in and around sleeping areas and other areas where ticks and fleas occur. Apply at weekly intervals if needed. Avoid treatment of kittens less than 4 weeks of age.

CHICKENS, TURKEYS, DUCKS, GEESE, GAME BIRDS for northern fowl mite and lice, and as a supplement to premises treatment for chicken mites and fleas, dust birds with 1 lb. of 5% dust per 100 birds. As a litter treatment to control northern fowl mite, chicken mite, lice, fleas and bed bugs, treat litter at rate of 1 lb. per 40 square feet.

Caution: Do not repeat applications within 4 weeks. Do not apply to birds within 7 days of slaughter or do not house birds in rearing houses within 7 days of slaughter. Do not apply directly to eggs or litter. Do not contaminate drinking water.

U.S. D.A. Registration No. 427-98

Net Weight - 1 Pound

MANUFACTURED BY
TEXAS PHENOTHIAZINE COMPANY
FORT WORTH, TEXAS

4 26 67