

334-270

# GCC-808

## GROUND CONTROL CHEMICALS

**DIRECTIONS:** Grubs of Japanese Beetle, Rhinoceros Beetle, Asiatic Garden Beetle, Oriental Beetle, Rose, European and Masked Chafer Beetles and Ants—use 5 quarts per acre (4 fl. oz. per 1,000 sq. ft.), apply in sufficient water for thorough coverage. Make application at any time when the ground is not frozen or soggy. Spring or fall treatments are preferred. Immediately after application work or wash into top few inches of soil. Treatment will kill a high percentage of the full-grown grubs in the turf at time of application and will control annual broods for 2 or 3 years.

Sod Webworms, Wireworms, Cicada-Killer Wasp, and Wild Bees—use 2½ qts. per acre (2 fl. oz. per 1,000 sq. ft.), apply in sufficient water for thorough coverage. After applying sprinkle lightly to wash the chemical down around the crowns of the plants. Do not water again for several days; then sprinkle the grass thoroughly to wash the chemical off the grass into the soil. One application should last for several weeks, repeat if needed. For the Sod webworm make application in late afternoon or evening.

Chinch Bugs and False Chinch Bugs—use 5 qts. per acre (4 fl. oz. per 1,000 sq. ft.), apply in sufficient water for thorough coverage, sprinkle lightly and then do not water again for several days.

Armyworms, Cutworms and Male Crickets—use 5 qts. per acre (2 fl. oz. per 1,000 sq. ft.), apply in sufficient

### Liquid Insecticide

**CONTROLS SOIL-INHABITING AND LEAF FEEDING INSECTS ATTACKING LAWNS AND ORNAMENTAL PLANTS — PLUS PRE-EMERGENCE CONTROL OF CRABGRASS**

**ACTIVE INGREDIENTS:**

Technical chlordane\* ..... 72.9%  
Petroleum distillate ..... 21.6%

**INERT INGREDIENTS:** ..... 5.5%

\*Equivalent to 43.7% octachloro-4,7-methanotetrahydroindane and 29.2% related compounds.

Contains not less than 8 lbs. technical chlordane per gallon at 68°F.

### CAUTION: KEEP OUT OF REACH OF CHILDREN

Harmful if Swallowed. Contact with skin can cause toxic symptoms. Avoid breathing spray mist. In case of contact with skin, wash with soap and water. Avoid contamination of feed and foodstuffs. Do not use on humans or household pets. Keep children and pets off treated lawns until insecticide has been washed into the soil by sprinkling or rain and grass is dry. Toxic to fish and wildlife. Do not contaminate any body of water, by direct application, cleaning of equipment or disposal of wastes and containers. Note: Birds feeding over treated lawn may be harmed.

**Do Not Store Near Foodstuffs**

An exclusive product of:

**Hyson PRODUCTS**

919 West 38th Street Chicago, Illinois 60609

ACCE

water for the application in applying the several days.

European Ear—use 2 qts. sufficient water when pe be effective fo

Pre-emergence acre (1½ pts. thorough cover month before after applicatio too late, will n This rate will insects that of years. Do SEED Local conditions STATE Agr Station of Specialists ments as the control of ot

Net Contents: