

[327-126]

TPC

DIRECTIONS:

This material is a concentrate. Do not use on animals without proper dilution with water.

For Lice and Ticks:

On Beef Cattle, Non-Lactating Dairy Cattle, Sheep and Goats, use at the rate of 1 gallon AN-I-MAL SPRAY per 100 gallons of water or 6½ ounces AN-I-MAL SPRAY per 5 gallons of water and apply a complete coverage spray making sure the animals are sprayed thoroughly. One treatment may be sufficient. Repeat treatment at 2-week intervals only if needed.

For Horn Flies:

On Beef Cattle and Non-Lactating Dairy Animals, use AN-I-MAL SPRAY at the rate of 1 to 1½ gallons per 100 gallons of water and apply complete coverage spray. Repeat at 2-week intervals if needed.

For Hogs:

Use AN-I-MAL SPRAY at the rate of 1 gallon in 100 gallons of water or 6½ ounces per 5 gallons of water. Apply a complete coverage spray to animals, pens, and litter. One treatment may be sufficient. Repeat application only if needed.

AN-I-MAL SPRAY

An emulsifiable concentrate containing Malathion for use in controlling LICE, TICKS- and HORN FLIES on Beef Cattle, Non-Lactating Dairy Cattle, Hogs, Sheep and Goats.

May be applied up to the day of slaughter when used as directed.

Residue tolerance: 4 ppm in meat and meat by-products.

Active Ingredients:

Malathion—	
0,0 dimethyl, dithiophosphate of diethyl mercaptosuccinate	57.00%
Xylene	28.00%
Inert Ingredients	15.00%
Total	100.00%

WARNING: Keep out of Reach of Children.

See side panel for additional precautions.

NET CONTENT: 128 FL. OZ. (1 GALLON)

WARNING: Do not use on animals under one month of age. Do not apply to Lactating Dairy Animals, or Non-Lactating Dairy Animals with 2 weeks of freshening. Do not apply to milk goats.

WARNING: Harmful if swallowed. Avoid contact with skin. Wash thoroughly with soap and water after using. Avoid inhalation of spray mist. Avoid contamination of feed, food stuffs, food containers, and watering troughs. Do not use in milk rooms or where milk is stored. Keep out of reach of children.

Malathion is a cholinesterase inhibitor and can cause toxic symptoms similar to those caused by other organic phosphate compounds. Should poisoning occur—call a physician immediately. ATROPINE IS ANTIDOTAL. 2-PAM is also antidotal and may be administered in conjunction with atropine. Do not re-use empty container. Destroy by burying with waste or burning. Stay away from smoke or fumes.

This product is toxic to fish. Keep out of lakes, streams and ponds. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

USDA Reg. No. 327-126



MANUFACTURED BY
TEXAS PHENOTHIAZINE COMPANY
P. O. BOX 4186 • FORT WORTH, TEXAS 76106



22, 1971
 P T E D
 MALATHION
 CONCENTRATE
 REGISTERED
 SUBJECT
 MEN 79