


ACCEPTED

OUTSTANDING



A DUST MIXTURE CONTAINING COPPER AND SEVIN® USED IN THE CONTROL OF "EARLY AND LATE BLIGHT," NAIL HEAD RUST, AND FOR THE CONTROL OF THE TOMATO FRUIT WORM, CUT WORM, CUCUMBER BEETLES, AND SQUASH BUGS.

ACTIVE INGREDIENTS:	
Copper (in Basic Copper Sulphate expressed as metallic)	6%
Carbaryl® (1-Naphthyl N-Methylcarbamate)	5%
INERT INGREDIENTS	
89%	
Total	100.00%


*Trademark of Union Carbide Corp. for the active ingredient 1-Naphthyl N-Methylcarbamate

CAUTION: Harmful if swallowed or inhaled. Do not breathe dust. Do not get in eyes or on skin. Wash thoroughly with soap and water after handling. Avoid contamination of feed and feedstuffs. During application, avoid dust to crops, food and storage crops that may be damaged or rendered unfit for sale or use or consumption.

WARNING: Keep out of reach of children. Do not use in or near buildings or structures. Do not use in or near food or feed storage areas.

For more information, contact your local extension agent or the manufacturer. Always use as directed.

CAUTION: Keep out of reach of children.
See caution above.
NET WEIGHT 4 POUNDS



DIRECTIONS: Read the entire label. Use strictly in accordance with label, cautions, warnings and directions. Use in conformity with Federal and State Regulations. Apply TPC C-D Dust when insects or their damage appear. Make applications in the morning or evening when it is cool. Treat upper and lower surfaces of all foliage for complete coverage. Repeat as necessary generally at 7-day intervals. Apply with a dust gun for best results. Do not apply in excess.

EARLY and LATE BLIGHT and NAIL HEAD RUST: Use TPC C-D DUST in the "cold frame" to control and aid in the prevention of blight which may invade the "cold frame." For Early Blight, Late Blight and Nail Head Rust, start dusting soon after the tomato plants come through the ground and repeat application at weekly intervals or as often as necessary for control. For control of Tomato Late Blight on maturing plants use at the rate of 50 lbs. C-D DUST per acre or 1 lb. per 1000 square feet.

CUT WORMS: These insects are a serious menace to young tomato plants in the field and have been successfully controlled with a light dusting of C-D DUST on the plants and a more liberal dusting to the ground immediately around the plant. In heavy infestations it may be necessary to use 40 lbs. C-D DUST per acre for control of Cut Worms, or 1.2 to 3.4 lb. per 1000 square feet.

TOMATO FRUIT WORM: Successful control of the Tomato Fruit Worm depends on the proper timing. Apply TPC C-D DUST at the time the tomatoes are beginning to set. Then follow with two more applications at six day intervals. Examine the field or patch five days after the third application and if any worms or worm eggs are present apply another treatment. Use at rate 1.2 to 3.4 lb. per 1000 square feet or 20 to 30 lbs. per acre.

CUCUMBER, SUMMER SQUASH, PICKLEWORM, MELON WORM: 1.4 to 1.2 lb. per 1000 sq. ft.

CUCUMBER BEETLES, SQUASH BUGS, CUCURBIT BEETLE: 1.2 lb. per 1000 sq. ft. or 10 to 20 lbs. per acre.

Do not apply when weather is rainy or too hot for the plants to tolerate.

Apply with a dust gun or a backpack sprayer. Do not use in or near buildings or structures. Do not use in or near food or feed storage areas. For more information, contact your local extension agent or the manufacturer. Always use as directed.

Manufactured by Texas Chemical Company, Houston, Texas