

ACCEPTED
April 13, 1972

LIVESTOCK WOUND BOMB

THE NEW KILLER

- Kills fast • Saves time
- Use bomb in any position
- Worms leave wound quickly

DIRECTIONS FOR USE

Hold bomb about 3 inches from wound or infested area. Be sure spout of container is directed toward the area to be treated. Press valve down to release spray. Wet thoroughly by spraying directly into and about 1 inch around wound.

Screw worm and Blowfly infestations: Use particular care to get spray into deep pockets caused by screw worms. Repeat at 5 to 7 day intervals as needed.

Ear tick control: Spray downward directly into animal's ear. Repeat at 5 to 7 day intervals as needed.

To prevent Screw worm and Blowfly infestations: Apply directly to all open or fresh wounds such as from dehorning, castrations, shear cuts, etc. Treat navels of newborn animals. Freely bleeding wounds may require several treatments. Repeat 5 to 7 day intervals as needed.

Fleece-worm infestations: Spray infested area liberally, working toward the center of the infestation. Shearing or removing the wool from the infested area is not necessary. Repeat at 5 to 7 day intervals as needed.

Blue color added for marking treated areas.

U. S. Patent No. 2,599,516.

FOR USE ON BEEF CATTLE,
DAIRY CATTLE,
SHEEP, GOATS AND SWINE.

KILLS SCREW WORMS, EAR TICKS
BLOWFLY MAGGOTS

Contains
***KORLAN**

Active Ingredients: Ronnel 0, 0-Dimethyl 0-(2,
4, 5,-trichlorophenyl) phosphorothioate 2.5%
Xylene 0.5%
Inert Ingredient. 97.0%

For All Ages of Livestock Newborn Tool

FOR EXTERNAL USE ONLY
LIVESTOCK REMEDY — NOT FOR HUMAN USE

CAUTION: Keep out of reach of children.
See side panel for additional precautions.

NET WEIGHT: 16 OZ. (1 LB.)

MANUFACTURED BY
TEXAS PHENOTHIAZINE COMPANY
FORT WORTH, TEXAS 76106

CAUTION: INJUROUS TO EYES. Do not get in eyes. Avoid breathing spray mists. Harmful if swallowed. Avoid contact with skin. Wash hands thoroughly after using. Avoid contamination of food and feed-stuff.

In case of human eye contact, flush eyes immediately with plenty of water and get prompt medical attention.

Ronnel is a cholinesterase inhibitor; should poisoning occur, obtain prompt medical aid. Atropine is antidotal.

WARNING

Korlan is a cholinesterase inhibitor. Do not use this product on animals simultaneously or within a few days before or after treatment with or exposure to cholinesterase inhibition drugs, pesticides or chemicals. Atropine is antidote. Consult veterinarian at first sign of an adverse reaction.

WARNING

Contents under pressure; do not puncture. Keep away from heat; store at room temperature. Never throw container into fire or incinerator. Destroy container when empty.

NOTICE: Since the use of this product is beyond the manufacturer's control, neither The Texas Phenothiazine Company nor the seller make any warranty expressed or implied concerning the use of this product other than indicated on the label. Buyer assumes all risk of use or handling of this material when such use or handling is contrary to label instructions.

*REGISTERED TRADEMARK OF
THE DOW CHEMICAL COMPANY

EPA Reg. No. 327-129

Lot No.

3-20-72