

TPC

# HEP-5 FIRE ANT KILLER

### WARNINGS

Keep out of reach of children. May be fatal if swallowed. Do not breathe vapor or dust. Do not get on skin. In case of contact with skin or eyes, flush with plenty of water; for eyes, get prompt medical attention. Wash thoroughly with soap and water after handling. Avoid contamination of feed or foodstuff. Do not apply in areas where food or feed crops are grown.

This product will kill fish and wild life. Birds and other wild life in treated areas may be killed. Do not apply where runoff is likely to occur. Do not apply where weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment or disposal of waste.

Apply this product only as specified on this label. Treatment may be repeated in 10 days if necessary for control.

Do not reuse containers; destroy or dispose of empty container in a safe place.

### DIRECTIONS

**CAUTION:** Keep children and pets off treated areas until material is washed into soil and area has thoroughly dried.

**FIRE ANTS:** Area treatment of lawns, cemeteries, golf courses and park areas may be made by using 3.5 lb. of Fire Ant Killer per each 1000 square feet


of a ... This would be equivalent to 1.25 lbs. of actual Heptachlor per acre. Application should be made with a fertilizer spreader, cyclone seeder or broadcast by hand using protective gloves or equipment. Wash thoroughly with soap and water after application. Do not treat food crop areas.

Mound treatment may be made in the above areas where infestation is light or for temporary control. Tear down the mound with a hoe or rake and apply 1 to 2 cupsful of FIRE ANT KILLER to the mound and to the ground within a 3 ft. area on all sides. Do not use in food crop areas.

For use in the control of Western Harvester Ant. Apply at the rate of 25 lbs. per acre or 3.5 lb. per 1000 square feet area. Use only on lawns, cemeteries, golf courses and non-food crop areas.

**NOTICE:** Since the use of this product is beyond the manufacturer's control, neither the Texas Phenothiazine Company nor the seller make any warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use or handling of this material when such use or handling is contrary to label instructions.

EPA Reg. No. 327-130




### Active Ingredients:

Heptachlor (heptachlorotetrahydro-4,	
7-methanoindene	5.00%
Related Compounds	1.95%
Inert Ingredients	93.05%
Total	100.00%

**WARNING:** ...

**NET WEIGHT 16 OZ. (1 LB.) AVOIR.**



Lot No. ... ATTACHED ...

12-20-71

*Ornamental Turf  
Non-food crop areas*

*Ornamental Turf  
Non-food crop areas*

TPC

MANUFACTURED BY  
**TEXAS PHENOTHIAZINE COMPANY**  
P. O. BOX 4186 • FORT WORTH, TEXAS 76106

TPC

334-270

ACCEPTED

# GCC-808

## GROUND CONTROL CHEMICALS

334-270

water for thorough coverage. For cutworms make application in late afternoon. Sprinkle lightly before applying the chemical and then do not water for several days.

### Liquid Insecticide

**CONTROLS SOIL-INHABITING AND LEAF FEEDING INSECTS ATTACKING LAWNS AND ORNAMENTAL PLANTS — PLUS PRE-EMERGENCE CONTROL OF CRABGRASS**

**ACTIVE INGREDIENTS:**

Technical chlordane\* ..... 72.9%  
Petroleum distillate ..... 21.6%

**INERT INGREDIENTS:** ..... 5.5%

\*Equivalent to 43.7% octachloro-4,7-methanotetrahydroindane and 29.2% related compounds.

Contains not less than 8 lbs. technical chlordane per gallon at 68°F.

### CAUTION: KEEP OUT OF REACH OF CHILDREN

Harmful if Swallowed. Contact with skin can cause toxic symptoms. Avoid breathing spray mist. In case of contact with skin, wash with soap and water. Avoid contamination of feed and foodstuffs. Do not use on humans or household pets. Keep children and pets off treated lawns until insecticide has been washed into the soil by sprinkling or rain and grass is dry. Toxic to fish and wildlife. Do not contaminate any body of water, by direct application, cleaning of equipment or disposal of wastes and containers. Note: Birds feeding over treated lawn may be harmed.

**Do Not Store Near Foodstuffs**

An exclusive product of:

**Hyman PRODUCTS**

91 West 38th Street Chicago, Illinois 60609

**European Earwigs, Fiery Skipper and Lucerne Moths**  
—use 2 qts. per acre (1½ fl. oz. per 1,000 sq. ft.), in sufficient water for thorough coverage. Make application when pests are first noticed. Application should be effective for several weeks, repeat if needed.

**Pre-emergence Crabgrass Control**—use 8 gals. per acre (1½ pts. per 1,000 sq. ft.), in sufficient water for thorough coverage. Apply in early spring, at least a month before crabgrass germinates. Immediately after application wash chemical into the soil. If applied too late, will not give adequate control of crabgrass. This rate will also control white grubs and other turf insects that feed on the grass roots for a period of years. **DO NOT APPLY TO NEWLY**

**SEED LAWNS**

Local conditions may affect the use of this chemical. Consult **STATE Agricultural Experimental Station or Extension Service** or **Soil Conservation Specialists**.

for specific recommendations as to dosage and time of application and for the control of other insects.

Net Contents: \_\_\_\_\_ Gallons U.S.D.A. Reg. No. 334-

Ants—  
Make  
not frozen  
preferred.  
into top  
percentage  
of applica-  
or 3 years.  
Wasp, and  
per 1,000  
ugh cover-  
the chem-  
ts. Do not  
the grass  
ss into the  
ral weeks,  
ke applica-  
5 qts. per  
n sufficient  
ly and then  
—use 5 qts.  
in sufficient