

ACCEPTED

MAR 12 1975

U.S. Department of Agriculture
 Agricultural Research Service
 Entomology and Plant Quarantine
 Beltsville, Maryland
 EPA Reg. No. 327-132

TPC

**DILUTION CHART FOR TPC 10% LINDANE
 EMULSIFIABLE CONCENTRATE**

Mix with water to make indicated
 volume of Lindane spray below

Amount of TPC 10% Lindane to use	For a 0.03% Lindane Spray	For a 0.046% Lindane Spray	For a 0.3% Lindane Spray
1 Gal.	333 ² / ₅	217 ¹ / ₂	33 ¹ / ₃
1 Qt.	83 ² / ₅	54 ³ / ₈	8 ¹ / ₃
1 Pt.	41 ³ / ₅	27 ¹ / ₅	4-1 6
8 oz.	20 ⁴ / ₅	13 ³ / ₅	2
4 oz.	10 ² / ₅	6 ⁴ / ₅	1 Gal.
1 oz.	2 ³ / ₅	1 ³ / ₅	1/4 Gal.

DIRECTIONS

For Horse Barns: To control flies use a finished spray containing 0.3% Lindane (1 gallon of TPC 10% Lindane in 33¹/₃ gallons of water). Apply spray at the rate of 1 gallon finished spray for each 500 square feet area of walls, ceilings, beams, window sills, and other places where insects may alight or crawl. Coarse sprays are desirable; power sprayers with agitation and less than 100 pound pressure are recommended. Spray should be applied so as to wet all treated surfaces to the point of run off. Feed, forage, and drinking troughs should not be sprayed. Drinking troughs should be washed out immediately if sprayed accidentally.

EZ-12
(improved)

LINDANE CONCENTRATE

Containing 10% Gamma Isomer

Active Ingredients:

Xylene	73.00%
Liquid Petroleum	10.00%
Gamma Isomer of Benzene	
Hexachloride from Lindane	10.00%
Inert Ingredients:	7.00%
	Total 100.00%

CAUTION: Keep out of reach of children.

See side panel for additional precautions.

NET CONTENT:

128 FL. OZ. (1 GALLON)

MANUFACTURED BY

TEXAS PHENOTHIAZINE COMPANY

P. O. BOX 4186 • FORT WORTH, TEXAS 76106

TPC

TPC

Do not use in dairy barns, milk rooms or poultry houses.

For Scab or Sarcoptic mange mites on beef cattle, hogs, sheep, and goats use a finished spray containing 0.046% Lindane. (1 gallon TPC 10% Lindane concentrate in 217¹/₂ gallons of water).

Use a sprayer giving about 400 pounds pressure and thoroughly spray each animal all over using 1 or more quarts of spray or the amount needed to ensure that animal's skin is thoroughly wet. Repeat application at 7-day intervals for one or more treatments.

Caution: Do not dip beef cattle, sheep, goats or hogs within 60 days of slaughter. Do not spray within 30 days of slaughter.

Do not treat sows within two weeks of farrowing nor for at least three weeks thereafter.

Avoid treating animals in cold, stormy weather. Avoid treating overheated or sick animals.

Do not use on emaciated animals or dairy animals.

For Ticks and Lice: To control ticks and lice on beef cattle sheep, goats and hogs use as directed for Sarcoptic mange. Use a finished spray containing 0.046% Lindane (1 gallon TPC 10% Lindane in 217¹/₂ gallons of water). Apply as directed for Sarcoptic mange. Repeat in 14 days only if needed for control.

Note: This product should not be used above the recommended percentage levels. Beef cattle should not be dipped in order to minimize the likelihood of possible residue problems which could result from this use.

WARNING

DO NOT USE ON YOUNG ANIMALS UNDER 3 MONTHS OLD. Do not use on animals of 3 to 6

months of age except as a spray and at no higher concentration than 0.03% Lindane. Do not dip animals of this age group.

CAUTION:

HARMFUL IF SWALLOWED. HARMFUL VAPOR. CONCENTRATE, VAPOR OR SPRAY MAY BE ABSORBED THROUGH THE SKIN. MAY PRODUCE IRRITATION OF SKIN AND EYES. Avoid breathing vapor or spray mist. Avoid contact with skin and eyes. In case of skin contact, wash immediately with soap and water. Do not use on household pets or humans.

Do not contaminate drinking water, milk, milking equipment feed or foodstuff.

CAUTION: DO NOT USE, POUR, OR SPILL NEAR OPEN FLAME.

WARNING

This product is toxic to fish and wildlife. Keep out of lakes, streams, and ponds.

Rinse equipment and containers and dispose of wastes by burying in non-crop lands away from water supplies. Containers should be disposed by punching holes in them and burying with wastes, or by burning. Keep out of smoke.

NOTICE: Since the use of this product is beyond the manufacturer's control, neither the Texas Phenothiazine Company nor the seller make any warranty expressed or implied concerning the use of this product other than indicated on the label. Buyer assumes all risk of use or handling of this material when such use or handling is contrary to label instructions.

EPA Reg. No. 327-132

EPA Est. No. 327-TX-1

2-3-75