

TRE



For use in control of Horn Flies use 2 tablespoonsful (not more than 2 ounces), sprinkle lightly on poll, neck (top and sides), along backs and flanks of infested animals. Rub dust in lightly by hand to allow powder to get beneath the hair. Repeat application no more often than once every three week period if necessary.

Do not use within 5 hours before milking or during milking. Do not treat animals under 1 month of age.

For use in the control of Horn Flies use as directed for dairy cattle and rub in lightly by hand to allow powder to get beneath the hair. Repeat in 10 to 14 days if necessary.

Repeat application thoroughly.

For use on hogs, pigs and livestock, use as directed. Repeat application as necessary.

ACCEPTED
 AUG 4 1972
 FOR EXHIBIT 100
 REPT. ACQUITTANCE

CONTAINING
 METHOXYCHLOR AND
 MALATHION

Active Ingredients
 Technical Methoxychlor 1.00
 Malathion 1.00
 Inert Ingredients 91.00
 Net Weight 1.00

Net Weight 1.00
 1.00
 1.00
 1.00

State of California and may violate state laws.

Harmful if swallowed. Avoid contact with skin. Wash thoroughly after using. Avoid inhalation of dust. Avoid contamination of feed, food, water troughs. Keep out of reach of children and animals. Protect fish and birds. Do not contaminate streams and rivers with dust.

For use on dairy cattle and hogs, use as directed. Repeat application as necessary.

For use on hogs, pigs and livestock, use as directed. Repeat application as necessary.

For use on hogs, pigs and livestock, use as directed. Repeat application as necessary.