

TPC

DIRECTION FOR USE:

The treatment is best applied with a 1-inch paint brush. UNINFESTED wounds should be thoroughly covered with a coating of the material.

INFESTED WOUNDS: Brush or smear on wound and surrounding area and into deep pockets twice weekly until healed. Use minimum quantity needed, not more than three teaspoonsful.

Some strains of screwworms may develop a resistance to lindane, and may not respond to treatment.

CAUTION:

Harmful if swallowed. Avoid breathing of vapor. May be absorbed through the skin. Do not get in eyes or human skin. Wash thoroughly with soap and water

**SCREWWORM SMEAR
EQ-335**

For use in treatment and to aid in preventing screwworm infestation in wounds of horses.

Active Ingredients:

Gamma Isomer of Benzene Hexachloride from Lindane	3.0%
Pine Oil	35.0%
Inert Ingredients:	62.0%

TOTAL 100.0%

CAUTION: Keep out of reach of children.

See side panel for additional precautions.

after using. Avoid contamination of feed and foodstuffs.

Do not use on newborn or young foals under 3 months of age. For foals older than 3 months, use the minimum amount of material necessary. Do not use more than 3 teaspoonsful of EQ-335 to a wound on a young foal, and do not make application more often than twice weekly whether the wound is infested or not.

Do not use on animals other than horses.

Do not use on horses to be used for food production.

Do not reuse container. Destroy by breaking when empty.

EPA Reg. No. 327 136

NET WEIGHT - 30 OZ. (1 LB. 14 OZ.) AVOIR.

Lot No.

8-18-72

TPC

TEXAS PHENOTHIAZINE COMPANY

P. O. BOX 1180 - FORT WORTH, TEXAS 76106

TPC

