

ACCEPTED

Sept. 19, 1970

334-44

UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND RODENTICIDE ACT
FOR ECONOMIC POISON REGISTERED
ED UNDER NO. SUBJECT
TO ATTACHED COMMENT

Hysan

Will control or kill insects if used as directed. The insect cannot live in its presence because it is a double acting poison.

Whether the insects eat the powder or merely come in contact with it, death results. It acts slowly and kills within 4 to 24 hours.

This gives the insect an opportunity to crawl back from whence it came, depositing some of the powder in its path, on other insects, and in the nests.

One of the ingredients has the power to draw out the insects from its hiding place, thus making it walk through the powder.

Contact as many insects as possible.

U. S. D. A. Reg. No. 334-44
062370

227-40-2

ROACH POWDER

AN EFFICIENT INSECTICIDE

For Destroying

Roaches, Silverfish, Ants, and Water Bugs

ACTIVE INGREDIENTS:

Sodium Fluoride 40.00%
Pyrethrins .11%

INERT INGREDIENTS:

59.89%

Total 100.00%

DANGER  POISON

Keep out of reach of children. May be fatal if swallowed. Causes eye and skin irritation. See caution statement on side panel.



Manufactured By

Hysan PRODUCTS CO.

919 West 38th Street Chicago, Illinois 60609

DIRECTIONS

It is best applied by means of a powder sprayer or bellows. It may be used by sifting direct from can. Blow or dust powder thoroughly in all cracks and crevices, mouldings, under sinks, around water pipes, radiators and runways and other hiding places of insects. Thorough and systematic application cannot be stressed too strongly.

Carefully brush any powder, visible after application is completed, into cracks or crevices or else remove it. Limit application to those surfaces that are not accessible to animals, pets and children.

Repeat application when necessary.

WARNING

May be fatal if swallowed/ Do not breathe dust. Do not contaminate feed or foodstuffs.

Keep out of reach of children and domestic animals.

Empty containers should be burned or buried.

ANTIDOTES --- 1 induce vomiting.

2. Administer milk or white of egg or 5 grams of egg albumin.

3. Combat collapse. Call a physician.