

ACCE
6
UNDER THE FED
FUNGICIDE AND
FOR ECONOMIC
ED UNDER NO.

334-20

USE DILUTIONS This product must always be diluted for use. Never use full strength. See dilution chart. Effective with Soft, Medium, Hard, Cold, or Hot Water.

FOR GENERAL CLEANING AND DISINFECTING: Use a 1/2% solution. For best results add the water to the DISINFECTANT.

FOR OPERATING ROOMS, NURSERIES, DELIVERY ROOMS, PATIENTS' ROOMS, MATERNITY WARDS— This powerful Hospital Disinfectant and Deodorant acts on germ-laden surfaces such as Metal Beds and Springs, Wheel Chairs, Linen Carts, Hampers, Refuse Cans, Dressing Carts, Clean-Up Carts, Mattress Covers, Furniture, Air Ducts, Toilet Seats, Telephones, Wash Basins, Urinals, Light Switches, Office Machinery, Laundry Chutes, Door Knobs, and other equipment, after the surface has been properly cleaned. Will aid in reducing the hazard of cross infection.

On Mats in front of operating rooms: Soak mat with a 2% solution.

On Plastic Mattress Covers: Clean with 2% solution and allow to air for 24 hours.

Bed Pans, and other utensils or instruments that may come in contact with patients: Use a 2% solution.

Floors in corridors, ward rooms, etc., should be wet mopped rather than dry swept to prevent dust from redistributing the germs. Use a 2% solution.

TOILET ROOMS: Use a 1% solution. Wash bathtubs, seats, bowls, urinals, sinks, floors, and baseboards with the solution at least twice a day.

Before cleaning the bowl, remove as much water as possible by pumping it over the trap.

GARBAGE CANS, REFUSE PAILS
Thoroughly saturate by sprinkling or spraying with a 1 1/2% solution. This will aid in repelling, and preventing the breeding of flies and check putrefaction action. Repeat as often as necessary.

FOOD PROCESSING AREAS
Before using, food products and packaging materials must be removed from the room or carefully protected. Use a 1/2% solution for general cleaning and disinfection. After use, equipment and utensils in the area must be thoroughly rinsed with potable water.

EPA Reg. No. 334-205 AA

MANUFACTURED BY



CHICAGO, ILLINOIS 60609

Kysan



Fyte 13

tuberculocidal
staphylocidal • fungicidal
salmonellacidal
pseudomonacidal

**Hospital
DISINFECTANT**

ACTIVE INGREDIENTS	38.71%
Isopropyl Alcohol, Sodium Dodecylbenzenesulfonate, Ortho-Benzyl-Para-Chlorophenol, Orthophenylphenol, Tetrasodium Ethylene Diamine Tetraacetate, Para-Tertiary-Amylphenol	
INERT INGREDIENTS	61.29%
Water, Propylene Glycol, Potassium Sulphite, Citrus Perfume.	
	100.00%

WARNING: KEEP OUT OF REACH OF CHILDREN. See side panel for further precautionary statements

Net Gal

This DISINFECTANT kills **STREPTOCOCCUS AUREUS** EASILY. This as surrounding. It is used in Hospital Rooms, Corridors, and Public Conveyances, Hotels, Institutions, Schools, to kill pathogenic organisms as well as high potency permits broad use thus is economical to use.

Equal caution and care should be used on BUSES, and Other Public Conveyances, FACTORIES, OFFICES, HOME AUDITORIUMS, and various rooms. Mopping or spraying with a DISINFECTANT on all floors, wall lockers, cabins, and pantries

Effective on odors from ambulances, nursing homes, post-ileostomy dressing and bandages; in the home, kitchen, industrial plant processes, for aging, fishing equipment, from smoke odors.

DILUTION CHART	
% SOLUTION	TO MAKE 1 GALLON
1/2% Solution	20 cc or 2/3 fl oz Remainder Water
1% Solution	40 cc or 1-1/3 fl oz Remainder Water
2% Solution	80 cc or 2-2/3 fl oz Remainder Water

RECOMMENDATIONS:
USE A 1/2% SOLUTION for disinfecting FLOORS, WALLS, EQUIPMENT, RECEPTACLES, RUBBER GOODS, and as ANTI-RUST AGENT.
USE A 1% SOLUTION for disinfecting Trichophyton interdigitale.
USE A 2% SOLUTION for disinfecting PSEUDOMONAS DISINFECTANT.
USE A 2% SOLUTION for disinfecting CLOTHES: Soak for 15 minutes.

WARNING:
HARMFUL IF SWALLOWED OR IN CONTACT WITH SKIN. AVOID SPILLAGE. Avoid contact with eyes, wash with flowing water. If irritation occurs, seek medical attention. If taken internally, give olive oil or other vegetable oil. **KEEP OUT OF REACH OF CHILDREN.**

021872

ACCE
6-
UNDER THE FLD
FUNGICIDE AND
FOR ECONOMIC
ED UNDER NO.

USE DILUTIONS This product must always be diluted for use. Never use full strength. See dilution chart. Effective with Soft, Medium, Hard, Cold, or Hot Water.

FOR GENERAL CLEANING AND DISINFECTING: Use a 1/2% solution. For best results add the water to the DISINFECTANT.

FOR OPERATING ROOMS, NURSERIES, DELIVERY ROOMS, PATIENTS' ROOMS, MATERNITY WARDS— This powerful Hospital Disinfectant and Deodorant acts on germ-laden surfaces such as Metal Beds and Springs, Wheel Chairs, Linen Carts, Hampers, Refuse Cans, Dressing Carts, Clean-Up Carts, Mattress Covers, Furniture, Air Ducts, Toilet Seats, Telephones, Wash Basins, Urinals, Light Switches, Office Machinery, Laundry Chutes, Door Knobs, and other equipment, after the surface has been properly cleaned. Will aid in reducing the hazard of cross infection.

On Mats in front of operating rooms: Soak mat with a 2% solution.

On Plastic Mattress Covers: Clean with 2% solution and allow to air for 24 hours

Bed Pans, and other utensils or instruments that may come in contact with patients: Use a 2% solution.

Floors in corridors, ward rooms, etc., should be wet mopped rather than dry swept to prevent dust from redistributing the germs. Use a 2% solution.

TOILET ROOMS: Use a 1% solution. Wash bathtubs, seats, bowls, urinals, sinks, floors, and baseboards with the solution at least twice a day.

Before cleaning the bowl, remove as much water as possible by pumping it over the trap.

GARBAGE CANS, REFUSE PAILS

Thoroughly saturate by sprinkling or spraying with a 1 1/2% solution. This will aid in repelling and preventing the breeding of flies and check putrefaction action. Repeat as often as necessary.

FOOD PROCESSING AREAS

Before using, food products and packaging materials must be removed from the room or carefully protected. Use a 1/2% solution for general cleaning and disinfection. After use, equipment and utensils in the area must be thoroughly rinsed with potable water.

EPA Reg. No. 334-205 AA

MANUFACTURED BY



CHICAGO, ILLINOIS 60609

Kysan



Kyte 13

tuberculocidal
staphylocidal • fungicidal
salmonellacidal
pseudomonacidal

**Hospital
DISINFECTANT**

ACTIVE INGREDIENTS 38.71%
Isopropyl Alcohol, Sodium Dodecylbenzenesulfonate, Ortho-Benzyl-Para-Chlorophenol,
Orthophenylphenol, Tetrasodium Ethylene Diamine Tetraacetate, Para-Tertiary-Amylphenol
INERT INGREDIENTS 61.29%
Water, Propylene Glycol, Potassium Sulphite, Citrus Perfume.
100.00%

WARNING: KEEP OUT OF REACH OF CHILDREN. See side panel for further precautionary statements

Net Gal

This DISINFECTANT kills strains of STAPHYLOCOCCUS AUREUS EASILY. This disinfectant is used in Hospital Operating Rooms, Corridors, and on Public Conveyances, Schools, Institutions, Schools, to destroy pathogenic organisms as well as high potency permits broad latitude thus is economical to use.

Equal caution and care should be used on BUSES, and Other Public Conveyances, FACTORIES, OFFICES, HOMES, AUDITORIUMS, and various other places. Effective on all floors, walls, lockers, cabins, and pantries.

Effective on odors from any nursing homes, ambulances, post-ileostomy dressing and bandages; in the home, kitchen, and industrial plant processes, food processing, fishing equipment, food smoke odors.

DILUTION CHART	
% SOLUTION	TO MAKE 1 GALLON
1/2% Solution	20 cc or 2/3 fl. oz. Remainder Water
1% Solution	40 cc or 1-1/3 fl. oz. Remainder Water
2% Solution	80 cc or 2-2/3 fl. oz. Remainder Water

RECOMMENDED USE:
USE A 1/2% SOLUTION for all disinfection, FLOORS, WALLS, EQUIPMENT, RECEPTACLES, RUBBER GOODS, and as ANTI-RUST AGENT.
USE A 1% SOLUTION for fungicide disinfection of Trichophyton interdigitale.
USE A 2% SOLUTION for TRICHOPHYTON INTERDIGITALE DISINFECTION.
USE A 2% SOLUTION for COLIFORM BACTERIA DISINFECTION: Soak for 15 minutes

WARNING:
HARMFUL IF SWALLOWED. AVOID SKIN CONTACT. AVOID SPLASHING INTO EYES. IF CONTACT WITH EYES, WASH WITH RUNNING WATER. IF IRRITATION PERSISTS, CONSULT A PHYSICIAN. IF TAKEN INTERNALLY, DO NOT INDUCE VOMITING. IF TAKEN INTERNALLY, GIVE OIL OR OTHER VEGETABLE OIL IMMEDIATELY. RINSE EMPTY CONTAINER THOROUGHLY AND DISCARD IT.
KEEP OUT OF REACH OF CHILDREN

021872