

**ACCEPTED**  
 Jul 13, 1977  
 UNDER THE FEDERAL INSECTICIDE  
 FUNGICIDE AND RODENTICIDE ACT  
 FOR ECONOMIC POISON REGISTERED  
 UNDER NO. 334-23

# GCC-690

GROUNDS CONTROL CHEMICALS



**READ ENTIRE LABEL USE STRICTLY IN ACCORDANCE WITH LABEL WARNINGS AND DIRECTIONS, AND IN CONFORMITY WITH FEDERAL AND STATE REGULATIONS**

A wettable powder formulation with a broad range of effectiveness on the common diseases of turf grasses such as:

- Brown Patch** ..... (Rhizoctonia)
- Melting Out** ..... (Helminthosporium)
- Snow Mold** (Fusarium sp. and Typhula spp)
- Fusarium Patch** ..... (Fusarium)
- Dollar Spot** ..... (Sclerotinia)
- Red Thread** ..... (Corticium)
- Fading Out** ..... (Curvularia)
- Cottony Blight** ..... (Pythium)
- Copper Spot** ..... (Gloeocercospora)

### DIRECTIONS

Certain turf diseases are more prevalent in early spring while others are prevalent in summer or fall. For control of all listed diseases, start treatment in April and continue through September.

#### GOLF GREENS & FINE TURF AREAS

**PREVENTIVE PROGRAM** - Use 2-1/2 pounds in 50 gallons water to thoroughly spray 10,000 sq. ft. Under normal conditions, apply at intervals of 7 to 14 days. During periods of high humidity with heavy rainfall, it may be necessary to repeat applications every 3 to 4 days.

## LAWN AND TURF FUNGICIDE

Controls

BROWN PATCH MELTING OUT SNOW MOLD	FUSARIUM PATCH DOLLAR SPOT RED THREAD	COTTONY BLIGHT FADING OUT COPPER SPOT
---	---	---

---

**ACTIVE INGREDIENTS:**

Folpet (N-(Trichloromethylthio) phthalimide) .....	60.0%
Thiram (Tetramethylthiuram disulfide) .....	10.0%
Cadmium carbonate* .....	5.0%
<b>INERT INGREDIENTS</b> .....	25.0%
*Cadmium (expressed as metallic) .....	3.1%

---

### CAUTION: KEEP OUT OF REACH OF CHILDREN

**WARNING:** Harmful if Swallowed! If taken internally induce vomiting and call a physician at once. Do not breathe dust or spray mist. May cause skin irritation. Do not get in eyes or on skin. Wash thoroughly after handling. Avoid contamination of feed and foodstuffs. Keep out of reach of children and domestic animals. Dispose of container promptly when empty.

**This product is toxic to fish.  
 Keep out of lakes, streams and ponds.**

---

**STORE IN COOL, DRY PLACE. PROTECT FROM EXCESSIVE HEAT  
 DO NOT STORE WITH FOODSTUFFS**

Net Contents: U.S.D.A. Reg. No. 334-256  
 An exclusive product of: *Hyscon* **PRODUCTS CO.**

**919 West 38th Street Chicago, Illinois 60609**

Jul 13, 1971  
334-20  
UNDER THE FEDERAL INSECTICIDE / FUNGICIDE AND RODENTICIDE / FOR ECONOMIC POISON REGIS- / ED UNDER NO.

# GCC-690

GROUNDS CONTROL CHEMICALS



## LAWN AND TURF FUNGICIDE

Controls

**BROWN PATCH**  
**MELTING OUT**  
**SNOW MOLD**

**FUSARIUM PATCH**  
**DOLLAR SPOT**  
**RED THREAD**

**COTTONY BLIGHT**  
**FADING OUT**  
**COPPER SPOT**

**ACTIVE INGREDIENTS:**

Folpet (N-(Trichloromethylthio) phthalimide) ..... 60.0%  
Thiram (Tetramethylthiuram disulfide) ..... 10.0%  
Cadmium carbonate\* ..... 5.0%

**INERT INGREDIENTS** ..... 25.0%

\*Cadmium (expressed as metallic) ..... 3.1%

### CAUTION: KEEP OUT OF REACH OF CHILDREN

**WARNING:** Harmful If Swallowed! If taken internally induce vomiting and call a physician at once. Do not breathe dust or spray mist. May cause skin irritation. Do not get in eyes or on skin. Wash thoroughly after handling. Avoid contamination of feed and foodstuffs. Keep out of reach of children and domestic animals. Dispose of container promptly when empty.

**This product is toxic to fish.**  
**Keep out of lakes, streams and ponds.**

**STORE IN COOL, DRY PLACE. PROTECT FROM EXCESSIVE HEAT**  
**DO NOT STORE WITH FOODSTUFFS**

Net Contents:

U.S.D.A. Reg. No. 34-256

An exclusive product of:

*Hyson* **PRODUCTS CO.**

919 West 38th Street

Chicago, Illinois 60609

**CURATIVE PROGRAM** - Use 5 pounds in 50 gallons of water to thoroughly spray 10,000 sq. ft. Make at least 3 successive applications at 5 to 7 day intervals and continue until disease is halted. Reseed areas destroyed by disease and continue treatments using the PREVENTIVE PROGRAM.

#### OTHER TURF AREAS

Use 2-1/2 pounds in 50 gallons of water to thoroughly spray 10,000 sq. ft. **SPRING AND FALL** - make applications at 10 to 14 day intervals. **SUMMER** - apply as needed depending upon moisture conditions favorable to diseases. Where disease is present, follow curative program listed above.

**SNOW MOLD PROGRAM** - Snow mold may be controlled by fall or early spring applications. The best time to apply is just before the first lasting snow. Additional treatments at the preventive rate may be needed if the snow melts in mid-winter or early spring.

**COMPATIBILITY:** Do not use with highly alkaline materials. This product is compatible with wettable powder formulations of insecticides such as DDT, Dieldrin, Aldrin, Heptachlor and Lead Arsenate.