

BEST DOCUMENT AVAILABLE

GCC
GMS-337

CHAMBERLAIN
MONTANA
MONTANA

GENERAL PURPOSE SOIL SUCRINATE

For Use in the Control of Weeds and Insects in Lawns, Turf, and Ornamental Plants

ACTIVE INGREDIENTS:

Sodium Chloride (NaCl) 90.0%

Sodium Metaborate (NaBO₂) 1.0%

Monuron (3,4-dichlorophenyl 1,1-dimethylurea) 2.4%

GENERAL INFORMATION:

Apply 1 pound per 1000 sq. ft. of area to be treated. On lawns, apply as a top-dress application and on turf as a pre-emergent application. In conditions where soil moisture is low, the product will give good control.

Directions: For use on lawns, apply 1 to 1 1/2 pounds per 1000 sq. ft. of area to be treated. For use on turf, apply 1 to 2 pounds per 1000 sq. ft. of area to be treated.

EXCLUSIONS OF USE: Do not use where no control is desired, and in areas where you do not want it. (See CAUTION statement.)

- Examples are as follows:
- Right-of-way, yards, and lawns.
 - Utility pipelines, telephone, telegraph and electric power lines, power sub-stations.
 - Highways, airports, and other areas where control of weeds is desired.
 - In and about sites—petroleum and other refineries, pipelines, lumberyard, mill, bonding, and other areas.
 - Agriculture—grain fields and elsewhere, farm buildings, drainage ditches and farm roads.
 - Other—airfields, gunnery, and landing fields; outdoor theatres; tennis courts; school playgrounds.

APPLICATION:

Note: It is essential that soil should be spread uniformly at rates given.

HOW TO APPLY: Broadcast by hand or apply with a portable broadcast sprayer or mechanical fertilizer spreader.

It is essential that soil should be spread uniformly at rates given.

HOW TO APPLY: Broadcast by hand or apply with a portable broadcast sprayer or mechanical fertilizer spreader.

Special instructions for use on lawns and turf. Do not use on lawns or turf where you do not want it. Do not use on lawns or turf where you do not want it. Do not use on lawns or turf where you do not want it.

Do not use on lawns or turf where you do not want it. Do not use on lawns or turf where you do not want it. Do not use on lawns or turf where you do not want it.

BEST DOCUMENT AVAILABLE

Net Content: 50 LBS.
U.S.D.A. P. O. No. 334-X

257

ACCEPTED

MAR 29, 1967

UNLAWFUL TO USE AS AN INSECTICIDE
OR FUNGICIDE OR WEED-KILLER
OR ECONOMIC POISON REGISTERED
UNDER NO.

DO NOT STORE WITH FOOD OR FEEDSTUFFS
Store in a cool, dry place.

Mfd. by: HYSAN PRODUCTS COMPANY
CHICAGO, ILLINOIS 60609

BEST DOCUMENT AVAILABLE

BEST DOCUMENT AVAILABLE

BEST DOCUMENT AVAILABLE

GCC
GMS-337

6/1/67

GENERAL PURPOSE
INSECTICIDE
(METHYL)
DIBIPHENYL



ACTIVE INGREDIENTS:
Sodium Chloride (5%)
Sodium Metaborate (10%)
Monuron (80%)
1-dimethylurea (5%)

GENERAL INSTRUCTIONS:
AMOUNTS TO USE: For Area
Plants - Apply 1 pound per 100 sq. ft.
as a follow-up application and under
conditions will give good control.
For Lawns - Apply 1/2 to 1 1/2
1 1/2 pounds per 100 sq. ft.
Johnson, E. J. & Co. (1967) Vol. 1
10m 1/2; 2 pounds per 100 sq. ft.

EXAMPLES OF USE: use where
control is desired, and in areas
wanted. (See CAUTION! statement)

Examples are as follows:
Roads - right-of-way, yards,
Utilities - pipelines, telephone,
power line poles; power sub-stations
Highways - shoulders, signposts,
Industrial Sites - pipelines, lumberyard, rail, houses.

Agriculture - grain, rice and
drainage ditches and farm roads
Other - airfields, runways, tennis
courts; tennis courts; school

Note: It is essential that soil
at rates given.

HOW TO APPLY: Broadcast
portable seed sower or machine

RECOMMENDED RATES:
U.S.D.A. Reg. No. 334-X

REGISTERED
MAR 29, 1967
U.S. DEPARTMENT OF AGRICULTURE
REGISTERED