

ACCEPTED

AUG 30 1967

UNITED STATES PATENT AND TRADE
MARK OFFICE
FUNGICIDE AND HERBICIDE ACT
FOR ECONOMIC POLICY REGISTERED
UNDER NO. 334-266 SUBJECT
TO ATTACHED COMMENTS.

GCC-615

GROUNDS CONTROL CHEMICALS



GENERAL INFORMATION:

This brand of weed killer has proven to be a very useful selective weed killer for control of the surface creeping or prostrate broadleaf weeds listed below. Although relatively slow in action, one of the advantages of this chemical is that it is safer to fine grasses. Can be used safely on most bent varieties, but use caution on Seaside, Arlington and Congressional bents which may be injured.

WEEDS CONTROLLED:

- | | |
|---------------------|---------------------------|
| Common Chickweed | (Stellaria Media) |
| Mouse ear Chickweed | (Cerastium vulgatum) |
| Red Clover | (Trifolium pratense) |
| White Clover | (Trifolium repens) |
| Ground Ivy | (Glechoma hederacea) |
| Stitchwort | (Stellaria graminea) |
| Knotweed | (Polygonum aviculare) |
| Pigweed | (Amaranthus Spp.) |
| Ragweed | (Ambrosia spp.) |
| Black Mustard | (Brassica nigra) |
| Shepherd's Purse | (Capsella bursa pastoris) |
| Lambsquarters | (Chenopodium album) |
| Cleavers (Bedstraw) | (Galium aparine) |
| Plantain | (Plantago spp.) |
| Wild Radish | (Raphanus raphanistrum) |
| Field Pennygrass | (Thlaspi arvense) |

CLEANING OF EQUIPMENT AFTER USE:

It is difficult to clean out GCCP completely from sprayers and other equipment; therefore do not use sprayers and other equipment for purposes where even trace amounts of this chemical may cause injury. If this cannot be done, the following suggestions are made for steel tanks:

1. Wash equipment thoroughly with water after use. Fill tank with water and add an equal amount of material and mix. The following material is safe for use and may be used:

(a) 1 qt. of household ammonia to 100 gals. of water, or

(b) 2 lbs. of soda ash per 100 gallons of water.

2. After cleaning, empty tank and leave to drain in tank form. Rinse with water. Do not use for spraying in all quantities until completely clean.

3. Do not use equipment for other purposes until completely clean.

Controls clover, chickweed, stitchwort, ground ivy, knotweed and other troublesome surface-creeping broadleaf weeds invading sports and ornamental lawns, including golf greens, fairways and tees; as well as on home lawns and other non-crop turf areas.

ACTIVE INGREDIENTS:	
Diethanolamine Salt of 2-(2-methyl 4 chlorophenoxy) propionic acid*	3.20%
INERT INGREDIENTS:	
Total	91.80%
100.00%	

*Equivalent to 5.5 lbs. 2-(2-methyl 4 chlorophenoxy) propionic acid or 1/2 lbs. MCPP acid per gallon.

CAUTION: KEEP OUT OF REACH OF CHILDREN

Avoid contact with skin, eyes or clothing. Avoid inhaling spray mist. Keep out of reach of children. Avoid spray drift to susceptible plants as this product may injure tomatoes, beans, peas, ornamentals, etc. (Coarse sprays are less likely to drift). Thoroughly clean spray equipment with a suitable chemical cleaner before using for other purposes (or do not use the same equipment for other purposes). Do not transport or store near seeds, fertilizers, insecticides or fungicides. Do not use near greenhouses. Avoid contamination of water intended for irrigation of crops or domestic purposes. Protect from freezing. It will freeze, thaw and remix before using.

DO NOT STORE WITH FOODSTUFFS
An exclusive product of

WILSON PRODUCTS
WILSON PRODUCTS COMPANY
1000 W. 10th Street
Tulsa, Oklahoma 74103

DIRECTIONS FOR USE:

Application Rate on:	Per 1,000 Sq. Ft.	Per Acre
Established greens and FINE turf lawns	6 to 8 oz.	2 to 2 1/2 gals.
Other established turf and fairways	10 to 12 oz.	3 to 3 1/2 gals.

HOW TO APPLY:

Apply as a water-mixed spray in 20 to 40 gallons water per acre (4 gals. per 1,000 sq. ft.) Apply with boom type sprayer to give thorough coverage. Smaller areas may be treated with a knapsack sprayer. Do not mow for several days before or after application.

It is wise to make sure that the lawn is properly nourished when killing weeds so as to encourage the grasses to heal and spread into the bare areas created from the weeds as they die. Therefore, lightly fertilize the lawn either before or after spraying with a good fertilizer.

If reseeding is necessary, it can be done 3 to 4 weeks after initial spraying.

WHEN TO APPLY:

For knotweed apply in spring or early summer when weeds are young for best results. For other weeds listed, apply in spring or fall, when weeds are growing actively.

Better weed control, especially of knotweed, has been attained with repeat applications at half recommended rate. This is preferred, especially on bent greens and aprons where there has been no previous experience.

CAUTION:

Do not apply when it is raining or when the grass is wet. Do not apply when it is very wet weather as drift may result.

U.S.D.A. Reg. No. 334-

NET CONTENTS GALLONS