

GENERAL INFORMATION

This product, a selective weed killer for lawn grasses, is a new post-emergence herbicide specially formulated for use on turf-grass for selective control of many broadleaf weeds commonly growing in lawns without injury to desirable grasses.

This chemical may be used on all types of turf including Merion, Kentucky Blue Grasses, Bermuda, fescues and rye grasses.

(This product has been used successfully on bentgrasses; however, it should not be used unless prior tests have shown it to be non-damaging under the particular conditions of use.)

Do not use on dichondra or on newly seeded lawns.

Weeds Controlled:

Buckthorn	Lambquarters	Red Clover
Clover	Mallow	Red Sorrell
Common	Mouse-ear	Stitchwort
Chickweed	Chickweed	Speedwell
Curled Dock	Morning Glory	White Clover
Dandelions	Pigweed	Wild Garlic
English Daisy	Plantain	Wild Onion
Ground Ivy	Purslane	Yarrow
Knotweed	Ragweed	
Polson Ivy	Polson Oak	

NOTE: MCPP acts more slowly than 2,4-D, so that some weeds will be controlled before others. Allow 3 to 4 weeks for complete weed killing action.

Either before or after application, it is wise to properly fertilize the lawn so as to encourage the lawn to heal rapidly as possible, filling in the bare areas where the weeds have died. Once the weeds have been removed, the bare areas should be reseeded. Bare areas may be reseeded 3 to 4 weeks after application.

WHEN TO APPLY:

Apply in spring, or fall, when weeds are actively growing. Control of knotweed is easier in spring or early summer, when weeds are young.

000-532

ORGANIC CHEMICALS



Controls many turf and lawn broadleaf weeds and surface-creeping broadleaf weeds without injury to most ornamental lawn grasses.

ACTIVE INGREDIENTS:

Dimethylamine salt of 2-(2-methyl-4-chlorophenoxy) Propionic acid* 0.312%
 Dimethylamine salt of 2,4-dichlorophenoxyacetic acid* 0.310%

INERT INGREDIENTS: 99.378%

*Equivalent to 0.25% of 2-(2-methyl-4-chlorophenoxy) propionic acid. Equivalent to 0.25% 2,4-dichlorophenoxyacetic acid.

EPA Reg. No. 334-272 AA

**WARNING: KEEP OUT OF REACH OF CHILDREN
 CONTENTS UNDER PRESSURE**
 SEE SIDE PANEL FOR FURTHER CAUTIONS.

Net Wt. 23oz. (1lb. 7oz.)

HySan

031772
 July 7, 1972
 UNDER THE FEDERAL INSECTICIDE
 FUNGICIDE AND RODENTICIDE ACT
 FOR ECONOMIC PEST CONTROL REGISTERED
 UNDER NO. 334-272-AA

PRECAUTIONS:

Do not apply when the turf is suffering from drought or during very hot weather or damage to grass may result.

Do not mow for several days before or after application.

Withhold sprinkling for 24 hours after application. Do not apply during periods of heavy rainfall.

Do not use on newly seeded areas. Do not use if you want clover in your turf. In some areas bentgrasses, carpet, buffalo and St. Augustine grasses may be injured; therefore do not use where these grasses predominate.

Local conditions may affect the use of this chemical. Consult competent state or local turf or weed specialists for specific recommendations as to dosage and time of application, and for control of other weeds.

DIRECTIONS FOR USE

Remove cover cap. Hold can in an inverted position with the sprayhead pointed downward toward weeds, depress spray valve. Apply spray lightly just so as to wet leaves. For best results hold spray can 6 to 10 inches away from weeds. Do not water or cut lawn for 24 hours after application. Repeat treatment if necessary.

WARNING:

KEEP OUT OF REACH OF CHILDREN.

Avoid contact with skin, eyes or clothing. AVOID INHALING SPRAY MIST.

After using, wash hands and other parts of the body contaminated with spray.

Avoid spray drift to susceptible plants as it may injure grapes, tomatoes, beans, peas and other vegetables, legumes, ornamentals, etc. Do not transport with or store near seeds, fertilizers, insecticides or fungicides. Do not use in or around greenhouses. Avoid contamination of water intended for irrigation or domestic purposes. Store at temperatures above 32° F. If allowed to freeze, thaw before using.

DO NOT STORE WITH FOODSTUFFS

Contents under pressure. Do not puncture. Do not use or store near heat or open flame. Exposure to temperatures above 130° Fahrenheit may cause bursting. Never throw container into fire or incinerator.

ACCEPTED WITH COMMENTS

GENERAL INFORMATION

This product, a selective weed killer for lawn grasses, is a new post-emergence herbicide specially formulated for use on turf grass for selective control of many broadleaf weeds commonly growing in lawns without injury to desirable grasses.

This chemical may be used on all types of turf including Merion, Kentucky Blue Grasses, Bermuda, fescues and rye grasses.

(This product has been used successfully on bentgrasses; however, it should not be used unless prior tests have shown it to be non-damaging under the particular conditions of use.)

Do not use on dichondra or on newly seeded lawns.

Weeds Controlled:

Buckhorn	Lamb-quarters	Red-top
Clover	Nutsedge	Red-top
Common	Moose ear	Stitchwort
Chickweed	Chickweed	Spurge
Cuscuta	Morning Glory	White Clover
Dandelions	Prairie	Wild Garlic
English Daisy	Plantain	Wintergreen
Ground Ivy	Purple	Yarrow
Knotted	Ragwort	
Plantain	St. John's	

NOTE: MCPP acts more slowly than 2,4-D. For some weeds it will be controlled by MCPP 3 to 4 weeks for complete eradication.

Either before or after application, it is wise to properly fertilize the lawn so as to encourage the lawn to heal rapidly as possible, filling in the bare areas where the weeds have died. Once the weeds have been removed, the bare areas should be reseeded. Bare areas may be reseeded 3 to 4 weeks after application.

WHEN TO APPLY:

Apply in spring or fall, when weeds are actively growing. Control of knotweed is easier in spring or early summer, when weeds are young.

GCC-532

GROUNDS CONTROL CHEMICALS



Controls many turf and lawn broadleaf weeds and surface-creeping broadleaf weeds without injury to most ornamental lawn grasses.

ACTIVE INGREDIENTS:

- Dimethylamine salt of 2-(2-methyl-4-chlorophenoxy) Propionic acid* 0.312%
- Dimethylamine salt of 2-(4-chlorophenoxy)acetic acid* 0.310%

INERT INGREDIENTS:

- *EPA Reg. No. 227-272-40
- 2-(2-methyl-4-chlorophenoxy) Propionic acid 0.312%
- 2-(4-chlorophenoxy)acetic acid 0.310%

EPA Reg. No. 334-272 AA

WARNING: KEEP OUT OF REACH OF CHILDREN
CONTENTS UNDER PRESSURE
SEE SIDE PANEL FOR FURTHER CAUTIONS.

Net Wt. 23 oz. (1 lb. 7 oz.)

MANUFACTURED BY

 CORPORATION, CHICAGO, ILLINOIS 60609

PRECAUTIONS:

Do not apply when the turf is suffering from drought or during zero frost weather or damage to grass may result.

Do not mow for several days before or after application.

Withhold sprinkling for 24 hours after application. Do not apply during periods of heavy rainfall.

Do not use on newly seeded areas. Do not use if you want clover in your turf. In some areas bentgrasses, carpet buffalo and St. Augustine grasses may be injured; therefore do not use where these grasses predominate.

Local conditions may affect the use of this chemical. Consult competent state or local turf or weed specialists for specific recommendations as to dosage and time of application, and for control of other weeds.

DIRECTIONS FOR USE

Remove cover cap. Hold can in an inverted position with the sprayhead pointed downward toward weeds, depress spray valve. Apply spray lightly just so as to wet leaves. For best results hold spray can 6 to 10 inches away from weeds. Do not water or cut lawn for 24 hours after application. Repeat treatment if necessary.

WARNING:

KEEP OUT OF REACH OF CHILDREN
 Apply to lawn with care. Do not apply to plants, shrubs, or trees. Avoid contact with hands, face, and eyes. Avoid contact with clothing. Do not use on plants, shrubs, or trees. Do not use on plants, shrubs, or trees. Do not use on plants, shrubs, or trees. Do not use on plants, shrubs, or trees.

DO NOT STORE WITH FOODSTUFFS
 Do not store with foodstuffs. Do not store with foodstuffs. Do not store with foodstuffs. Do not store with foodstuffs. Do not store with foodstuffs. Do not store with foodstuffs.