

DIRECTIONS

RATS: 4oz. to 1 lb
Place a small amount, ~~in a~~
on a piece of paper in locations frequent-
ed by rats or where rats have been known
to feed. Keep out of reach from dogs,
cats, poultry, livestock, etc. by using bait
boxes, cages, or other suitable means.
Baits should be examined regularly and
replenished before completely eaten. If
bait becomes dirty, it should be replaced.

KEEP RAT BAIT IN BAIT BOX
OR OTHER PROTECTED PLACE

MICE:
Follow the same general directions as for
rats except that the amount should be
smaller (1 tablespoonful) and more re-
placements should be made.

KEEP BAIT IN BAIT BOX
OR OTHER PROTECTED PLACE

Mice are
more difficult to control than rats and
complete control may take a longer period
of baiting.

KEEP BAIT IN BAIT BOX
OR OTHER PROTECTED PLACE

331-283
ACCEPTED
Jul 10 1968
THE FEDERAL INSECTICIDE
AND FUNGICIDE ACT
REGISTERED SUBJECT
TO ATTACHED COMMENTS.



KILLS RATS AND MICE

CONTAINS WARFARIN*

Ingredients:

Active: Warfarin	
3-(α -acetylbenzyl)-4-hydroxycoumarin*	.025%
Inert	99.975%
Total	100.00%

*Warfarin is a patented product discovered in the
laboratories of Dr. Karl Paul Link, Biochemistry De-
partment, University of Wisconsin.
Wisconsin Alumni Research Foundation Compound #42
—U. S. Patent #2,427,578

CAUTION
KEEP OUT OF REACH
OF CHILDREN. SEE
SIDE PANEL FOR
FURTHER CAUTION
STATEMENTS

IMPORTANT

Baiting should be continued as long as
evidence of feeding is observed. Except
in extremely heavy infestations, this should
be from five days to two weeks. Where
a continuous source of infestation is
present from nearby dumps or fields, per-
manent bait stations should be used and
bait placed in them as often as necessary.

CAUTION

KEEP OUT OF REACH
OF CHILDREN

This rat killer contains as its active in-
gredient an anti-coagulant chemical,
therefore, it must be protected from or
placed where humans, livestock, poultry or
pets will not accidentally eat them. In
case it is ~~accidentally~~ eaten, give a table-
spoonful of salt in a glass of warm water
and repeat until vomit fluid is clear. Call
a physician immediately.

Note to Physicians:

When a human has been known to have
accidentally ingested any of this rat killer,
blood transfusions combined with intra-
venous injections and oral doses of Vitamin
K are indicated as in the case of hem-
orrhage caused by overdoses of Dicumarol.

U. S. D. A. Reg. No. 334-

Net Contents 1 lb.