Reg # 334-381
PM17
p 1/1

20.
24oz
32.

DIRECTIONS FOR USE

SHAKE BEFORE USING

ake well before each use. Remove protective cap. Aim spray opening away from person. Push ton to spray.

NTROL HOUSEHOLD PESTS

Houseflies, Mosquitoes and Gnats. Close all doors and windows. Point nozzle upward, direct ay mist to all parts of the room, especially windows and other light sources which attract these ects. Fill the room with mist, then leave the treated area. Keep the room closed for at least minutes. Ventilate the room when treatment is completed.

Crawling Insects: Roaches, earwigs, grain mites, sowbugs, spiders and centipedes. Spray roughly into hiding places, cracks, crevices, moist areas, openings around pipes and sinks, der refrigerators, baseboards and storage areas. Spray directly on insects where possible.

ts: Spray trails, nests and points of entry. Spray on ants where possible. Repeat as necessary.

TDOORS—TO REDUCE ANNOYANCE FROM FLYING INSECTS. Reduce mosquitoes, gnats, dges, houseflies, biting flies and flying moths (millers) by spraying their resting areas. Follow cautions listed above. Hold container 18 to 24 inches from shrubs, bushes and grassy areas. ay these areas with a slow sweeping motion at the rate of from 1 to 3 seconds per cubic yard space while moving away from the treated areas. To reduce annoyance from these flying in cts, treat open porches and patios in a similar manner. Spray only when the air is still. Allow ay to dissipate before occupying the treated area. Repeat as necessary.

TDOORS—TO PROTECT ORNAMENTAL PLANTS. To control aphids, exposed thrip, adult te flies, leafhoppers, two spotted mites, plant bugs, spittlebugs and Japanese beetle on smaller ododendrons, pines, spruce, hemlock, spirea, dogwood, palm, mimosa, [illegible], mymus, oaks, maples, locust, forsythia and althea, and mimosa webworm larvae on mimosa. ay directly on insects. Hold container 18 inches from the infested plants. Spray in short spray ts of one to three seconds or spray at the rate of one linear foot of growth per second. Direct ay to all sides covering foliage until slightly moist. Avoid heavy wetting. Repeat as necessary obtain control.

ay pecnies, iris, [illegible], snapdragons, [illegible], chrysanthemums, nasturtiums, daisy, aniums, marigolds, daphne, fatshedera, ivy, lobira, queen sago and bread leaf, acacia for con- of aphids, exposed thrips, adult white flies, leafhoppers, plant bugs, spittlebugs and Japanese ettles. Spray as directed above.

E WITH CAUTION on azaleas, junipers, roses, including hybrid tea roses, flowering quince and camellias. These plants may be subject to leaf or flower burn under certain conditions of use.

adult whiteflies, two spotted mites be sure to direct spray to contact these pests. Thoroughly er the infested area, use several repeat applications at two day intervals if necessary.

OTECT HOUSE PLANTS. Spray infested African violets, philodendron, geraniums, begonias, ns for listed pests. Leave the sprayed area when done indoors, and ventilate area after treat nt is completed.

ISANCE PESTS—ON OR ENTERING BUILDINGS

ape: Applications should be made in late evening when insects are at rest. Spray liberally into ing and breeding places, contacting as many insects as possible. Repeat as necessary. Spray ectly on stray wasps that enter buildings.

ts: Spray trails, nests and points of entry. Spray on ants where possible. Repeat as necessary.

dera: Direct short burst of spray on spiders and repeat as necessary.

nwigs: Spray directly on earwigs in crevices and other hiding places on outdoor porches, patio ar and doorway entrances.

CAUTION: KEEP OUT OF REACH OF CHILDREN

rmful if swallowed. Avoid contamination of feed and foodstuffs. Do not use on edible crops. en treating areas where food is prepared or processed, foods should be removed or covered ing treatment or thoroughly cleaned before using. When using the product in these areas, ly only when the facility is not in operation. Remove pets, birds and cover fish aquariums before aying. Contents under pressure. Do not use near heat or open flame. Exposure to temperatures ove 130°F may cause bursting. Avoid inhalation of spray mist. Avoid contact with skin.

STORAGE AND DISPOSAL

ORE in a cool, dry area away from flames, sparks, and other sources heat and ignition. Containers heated above 130°F (54°C) may burst. r DISPOSAL replace cap and discard container in trash. Do not cinerate or puncture.

PA Reg. No. 334-33[illegible] EPA Est. 334-IL-2

MANUFACTURED BY

hysan CORPORATION

MAIN OFFICE

CHICAGO, IL 60609

Arlington, TX. 76011 • Atlanta, GA 30319

Blue Island, IL. 60406

OERA Made in U.S.A.

PS 75

INSECT KILLER

KILLS: Houseflies, mosquitoes, gnats, wasps, roaches, earwigs, two-spotted mites, sowbugs, spiders, centipedes, ants, biting flies, aphids (plant lice), leafhoppers, exposed thrips, adult whiteflies, plant bugs, Japanese beetles, spittlebugs.

ACTIVE INGREDIENTS:	
*[illegible] 3 furylmethyl 2,2 dimethyl 3	
[illegible]	0.2[illegible]% w/w
Related compounds	0.028%
d-trans Allethrin (allyl homolog of Cinerin I)	0.150%
Related compounds	0.012%
[illegible] hydrocarbons	0.272%
INERT INGREDIENTS:	99.338%
TOTAL	100.000%

*Lis. trans-cis-ratio max. 30% (±) cis and min. 70% (±) trans.
TUS Patent No. 3,465,007

NET WT. 17 oz. (1 lb. 1 oz.) (481 g)

KEEP OUT OF REACH OF CHILDREN

CAUTION

CONTENTS UNDER PRESSURE.

SEE BACK PANEL FOR FURTHER PRECAUTIONS.

ACCE[illegible]

DEC 15 1984

Under the [illegible] Fungicide [illegible] contained [illegible] registered under [illegible] Reg. No. 334-381