

# LINDANE 75 WP DYED SEED TREATER

Agricultural Insecticide

Wettable Powder

ACTIVE INGREDIENT: LINDANE (Gamma Isomer of Benzene Hexachloride) 75.0%  
INERT INGREDIENTS: 25.0%  
100.0%

FOR AGRICULTURAL USE ONLY

KEEP OUT OF REACH OF CHILDREN

## WARNING

### Statement of Practical Treatment

**If swallowed** — Drink one or two glasses of water and induce vomiting by touching the back of the throat with finger. Repeat until vomit fluid is clear. Call a physician immediately. Do not induce vomiting or give anything by mouth to an unconscious person.

**If inhaled** — Remove victim to fresh air. Apply respiration if indicated.

**If on skin** — Remove contaminated clothing and wash affected areas with soap and water.

**If in eyes** — Flush eyes with plenty of water. Call a physician if irritation persists.

See side panel for additional precautionary statements.

STORE IN A COOL, DRY PLACE  
STIR CONTENTS THOROUGHLY INTO  
WATER BEFORE USE

**POUNDS NET**  
Made in U.S.A. by


AGRICULTURAL PRODUCTS COMPANY, Ft. Worth, TX 76112

## PRODUCT INFORMATION:

**LINDANE 75 WP SEED TREATER** is a microfine wettable powder agricultural insecticide designed especially for treatment of seeds against wireworms, seed corn maggots, and other soil-borne insects. It contains an effective bonding agent to adhere Lindane on the seeds during sacking, storing, and planting, and is applied for wet application through slurry treaters.

## SEED TREATMENT DOSAGE RATES

Type of Seed	Ounces per 100 pounds
Alfalfa Clover	1.02
Berley	1.02
Beets	1.02
At least 1 Standard & Eckbreck Large Lentils	1.02
Standard & Eckbreck Large Lentils	1.02
Broccoli, Brussels Sprouts, Cabbage, Cauliflower, Radish	1.02
Cantaloupe, Cucumber, Pumpkin, Squash, Water-melon	1.02
Corn	1.02
Cotton, Acid or Machine Dressed	1.02
Cowpeas	1.02
Fava	2.02
Lettuce	1.02
Milo, Soy, Sun	1.02
Oats	1.02
Onion	1.02
Onions	1.02
Peanut	1.02
Rye	1.02
Safflower	2.02
Sorghum	1.02
Soybeans	1.02
Sprach	1.02
Sudan Grass	1.02
Sugar Beets	1.02
Sunflower	2.02
Tanqueret	1.02
Wheat	1.02



**ONLY SEED OF HIGH GERMINATION** should be used. Poor seed may suffer from the effects of the insecticide. The effect of the insecticide on the seed is dependent on the soil temperature and the moisture content of the soil. The effect of the insecticide on the seed is dependent on the soil temperature and the moisture content of the soil.

**IF SOIL TEMPERATURE IS EXTREMELY HIGH**, the insecticide may be washed away from the seed. The insecticide may be washed away from the seed if the soil temperature is extremely high.

### NOTICE — READ CAREFULLY

**CONDITIONS OF SALE** — This product is sold as a "best" product. Users are deemed to have accepted the product as such and are hereby notified that the product is sold as is, without any warranty, expressed or implied.

**WARRANTY LIMITATION** — The manufacturer's liability is limited to the chemical description on the label. The manufacturer is not liable for any damage or loss resulting from the use of this product.

**OF MERCHANTABILITY** — The manufacturer is not liable for any damage or loss resulting from the use of this product. There is no implied warranty of merchantability.

**INHERENT RISKS** — The user is advised that the use of this product is subject to the risk of injury and should be used with care. The user is advised that the use of this product is subject to the risk of injury and should be used with care.

**LIMITATION OF LIABILITY** — The manufacturer is not liable for any damage or loss resulting from the use of this product. The manufacturer is not liable for any damage or loss resulting from the use of this product.

EPA Reg No 43789-40  
EPA Est 43789-TX-01