AQUA – KILL RESIDUAL HOUSEHOLD INSECT KILLER

(Unscented, Country or Cedar Fragrance)

- CONTAINS PYRENONE® AND PERMANONE®
- CONTROLS RESIDENTIAL AND GARDEN INSECTS
- SEEKS OUT BUGS WHERE THEY HIDE
- KILLS ON CONTACT
- FOR HOUSEHOLD USE
- NON-STAINING

)

- A BROAD SPECTRUM INSECTICIDE HIGHLY EFFECTIVE AGAINST ANTS, (INCLUDING CARPENTER ANTS), BEES, BEETLES (INCLUDING CARPET, FLOUR, GRAIN AND GROUND BEETLES), CENTIPEDES, CLOVER MITES, COCKROACHES, CRICKETS (INCLLUDING MOLE CRICKETS), EARWIGS, FIREBRATS, FLEAS, MILLLIPEDES, MOTHS, PILLBUGS, SCORPIONS, SILVERFISH, SOWBUGS, SPIDERS, TICKS, WASPS, AND OTHER PESTS LISTED ON THE LABEL.
- KILLS LICE AND THEIR EGGS
- FOR USE ON BEDDING, FURNITURE AND OTHER INANIMATE OBJECTS INFESTED WITH LICE
- NO UNPLEASANT ODOR
- WATER-BASED FORMULA
- FOR USE ON DOGS AND ON PREMISES TO CONTROL FLEAS AND TICKS
- CONTROLS FLEAS AND TICKS ON DOGS FOR 14 DAYS
- PROTECTS FOR UP TO (4 WEEKS, 12 WEEKS, 3 MONTHS)

ACTIVE INGREDIENTS

Pyrethrins	0.10%
*Piperonyl Butoxide, Technical	0.50%
**Permethrin	0.20%
INERT INGREDIENTS	99.20%

*Equivalent to min. 0.4% (butylcarbityl)(6-propylpiperonyl) ether and 0.1% related compounds.

**(3-phenoxyphenyl)methyl(+/-) cis/trans 3-(2,2-dichloroethenyl) 2,2-dimethyleyelo-propanecarboxylate. Cis/trans ratio: min 35% (+/-) cis and max. 65% (+/-) trans Pyrenone and Permanone – Registered Trademarks of AgrEvo Environmental Health, Inc.

EPA REG. NO. 334-570 EPA EST. NO. 34702-OH-1

KEEP OUT OF REACH OF CHILDREN CAUTION STATEMENT OF PRACTICAL TREATMENT

IF INHALED: Remove victim to fresh air. If not breathing, give artificial respiration, preferably mouth to mouth. Get medical attention

See Rear Panel for Use Cautions

ACCEPTED

Under the Follows Insecticide. Franchische, and Indonticide Recus unsweller, and insecticide Reregiment of \$34-570 NET CONTENTS:

PRECAUTIONARY STATEMENTS Hazards to Humans & Domestic Animals CAUTION

Harmful if inhaled. Avoid breathing vapor or spray mist. Remove contaminated clothing and wash before reuse. Avoid contamination of feed and foodstuffs. Remove pets and birds and cover fish aquariums before spraying. Vacate room after treatment and ventilate before reoccupying. Do not allow children or pets to contact treated areas until surfaces are dry.

Environmental Hazards

This product is toxic to fish. Do not apply directly to any body of water. Do not contaminate water when disposing of equipment washwaters.

Physical or Chemical Hazards

Contents under pressure. Do not use or store near heat or open flame. Do not puncture or incinerate container. Exposure to temperatures above 130 degrees F may cause bursting. Do not use this product in or on electrical equipment due to the possibility of shock hazard.

Directions for Use

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

In the home, all food processing surfaces and utensils should be covered during treatment or thoroughly washed before use. Exposed food should be covered or removed.

Do not use in rooms/areas while occupied by patients, the elderly or infirm. Do not apply to classrooms when in use.

SHAKE WELL BEFORE USING

Remove protective cap, hold container upright and spray from a distance of 12 to 15 inches. **INDOORS:**

Cluster files, houseflies, fruit flies, mosquitoes, gnats, wasps, hornets, bees, black flies, skipper flies, blow flies, and small flying moths: Close all doors and windows and direct spray upward into center of room with a slow sweeping motions. Spray 5-10 seconds for average room, moving toward door. Direct spray to all parts of the room, especially windows and other light sources which attract these insects. Keep nozzle in constant motion. Leave room and keep area closed for 15 minutes after spraying. Ventilate room thoroughly before re-entry. Sweep up and discard fallen insects. Repeat as necessary.

Cockroaches, waterbugs, palmetto bugs, silverfish, firebrats, crickets, scorpions, spiders, millipedes, centipedes, sowbugs and pillbugs: Spray into hiding places such as cracks and crevices, behind sinks, cabinets, along baseboards and floors, around drains and plumbing, hitting insects with spray whenever possible. Repeat treatment as necessary.

Ants: Spray trails, nests and points of entry. Spray on ants where possible. Repeat as necessary.

Carpet Beetles: Spray directly on beetles, floor, baseboards, shelves and carpets. Repeat treatment as necessary.

Bedbugs: Spray mattresses, particularly around tufts and seams. Take beds apart and spray into all joints. Treat baseboards, moldings and floors. Repeat treatment as necessary. Allow all sprayed articles to dry thoroughly before use.

To control clothes moths and carpet beetles: on wool and other proteinaceous fibers. Remove infested articles from storage, brush thoroughly, and air for several hours in sunlight, is possible. Apply spray liberally to empty chests, closets, bureaus and other storages directing spray into cracks, crevices and joint. After airing, infested articles may be treated by spraying lightly. Keep nozzle at least feet from fabric to avoid staining. Fleas in Buildings - Space Spray: Hold can 36 inches above floor and direct spray toward floor and lower walls at a rate of 10 seconds per 100 sq. ft. making sure that all floor, sofa and chair surfaces are contacted. Leave room and keep area closed for 15 minutes. Open and ventilate before reoccupying. Repeat treatment after 7 days or as needed.

Fleas & Ticks (Adults and Larvae): Remove and thoroughly clean or destroy pet bedding. Spray pet sleeping quarters, bed, entire floor and floor covering, with special attention to cracks and crevices around baseboards and window and doorframes. Infested upholstered furniture and mattresses may also be treated with this product. A test application should be made to upholstery or drapery fabrics in an inconspicuous place before use.

To Control Fleas, Ticks and Lice on Dogs: Starting at the head, stroke against the lay of the hair spraying the parted fur in short bursts, directly behind the hand to insure penetrating the coat. Dampen the fur back to the tail. Repeat application weekly if necessary to control infestation.

Note: Not recommended for use on puppies less than four weeks old.

To Kill Lice and Louse Eggs: Spray in an inconspicuous area to test for possible staining or discoloration. Inspect again after drying, then proceed to spray entire area to be treated. Hold container upright with nozzle away from you. Depress valve and spray from a distance of 8 to 10 inches. Spray each square foot for three seconds. Spray only those garments and parts of bedding, including mattresses and furniture that cannot be either laundered or dry-cleaned. Allow all sprayed articles to dry thoroughly before use. OUTDOORS:

Point spray nozzle away from face. Hold can on slight downward angle. For best results, spray when air is calm. Allow a few minutes for product to take effect. Spray with wind if breeze is blowing.

Flying Insects: Flies, Mosquitoes, Gnats, Small Flying Moths, Hornets, Wasps and Yellow Jackets. Allow fog to drift over area. Direct spray toward bushes, or grass from at least 4 to 6 feet away. Give special attention to breeding areas such as garbage cans and compost piles. Repeat as necessary.

Mud Daubers and Wasp Nests: Spray mist directly on insects and their nests from approximately 2 feet away. Contacted insects will fly away from fog. Applications should be made in late evening when insects are at rest. Spray into hiding and breeding places contacting as many insects as possible.

Crawling Insects: Ants, Cockroaches, including Asian Cockroaches, Crickets, Moles Crickets, Centipedes and Silverfish. Spray infested surface of patio or picnic area, hitting as many insects as possible. Also spray legs of tables and chairs to repel ants. Spray anthills for best control.

For the Control of Garden Insects: For use on roses, dahlias, asters and other ornamentals to kill Japanese Beetles hit by spray, as well as Whiteflies, Aphids, exposed Thrips, Spiders, Red Mites and Leafminers. May be used on plants, as well as a multipurpose spray in small gardens and for spot treatments of incipient insect infestations to prevent spreading in large garden areas. May also be used on other plants such as African violets, asters, azaleas, begonias, camellias, carnations, crysanthemums, dahlias, delphiniums, dogwood, English ivy, euonymus, fuschia, geraniums, crassula, grape vine, Kentia palm, laurel, marigolds, rhododendions, roses. rubber plants, snapdragons, stocks, wandering. Jews and zinnias.

Effective when used as directed against Whiteflies, Aphids, Spider and Red Mites, Exposed Thrips, Lace Bugs, Japanese Beetles, Pavement Ants, Clover Mites and Army Worms. When spraying plants, do not operate closer than 18 inches. Use sweeping motion. Spray in one-second bursts. Be cautious about wetting tender foliage, young plants and new growth. Do not spray plant when temperatures exceed 90 degrees F. Contact insects directly whenever possible. Repeat treatment as infestations occur.

STORAGE AND DISPOSAL

Store in a cool, dry area. Do not transport or store below 32 degrees F.
Wrap container in several layers of newspaper and dispose of in trash. Do not incinerate or puncture.

Manufactured by: Hysan/Amp, a division of Specialty Chemical Resources, Inc 9055 Freeway Drive Macedonia, OH 44056 1-800-321-9500