

DANGER

KEEP OUT OF REACH OF CHILDREN. MAY CAUSE CHEMICAL BURNS. DO NOT GET IN EYES, SKIN AND CLOTHING. MAY BE HARMFUL OR FATAL IF SWALLOWED.

STRONG OXIDIZING AGENT. NEVER MIX WITH ANY MATERIALS (INCLUDING OTHER POOL CHEMICALS) OTHER THAN WATER. CONTACT WITH FOREIGN MATERIAL, HEAT, FRICTION, MOISTURE, OR CONTAMINATION FROM ANY SOURCE MAY RESULT IN EXPLOSION OR FIRE OF GREAT INTENSITY WITH THE LIBERATION OF HAZARDOUS AND IRRITATING GASES. USE ONLY A CLEAN DRY CUP OR DISPENSER. DO NOT REUSE CONTAINER. RINSE EMPTY CONTAINER THOROUGHLY WITH WATER BEFORE DISCARDING.


ANTIDOTE

EXTERNAL — Wash thoroughly with water. For eyes get prompt medical attention.

INTERNAL — If swallowed, drink promptly a large quantity of milk, egg whites, gelatin solution or, if these are not available, drink large quantities of water. Avoid alcohol. Call physician immediately. (Note to physician: Mucosal damage may contraindicate the use of gastric lavage.)

100 LB.
NET WT.

ACCEPTED
FFB 27 1974
335-195
UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND RODENTICIDE ACT
FOR ECONOMIC POISON REGISTERED
ED UNDER NO. _____ SUBJECT
TO ATTACHED COMMENTS.



POOL-CL



Concentrated Chlorine for Sw

ALGAEICIDE • BACTERICIDE • D

ACTIVE INGREDIENT: Sodium Dichloro-s-Triazinetrione (Sodium
AVAILABLE CHLORINE 63%

DANGER: KEEP OUT OF REACH OF
See Back Panel for Additional Precautions and Ar

PENN SWIM DEPARTMENT • PENN WALT CORPOF



POOL-CLOR

Chlorine for Swimming Pools

BACTERICIDE • DISINFECTANT

100% Dichloro-s-Triazinetrione (Sodium Dichloroisocyanurate) 100%
AVAILABLE CHLORINE 63%

KEEP OUT OF REACH OF CHILDREN
Read Label for Additional Precautions and Antidote Statements

DIRECTIONS FOR USE

Pool-Clor Concentrated Chlorine is a fast dissolving granular powder for chlorination of swimming pool water.

For pool start-up, first adjust the pH of water to 7.2-7.6 with a proper additive such as Pennswim Aquet® or Dry Acid according to the instructions on their label, then add at least 4 oz. of Pool-Clor Concentrated Chlorine for each 10,000 gallons of water to satisfy the chlorine demand of the water and reach the initial chlorine residual of 0.6 ppm (stabilized pool required 1.0 ppm). For the stabilization of the chlorine in the pool, add Pennswim Pool Stabilizer (following the directions for use on its label) at the beginning of the pool season.

To prevent the growth of algae, superchlorinate at the start of the season, and then once every one or two weeks with PENNSWIM BIGSHOT, used according to the instructions on its label.

Continue with 2 oz.-3 oz. of POOL-CLOR Concentrated Chlorine per 10,000 gallons of water as necessary to maintain the minimum chlorine residuals. Health regulatory agencies usually require 0.6 to 1.0 ppm for non-stabilized and 1.0 to 1.5 ppm for stabilized pools, determined by daily or more frequent checks with a reliable chlorine test kit. pH of the pool water should be maintained in the 7.2-7.6 range as determined by daily checks with a test kit. Variations in temperature, sunlight, organic matter (leaves, grass, dust, etc.) and bathing loads will influence the rate of chlorine consumption and determine Pool-Clor Concentrated Chlorine dosages.

This product is toxic to fish. Treated effluent should not be discharged where it will drain into lakes, streams, ponds or public water.

EPA Reg. No. 335-195

Printed in U.S.A.

Made in U.S.A.

4-6650A-01

T • PENN WALT CORPORATION • PHILA., PA. 19102

DANGER

KEEP OUT OF REACH OF CHILDREN. MAY CAUSE CHEMICAL BURNS. DO NOT GET IN EYES, SKIN AND CLOTHING. MAY BE HARMFUL OR FATAL IF SWALLOWED.

STRONG OXIDIZING AGENT. NEVER MIX WITH ANY MATERIALS (INCLUDING OTHER POOL CHEMICALS) OTHER THAN WATER. CONTACT WITH FOREIGN MATERIAL, HEAT, FRICTION, MOISTURE, OR CONTAMINATION FROM ANY SOURCE MAY RESULT IN EXPLOSION OR FIRE OF GREAT INTENSITY WITH THE LIBERATION OF HAZARDOUS AND IRRITATING GASES. USE ONLY A CLEAN DRY CUP OR DISPENSER. DO NOT REUSE CONTAINER. RINSE EMPTY CONTAINER THOROUGHLY WITH WATER BEFORE DISCARDING.

ANTIDOTE

EXTERNAL — Wash thoroughly with water. For eyes get prompt medical attention.

INTERNAL — If swallowed, drink promptly a large quantity of milk, egg whites, gelatin solution or, if these are not available, drink large quantities of water. Avoid alcohol. Call physician immediately. (Note to physician: Mucosal damage may contraindicate the use of gastric lavage.)

100 LB.
NET WT.

ACCEPTED
FEB 27 1974

335-195

UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND RODENTICIDE ACT
FOR ECONOMIC POISON REGISTRATION
ED UNDER NO. 335-195 SUBJECT
TO ATTACHED COMMENTS

PENNSWIM

POOL-CLOR

Concentrated Chlorine for Swimming Pools

ALGAECIDE • BACTERICIDE • DISINFECTANT

ACTIVE INGREDIENT: Sodium Dichloro-s-Triazinetrione (Sodium Dichloroisocyanurate)
AVAILABLE CHLORINE 63%

DANGER: KEEP OUT OF REACH OF CHILDREN!
See Back Panel for Additional Precautions and Antidote Statements

PENNSWIM DEPARTMENT • PENN WALT CORPORATION • PHILADELPHIA, PA.



ACCEPTED
8 27 1974
35-195
THE FEDERAL INSECTICIDE
AND RODENTICIDE ACT
ECONOMIC POISON REGISTRATION
NO... CONTACT
RED COMMENTS.



RECEIVED
20 FEB 1974
FEDERAL BUREAU OF INVESTIGATION
WASHINGTON, D.C.

POOL-CLOR

Concentrated Chlorine for Swimming Pools

ALGAEICIDE • BACTERICIDE • DISINFECTANT

INGREDIENT: Sodium Dichloro-s-Triazinetrione (Sodium Dichloroisocyanurate) 100%
AVAILABLE CHLORINE 63%

DANGER: KEEP OUT OF REACH OF CHILDREN
See Back Panel for Additional Precautions and Antidote Statements

DIRECTIONS FOR USE

Pool-Clor Concentrated Chlorine is a fast dissolving granular powder for chlorination of swimming pool water.

For pool start-up, first adjust the pH of water to 7.2-7.6 with a proper additive such as Pennswim Aquet[®] or Dry Acid according to the instructions on their label, then add at least 4 oz. of Pool-Clor Concentrated Chlorine for each 10,000 gallons of water to satisfy the chlorine demand of the water and reach the initial chlorine residual of 0.6 ppm (stabilized pool required 1.0 ppm). For the stabilization of the chlorine in the pool, add Pennswim Pool Stabilizer (following the directions for use on its label) at the beginning of the pool season.

To prevent the growth of algae, superchlorinate at the start of the season, and then once every one or two weeks with PENNSWIM BIGSHOT, used according to the instructions on its label.

Continue with 2 oz.-3 oz. of POOL-CLOR Concentrated Chlorine per 10,000 gallons of water as necessary to maintain the minimum chlorine residuals. Health regulatory agencies usually require 0.6 to 1.0 ppm for non-stabilized and 1.0 to 1.5 ppm for stabilized pools, determined by daily or more frequent checks with a reliable chlorine test kit. pH of the pool water should be maintained in the 7.2-7.6 range as determined by daily checks with a test kit. Variations in temperature, sunlight, organic matter (leaves, grass, dust, etc.) and bathing loads will influence the rate of chlorine consumption and determine Pool-Clor Concentrated Chlorine dosages.

This product is toxic to fish. Treated effluent should not be discharged where it will drain into lakes, streams, ponds or public water.

Printed in U.S.A.

EPA Reg. No. 335-195

Made in U.S.A.

4-6650A-01

DEPARTMENT • PENN WALT CORPORATION • PHILA., PA. 19102