



# STERICIDE SANITIZER<sup>®</sup>

## Disinfectant Deodorizer Sanitizer

FOR INDUSTRIAL USE ONLY

NET CONTENTS 15455 GALS.

### DIRECTIONS FOR USE:

**HOUSEHOLD** For sanitizing and disinfecting floors, walls and inanimate hard surfaces. For Schools, Homes, Locker Rooms, Garbage Pails, Sink Tops, Corridors, Classrooms, Offices and Shower Stalls. Apply solution with mop or cloth. Sanitize with 1 ounce Stericide Sanitizer to 4 gallons of water. Disinfect with 3 ounces Stericide Sanitizer to 5 gallons of water. Kills Staphylococcus aureus and Salmonella choleraesuis.

**HOSPITALS** For floors, walls and other hard surfaces. In Nursing Homes, Institutions, as well as Sink Tops, Garbage Pails, Telephones and Restrooms, use Stericide Sanitizer at 3 1/2 ounces per 5 gallons of water. At this level Stericide Sanitizer will be effective against Pseudomonas aeruginosa (PRD) 10.

**FOOD PROCESSING EQUIPMENT** For sanitization of food processing equipment and food utensils dilute 1 ounce Stericide Sanitizer per 4 gallons of water to provide 200 ppm of active quaternary. At this level, No Potable Water Rinse is required.

**DAIRIES** To sanitize dairy equipment, first clean and rinse the equipment thoroughly. Then apply sanitizing solution containing 1 ounce Stericide Sanitizer to 4 gallons of water (200 ppm). At this level, No Potable Water Rinse is required. Follow recommendations of local Health Board.

**MEAT AND POULTRY PLANTS** For sanitizing cleaned food processing equipment or utensils in Federally inspected meat and poultry processing plants dilute 1 ounce Stericide Sanitizer per 4 gallons of water (200 ppm). At this concentration, No Potable Water Rinse is required after thorough draining of the sanitizing solution.

**COLD DISINFECTION** For disinfection of previously cleaned surgical instruments, barber tools and dental equipment, simply submerge instruments into solution containing 1 1/2 ounce Stericide Sanitizer per gallon of water.

**RESTAURANT AND BAR RINSE** Dishes, glassware, silverware, cooking utensils, wash with soap or synthetic detergent, rinse thoroughly, and immerse in a sanitizing solution containing 1 ounce Stericide Sanitizer to 4 gallons of water. No Potable Water Rinse Required.

**ACTIVE INGREDIENTS:** n-Alkyl (60% C-30 - C-5) - C-5 - C) dimethyl benzyl ammonium chloride 5%  
n-Alkyl (68% C-32 - C-18) methyl ethyl benzyl ammonium chloride 5%  
**INERT INGREDIENTS:** 90%

**DANGER:** KEEP OUT OF REACH OF CHILDREN. CORROSIVE. CAUSES EYE DAMAGE AND SKIN IRRITATION. Do not get in eyes, skin or on clothing. Protect eyes and skin when handling. Harmful if swallowed. Avoid contamination of food, drink, or a concentrate and should be diluted before using.

**FIRST AID:** In case of contact, immediately flush eyes or skin with plenty of water for at least 15 minutes. For eyes, call a physician. Remove and wash contaminated clothing before reuse. If swallowed, drink promptly a large quantity of milk and water or great quantities of water if these are not available. Drink large quantities of water. Call a physician immediately.

**NOTE TO PHYSICIAN:** Potable water contamination may contraindicate use of gastric lavage. Mercuric agents, in case of shock, hypotension, hypopressure, and convulsions may be needed.

Rinse empty container thoroughly with water and discard.

PENNCHEM DEPARTMENT • PENNWALT CORPORATION • PHILADELPHIA, PENNSYLVANIA 19102

PCI 52917 - Robert Krain  
8-18-75 - mtr  
Pennwalt Stericide Sanitizer

Produced from  
3 3/4" x 20"