

BEST DOCUMENT AVAILABLE



STERICIDE™ SANITIZER



DIRECTIONS FOR USE:

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

FOOD PROCESSING EQUIPMENT—Clean equipment with a good detergent and follow with water rinse prior to sanitizing. For sanitization of pre-cleaned food processing equipment and utensils dilute 1 ounce Stericide Sanitizer per 4 gallons of water to provide 200 ppm of active quaternary for 1 minute contact time. At this level No Potable Water Rinse is required providing the sanitizing solution is adequately drained. A fresh sanitizing solution should be prepared daily for mechanical as well as for manual applications and should not be re-used except for cleaning or pre-rinse of dirty equipment.

DAIRY AND MUSHROOM FARMS—To sanitize dairy or mushroom equipment first clean and rinse equipment thoroughly. Then apply sanitizing solution containing 1 ounce Stericide Sanitizer to 4 gallons of water (200 ppm). At this level No Potable Water Rinse is required after complete draining. The Stericide Sanitizer can also be used to disinfect floors, walls and inanimate hard surfaces on the mushroom premises or in areas where compost and mushrooms are not present.

MEAT AND POULTRY PLANTS—For sanitizing cleaned food processing equipment or utensils in Federally inspected meat and poultry processing plants, dilute 1 ounce Stericide Sanitizer per 4 gallons of water (200 ppm). At this concentration, No Potable Water Rinse is required after thorough draining of the sanitizing solution.

COLD DISINFECTION—For disinfection of previously cleaned surgical instruments, barber tools and dental equipment, simply submerge instruments into solution containing 1 ounce Stericide Sanitizer per gallon of water for 10 minutes. Prepare a fresh solution daily or when use solution becomes visibly dirty.

RESTAURANT AND BAR RINSE—Dishes, glassware, silverware, cooking utensils, wash with soap or synthetic detergent, rinse thoroughly and immerse in a sanitizing solution containing 1 ounce Stericide Sanitizer to 4 gallons of water. No Potable Water Rinse Required if drained adequately.

Disinfectant Deodorizer Sanitizer

ACTIVE INGREDIENTS: n-Alkyl (60% C₁₂, 30% C₁₄, 5% C₁₆, 5% C₁₈)
dimethyl benzyl ammonium chlorides 5%
n-Alkyl (68% C₁₂, 32% C₁₄) dimethyl ethylbenzyl ammonium chlorides 5%
INERT INGREDIENTS: 90%

KEEP OUT OF REACH OF CHILDREN
DANGER

CORROSIVE CAUSES EYE DAMAGE AND SKIN IRRITATION Do not get in eyes, skin or clothing. Protect eyes and skin when handling. Harmful if swallowed. Avoid contamination of food. Stericide is a concentrate and should be diluted before using. (See side panel for additional precautionary statements)

Net Contents _____ Gals. _____ L

FOR INDUSTRIAL USE ONLY

ONLY FOR SALE TO, USE, AND STORAGE BY SERVICE PERSONS

4 97141 04 A4

Made and Printed in U.S.A.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

HOSPITALS—For floors, walls and other hard nonporous surfaces. In Nursing Institutions, as well as Sink Tops, Garbage Pails, Telephones and Restrooms, use Sanitizer at 3 1/2 ounces per 5 gallons of water. At this level Stericide Sanitizer is effective against Pseudomonas aeruginosa PRD-10. Remove gross filth or dirt. Pre-clean all surfaces. Apply with a cloth or mop. Thoroughly wet all surface surfaces must remain wet for 10 minutes. Fresh solution should be prepared daily. The use solution becomes visibly dirty.

STATEMENT OF PRACTICAL TREATMENT: In case of contact, immediately flush or skin with plenty of water for at least 15 minutes. For eyes, call a physician and wash contaminated clothing before re-use. If swallowed, do not induce vomit; promptly a large quantity of milk, egg whites, gelatin solution or if these are not available, drink large quantities of water. Call physician immediately.

NOTE TO PHYSICIAN: Probable mucosal damage may contraindicate the use of lavage. Measure against circulatory shock, respiratory depression and comatose needed.

Rinse empty container thoroughly with water and discard.

SPILL OR LEAK: In case of spill, flush up area with water spray.

STORAGE AND DISPOSAL: Precautions: Do not contaminate water, food or feed by storage or disposal. Open dumping is prohibited. Do not reuse empty container. **Pesticide Disposal:** Pesticide spray mixture, or residue that cannot be used or processed should be disposed in a landfill approved for pesticides away from supplies.

Container Disposal: Triple rinse with water and dispose of in an approved landfills. General: Consult federal, state or local disposal authorities for approved alternatives.

EPA REG. NO. 335-202

EPA EST. NO. 335-OH-1



BK DEPARTMENT • PENNWALT CORPORATION • PHILADELPHIA, PENNSYLVANIA 19102

ACCEPTED
with COMMENTS
in EPA Letter Dated:

APR 26 1984

APR 13 1984

JOB 137
Pennwalt—Stericide Sanitizer

Under the Federal Insecticide, Fungicide, and Rodenticide Act
as amended, this product is
registered under EPA Reg. No.
335-202

Registration Division (TS-767)

BEST DOCUMENT AVAILABLE

STERICIDE™ SANITIZER



Disinfectant Deodorizer Sanitizer

ACTIVE INGREDIENTS: n-Alkyl (60% C₁₂, 30% C₁₄, 5% C₁₂, 5% C₁₆)
dimethyl benzyl ammonium chlorides 5%
n-Alkyl (68% C₁₂, 32% C₁₄) dimethyl ethylbenzyl ammonium chlorides 5%
INERT INGREDIENTS: 90%

KEEP OUT OF REACH OF CHILDREN
DANGER

CORROSIVE CAUSES EYE DAMAGE AND SKIN IRRITATION Do not get in eyes, skin or clothing. Protect eyes and skin when handling. Harmful if swallowed. Avoid contamination of food. Stericide is a concentrate and should be diluted before using. (See side panel for additional precautionary statements)

Net Contents _____ Gals. _____ L

FOR INDUSTRIAL USE ONLY

ONLY FOR SALE TO, USE, AND STORAGE BY SERVICEPERSONS

4-93498-04 A4

Made and Printed in U.S.A.

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

HOSPITALS—For floors, walls and other hard nonporous surfaces. In Nursing Home Institutions, as well as Sinks, Garbage Pails, Telephones and Restrooms, use Stericide Sanitizer at 3 ounces per 5 gallons of water. At this level Stericide Sanitizer is effective against *Pseudomonas aeruginosa* PHD 10. Remove gross filth or heavy Preclean all surfaces. Apply with a cloth or mop. Thoroughly wet all surfaces. Surfaces must remain wet for 10 minutes. Fresh solution should be prepared daily or the use solution becomes visibly dry.

STATEMENT OF PRACTICAL TREATMENT: In case of contact, immediately flush or skin with plenty of water for at least 15 minutes. For eyes, call a physician. He and wash contaminated clothing before re-use. If swallowed, do not induce vomiting. promptly a large quantity of milk, egg whites, goat milk solution or if these are not available, drink large quantities of water. Call physician immediately.

NOTE TO PHYSICIAN: Probable mucosal damage may contraindicate the use of g lavage. Measure against circulatory shock, respiratory depression and convulsions if needed.

Rinse empty container thoroughly with water and discard.

SPILL OR LEAK: In case of spill, flush spill area with water spray.

STORAGE AND DISPOSAL: Prohibitions: Do not contaminate water, for feed by storage or disposal. Open dumping is prohibited. Do not reuse empty container.

Pesticide Disposal: Pesticide spray mixture or residue that cannot be used or then processed should be disposed in a landfill approved for pesticides away from supplies.

Container Disposal: Triple rinse with water and dispose of in an approved landfill.

General: Consult federal, state or local disposal authorities for approved alternate procedures.

EPA REG NO 335-202

EPA EST NO 335-OH-1



PENNWALT CORPORATION • PHILADELPHIA, PENNSYLVANIA 19102

ACCEPTED
with COMMENTS
to EPA Letter Dated:

APR 26 1984

JOB 137
Pennwalt—Stericide Sanitizer 1-2

APR 13 1984

Under the Fungicide, and Herbicide Act
as amended, for the pesticide
registered under EPA Reg. No.

335-202

Registration Division (TS-767)

26 APR 1984

Pennwalt Corporation
3 Parkway
Philadelphia, PA 19102

Attention: G. R. Dychala
Group Leader

Gentlemen:

Subject: Stericide[®] Sanitizer
EPA Registration No. 335-202
Your Submission dated April 12, 1984

The amendment referred to above, submitted in connection with registration under FIFRA, is acceptable provided that you:

1. Make the labeling changes listed below before you release the product for shipment bearing the amended labeling:
 - a. Revise the statement (left panel):
 ...hard surfaces on the mushroom premises or in areas where compost and mushrooms are not present.
 to read:
 ...hard surfaces on the mushroom premises only in areas where compost and mushrooms are not present.
2. Submit five (5) copies of your final printed labeling before you release the product for shipment. Refer to the A-79 Enclosure for a further description of final printed labeling.

Your release for shipment of the product bearing the amended labeling constitutes acceptance of these conditions.

CONCURRENCES							
SYMBOL ▶							
SURNAME ▶							
DATE ▶							

335-202

-2-

A stamped copy of the labeling is enclosed for your records.

Sincerely yours,



John H. Lee
Product Manager (31)
Disinfectants Branch
Registration Division (TS-767C)

Enclosure

RD:DIS:DCR-04915;J.Lee:cml:RAVEN:557-2226;RD-53:4/20/84;Del.5/4/84
REVISED:DCR-04933;J.Lee:eg:Raven:557-2226;RD-53:4/25/84;Del.5/4/84