



335-209

CALED COOLING WATER TREATMENT

RECIRCULATING WATER COOLING TOWER ALGAEICIDE

Concentrated algaecide for industrial re-circulating water cooling towers

Helps keep water cooling tower systems clear and free of visible algae slimes. Inhibits the growth of bacteria.

ACTIVE INGREDIENTS:

n-Alkyl (60%C ₁₄ , 30%C ₁₆ , 5%C ₁₂ , 5%C ₁₈) dimethyl benzyl ammonium chlorides	5%
n-Alkyl (68%C ₁₂ , 32%C ₁₄) dimethyl ethylbenzyl ammonium chlorides	5%

INERT INGREDIENTS:	90%
	<hr/> 100%

NET CONTENTS _____ EPA Reg. No. 10666-2

DANGER

Keep out of Reach of Children

See other precautionary statements on right side panel.

CALED DEPARTMENT

 BRENTWOOD, MD. 20722

When Caled Cooling Treatment is used as directed, it will aid in controlling and/or preventing the development of unsanitary slime conditions which are unsightly, produce foul odors and which are often the cause of equipment breakdown.

Caled Cooling Treatment is compatible with virtually all corrosion inhibition additives, but should not be used in the presence of soap or other anionic detergents. Caled Cooling Treatment at use dilution is non-corrosive to inanimate hard surfaces.

ACCEPTED

3 28 1973

FEDERAL INSECTICIDE FUNGICIDE AND RODENTICIDE ACT FOR ECONOMIC POISON REGISTERED UNDER NO. 10666-2 SUBJECT TO ATTACHED COMMENTS

DANGER

Keep out of Reach of Children

Causes severe eye and skin damage. Do not get in eyes, on clothing. Wear goggles or face shield, and rubber gloves when handling. Harmful or fatal if swallowed. Avoid contamination of food.

FIRST AID

In case of contact, immediately flush eyes or skin with plenty of water for at least 15 minutes. For eyes, call a physician. Remove and wash all contaminated clothing before reuse. If swallowed, drink milk, egg whites, gelatin solution, or if these are not available, drink large quantities of water. Call a physician.

Rinse empty container thoroughly with water and discard.

NOTE TO PHYSICIAN

Probable mucosal damage may contraindicate the use of gastric lavage. Measures against circulatory shock, respiratory depression and convulsion may be needed.