



PENNCHEM

Aug 27, 1975

ACCOMPLISHTM

PAT. PENDING

Low Foam Concentrated Acid-Anionic Surfactant Sanitizer

DIRECTIONS FOR USE

DIRECTIONS FOR USE: To prepare a sanitizing solution, add one fluid ounce of ACCOMPLISH to 2 gallons of water which will provide 200 ppm of available sanitizer as determined by the use of Pennsan XXX Test Kit according to its instructions (Modified for Accomplish). Allow 2 minutes for exposure time. Discard sanitizing solution when test indicates 100 ppm.

DANGER: CORROSIVE TO TISSUES. CAUSES EYE DAMAGE AND SKIN IRRITATION. DO NOT GET IN EYES, ON SKIN OR ON CLOTHING. WEAR GOGGLES OR FACE SHIELD, AND RUBBER GLOVES WHEN HANDLING. HARMFUL OR FATAL IF SWALLOWED. AVOID CONTAMINATION OF FOOD. AVOID BREATHING SPRAY OR FOGGING MIST.

FIRST AID: In case of contact, immediately flush eyes or skin with plenty of water for at least 15 minutes. For eyes, call a physician. Remove and wash contaminated clothing before reuse.

If swallowed, drink promptly a large quantity of milk, egg whites, gelatin solution or if these are not available, drink large quantities of water. Call a physician immediately.

NOTE TO PHYSICIAN: Probable mucosal damage may contraindicate the use of gastric lavage. Measures against circulatory shock, respiratory depression and convulsion may be needed.

ACTIVE INGREDIENTS:	
Phosphoric Acid	37.50%
Dodecylbenzene Sulfonic Acid	4.85%
Sodium Mono (1-alkenyl*) Phenoxybenzene Disulfonate	3.15%
Sodium Di (1-alkenyl*) Phenoxybenzene Disulfonate	1.35%
Sodium Methyl and Dimethyl Naphthalene Sulfonate	5.00%
INERT INGREDIENTS:	48.15%

*C₉₋₁₀

DANGER:

Keep Out of Reach of Children.

Read First Aid and Additional
Precautions on Side Panel.

EPA Reg. No. 335-211
EPA EST. NO. 335-IL-1

*Rinse empty container thoroughly
with water and discard it.*

DAIRY AND FOOD PLANTS

TANK TRUCK, HOLDING TANKS, SILOS, VATS AND PLANT EQUIPMENT: Sanitize pre-cleaned equipment with ACCOMPLISH solution (1 oz./2 gallons of water). Drain ACCOMPLISH solution from equipment.*

CIP LINES—CLOSED SYSTEM: Flush entire system with tepid water until water runs clear, then sanitize with ACCOMPLISH (1 oz./2 gal. of water) by circulating solution through lines. Drain sanitizing solution from the system.*

BEVERAGE PLANTS

PRE-MIX AND POST-MIX VENDING MACHINES, PLANT EQUIPMENT: Sanitize pre-cleaned surfaces with ACCOMPLISH solution (1 oz./2 gallons of water). Drain sanitizing solution from the system.*

EATING AND DRINKING ESTABLISHMENTS

Sanitize pre-cleaned utensils and glassware with ACCOMPLISH (1 oz./2 gallons of water). Drain sanitizing solution from utensils and glassware.*

DAIRY FARMS

BULK MILK COOLING TANKS: Prior to filling the tank with milk, sanitize clean tank with ACCOMPLISH solution (1 oz./2 gallons of water) by brush, circulation or spray method. Drain sanitizing solution from tank.*

FARM DAIRY UTENSILS: Prior to milking, sanitize clean utensils with ACCOMPLISH solution (1 oz./2 gallons of water) by rinse or spray.*

PIPELINE MILKERS: After cleaning by recommended procedure, rinse with warm water and sanitize with ACCOMPLISH solution (1 oz./2 gallons of water).*

UDDER SANITATION: Wash and massage udders before milking with solution of 1 oz. of ACCOMPLISH per 2 gallons of warm water (110°-115°F.) Individual disposable paper towels for each cow should be used. After milking, dip each teat in dip cup containing ACCOMPLISH solution (1 oz. per 2 gallons of water). Never use the same solution for more than 3-4 animals. Replace solution if it becomes cloudy or visibly dirty. Never dip used towel back in the sanitizing solution.

INFLATIONS: Brush inflations after milking with warm (110°-120°F.) ACCOMPLISH solution (1 oz./2 gallons of water) using good inflation brush. Rinse thoroughly and store as directed by sanitarians or field men. Prior to milking, sanitize with ACCOMPLISH solution (1 oz./2 gal-

ACCEPTED

PENNCHEM

335-211

11-27-1975

ACCOMPLISH™

PAT. PENDING

Low Foam Concentrated Acid-Anionic Surfactant Sanitizer

DIRECTIONS FOR USE

DIRECTIONS FOR USE: To prepare a sanitizing solution, add one fluid ounce of ACCOMPLISH to 2 gallons of water which will provide 200 ppm of available sanitizer as determined by the use of Pennsan XXX Test Kit according to its instructions (Modified for Accomplish). Allow 2 minutes for exposure time. Discard sanitizing solution when test indicates 100 ppm.

DANGER: CORROSIVE TO TISSUES. CAUSES EYE DAMAGE AND SKIN IRRITATION. DO NOT GET IN EYES, ON SKIN OR ON CLOTHING. WEAR GOGGLES OR FACE SHIELD, AND RUBBER GLOVES WHEN HANDLING. HARMFUL OR FATAL IF SWALLOWED. AVOID CONTAMINATION OF FOOD. AVOID BREATHING SPRAY OR FOGGING MIST.

FIRST AID: In case of contact, immediately flush eyes or skin with plenty of water for at least 15 minutes. For eyes, call a physician. Remove and wash contaminated clothing before reuse.

If swallowed, drink promptly a large quantity of milk, egg whites, gelatin solution or if these are not available, drink large quantities of water. Call a physician immediately.

NOTE TO PHYSICIAN: Probable mucosal damage may contraindicate the use of gastric lavage. Measures against circulatory shock, respiratory depression and convulsion may be needed.

ACTIVE INGREDIENTS:

Phosphoric Acid	37.50%
Dodecylbenzene Sulfonic Acid	4.85%
Sodium Mono (1-alkenyl*)	
Phenoxybenzene Disulfonate	3.15%
Sodium Di (1-alkenyl*)	
Phenoxybenzene Disulfonate	1.35%
Sodium Methyl and Dimethyl	
Naphthalene Sulfonate	5.00%

INERT INGREDIENTS: 48.15%

*C₉₋₁₀

DANGER:

Keep Out of Reach of Children.

Read First Aid and Additional Precautions on Side Panel.

EPA Reg. No. 335-211

EPA EST. NO. 335-IL-1

Rinse empty container thoroughly with water and discard it.

DAIRY AND FOOD PLANTS

TANK TRUCK, HOLDING TANKS, SILOS, VATS AND PLANT EQUIPMENT: Sanitize pre-cleaned equipment with ACCOMPLISH solution (1 oz./2 gallons of water). Drain ACCOMPLISH solution from equipment.*

CIP LINES—CLOSED SYSTEM: Flush entire system with tepid water until water runs clear, then sanitize with ACCOMPLISH (1 oz./2 gal. of water) by circulating solution through lines. Drain sanitizing solution from the system.*

BEVERAGE PLANTS

PRE-MIX AND POST-MIX VENDING MACHINES, PLANT EQUIPMENT: Sanitize pre-cleaned surfaces with ACCOMPLISH solution (1 oz./2 gallons of water). Drain sanitizing solution from the system.*

EATING AND DRINKING ESTABLISHMENTS

Sanitize pre-cleaned utensils and glassware with ACCOMPLISH (1 oz./2 gallons of water). Drain sanitizing solution from utensils and glassware.*

DAIRY FARMS

BULK MILK COOLING TANKS: Prior to filling the tank with milk, sanitize clean tank with ACCOMPLISH solution (1 oz./2 gallons of water) by brush, circulation or spray method. Drain sanitizing solution from tank.*

FARM DAIRY UTENSILS: Prior to milking, sanitize clean utensils with ACCOMPLISH solution (1 oz./2 gallons of water) by rinse or spray.*

PIPELINE MILKERS: After cleaning by recommended procedure, rinse with warm water and sanitize with ACCOMPLISH solution (1 oz./2 gallons of water).*

UDDER SANITATION: Wash and massage udders before milking with solution of 1 oz. of ACCOMPLISH per 2 gallons of warm water (110 -115 F.) Individual disposable paper towels for each cow should be used. After milking, dip each teat in dip cup containing ACCOMPLISH solution (1 oz. per 2 gallons of water). Never use the same solution for more than 3-4 animals. Replace solution if it becomes cloudy or visibly dirty. Never dip used towel back in the sanitizing solution.

INFLATIONS: Brush inflations after milking with warm (110 -120 F.) ACCOMPLISH solution (1 oz./2 gallons of water) using good inflation brush. Rinse thoroughly and store as directed by sanitarians or field men. Prior to milking, sanitize with ACCOMPLISH solution (1 oz./2 gal. of water).*