

DIRECTIONS

**WHERE TO USE
BARRENIZE**

BARRENIZE should be used only in places where the complete control of vegetation is desired. It is non-selective! Treated areas may remain free of vegetation for a year or longer. Use BARRENIZE wherever weeds create a fire-hazard, a maintenance problem, a nuisance or an unsightly appearance. It is successfully used about Industrial Sites, Lumber Yards, Substations and Pole Yards, Petroleum Installations, Warehouses, Storage Yards, Highway Markers, Guard Rails, Bridge Abutments, Rail Sidings, or wherever a weed-free surface is desired.

HOW TO APPLY

BARRENIZE is applied DRY. It is a granular, dust-free material—ready for use when you receive it. Any mechanical spreader capable of delivering the low rates prescribed, may be used. Spot treatments can be made by hand. The most simple and efficient method of application is to use the special PCB Spreader. We strongly recommend using this hand-operated Spreader as it is specifically designed for the even and uniform application of BARRENIZE.

WHEN TO APPLY

Fall, winter, or early spring applications are recommended for maximum efficiency. Late spring and summer applications are not recommended for areas where summer rainfall is normally sufficient to dissolve and activate the material. BARRENIZE kills plants through their root systems. (Rainfall normally is required to carry it to the roots.) Applications made during dry seasons may be activated by thoroughly wetting the soil after treatment.



ACC
FEB 2 1967
337-18
FEDERAL INSPECTION
FEDERAL AND ROYALTY ACT
FEDERAL POISON REGISTRY
FEDERAL NO. SUBJECT
TO ATTACHE COMMENTS

BEST DOCUMENT AVAILABLE

BARRENIZE

CAUTION: Keep out of the reach of children
SOIL TOXICANT

**NON-SELECTIVE WEED KILLING COMPOUND
FOR DRY APPLICATION
EASY TO USE**

ACTIVE INGREDIENTS	
Barrenize (active ingredient) (N.B. 100%)	95
Inert (Treated with Dieldrin) (N.B. 100%)	5
Manufactured by (N.B. 100%)	4
Manufactured by (N.B. 100%)	2

NET CONTENTS POUNDS
SOLD BY

Lester Laboratories
INCORPORATED
ATLANTA, GEORGIA
PLANT: EAST POINT, GEORGIA

**TO CONTROL
ANNUAL WEEDS**

Apply 1/2 to 1 pound of BARRENIZE per 100 sq. ft. of soil surface. Applications should be made before seedlings appear or when young plants are tender and most susceptible to chemical action. BARRENIZE gives non-selective control for an extended period. Length of residual action is dependent on rainfall and type of soil. In arid regions, applications should be made just prior to rainy season.

**TO CONTROL
PERENNIAL WEEDS**

Apply 1 to 2 pounds of BARRENIZE per 100 sq. ft. during the fall, winter, or early spring months. Fall applications are preferable for deep-rooted perennials. Late spring and summer applications are not recommended in areas where summer rainfall may be insufficient to dissolve the material.

IMPORTANT

This material is not selective in action and may be toxic to all types of vegetation and may render the entire treated area totally or partially unproductive for one or more years. Care should be taken to confine the use or application to the particular area intended to be treated and to avoid its contact with lawns, trees, shrubs, crops and other desirable plants which are not intended to be destroyed or injured. This includes precautions in treating areas which may be underlaid by roots of adjacent valuable growths. Careless application of this material—washing by water run-off, for example, where desirable plants are growing or which will be used for later planting may result in injury to desirable plants. Avoid contamination of irrigation waters. Mechanical equipment should be thoroughly cleaned after use.

CAUTION

May cause irritation of eyes, nose, throat and skin. Avoid contact with skin, eyes and clothing. Avoid contamination of feed and foodstuffs. Keep away from children and domestic animals.

DIRECTIONS

WHERE TO USE BARRENIZE

BARRENIZE should be used only in places where the complete control of vegetation is desired. It is non-selective! Treated areas may remain free of vegetation for a year or longer. Use BARRENIZE wherever weeds create a fire hazard, a maintenance problem, a nuisance or an unsightly appearance. It is successfully used about Industrial Sites, Lumber Yards, Substations and Pole Yards, Petroleum Installations, Warehouses, Storage Yards, Highway Markers, Guard Rails, Bridge Abutments, Rail Sidings or wherever a weed free surface is desired.

HOW TO APPLY

BARRENIZE is applied DRY. It is a granular, dust-free material—ready for use when you receive it. Any mechanical spreader capable of delivering the low rates prescribed, may be used. Seed treatments can be made by hand. The most simple and efficient method of application is to use the special PCB Spreader. We strongly recommend using this hand operated Spreader as it is specifically designed for the even and uniform application of BARRENIZE.

WHEN TO APPLY

Fall, winter or early spring applications are recommended for maximum efficiency. Late spring and summer applications are not recommended for areas where summer rainfall is normally sufficient to dissolve and activate the material. BARRENIZE kills plants through their root system. (Rainfall normally is required to carry it to the root zone.) Applications made during dry seasons may be activated by thoroughly wetting the soil after treatment.



ACC
FEB 2 1967
337-18
FEDERAL INSECTICIDE
AND RODENTICIDE ACT
FOR REG. ORIC. POISON REG. SYM-
BUL. 10. SUBJECT
TO ATTACH. COMMENTS.

QUALITY PURITY
CHEMICAL INTEGRITY

BEST DOCUMENT AVAILABLE

BARRENIZE

CAUTION: Keep out of the reach of children
SOIL TOXICANT

**NON-SELECTIVE WEED KILLING COMPOUND
FOR DRY APPLICATION
EASY TO USE**

ACTIVE INGREDIENT
100% BARRENIZE
NET CONTENTS: 10 POUNDS
SOLD BY

NET CONTENTS POUNDS
SOLD BY

Laboratories
INCORPORATED
ATLANTA, GEORGIA
PLANT: EAST POINT, GEORGIA

TO CONTROL ANNUAL WEEDS

Apply 1/2 to 1 pound of BARRENIZE per 100 sq. ft. of soil surface. Applications should be made before seedlings appear or when young plants are tender and most susceptible to chemical action. BARRENIZE gives non-selective control for an extended period. Length of residual action is dependent on rainfall and type of soil. In arid regions applications should be made just prior to rainy season.

TO CONTROL PERENNIAL WEEDS

Apply 1 to 2 pounds of BARRENIZE per 100 sq. ft. during the fall, winter or early spring months. Fall applications are preferable for deep-rooted perennials. Late spring and summer applications are not recommended in areas where summer rainfall may be insufficient to dissolve the material.

IMPORTANT

This material is not selective in action and may be toxic to all types of vegetation and may render the entire treated area totally or partially unproductive for one or more years. Care should be taken to confine the use or application to the particular area intended to be treated and to avoid its contact with lawns, trees, shrubs, crops and other desirable plants which are not intended to be destroyed or injured. This includes precautions in treating areas which may be underlaid by roots of adjacent valuable growths. Careless application of this material or washing by water run-off, to areas where desirable plants are growing or which will be used for later planting may result in injury to desirable plants. Avoid contamination of irrigation waters. Mechanical equipment should be thoroughly cleaned after use.

CAUTION

May cause irritation of eyes, nose, throat and skin. Avoid contact with skin, eyes and clothing. Avoid contamination of feed and foodstuffs. Keep away from children and domestic animals.

DIRECTIONS

WHERE TO USE BARRENIZE

BARRENIZE should be used only in places where the complete control of vegetation is desired. It is non-selective! Treated areas may remain free of vegetation for a year or longer. Use BARRENIZE wherever weeds create a fire-hazard, a maintenance problem, a nuisance or an unsightly appearance. It is successfully used about Industrial Sites, Lumber Yards, Substations and Pole Yards, Petroleum Installations, Warehouses, Storage Yards, Highway Markers, Guard Rails, Bridge Abutments, Rail Sidings, or wherever a weed-free surface is desired.

HOW TO APPLY

BARRENIZE is applied DRY. It is a granular, dust-free material—ready for use when you receive it. Any mechanical spreader, capable of delivering the low rates prescribed, may be used. Spot treatments can be made by hand. The most simple and efficient method of application is to use the special PCB Spreader. We strongly recommend using this hand-operated Spreader as it is specifically designed for the even and uniform application of BARRENIZE.

WHEN TO APPLY

Fall, winter or early spring applications are recommended for maximum efficiency. Late spring and summer applications are not recommended for areas where summer rainfall is normally insufficient to dissolve and activate the material. BARRENIZE kills plants through their root systems. (Rainfall normally is required to carry it to the root zone.) Applications made during dry seasons may be activated by thoroughly wetting the soil after treatment.



BARRENIZE

CAUTION: Keep out of the

SOIL TOXIC

NON-SELECTIVE WEED KILLER

FOR DRY APPLICATIONS

EASY TO USE

ACTIVE INGREDIENTS

Disodium Tetraborate Pentahydrate
Disodium Tetraborate Dihydrate
Monuron (3-(4-Chlorophenyl)-1,1-Dimethylurea)
(Boron Trioxide (B₂O₃)) (equivalent)
Inert ingredients

NET CONTENTS

SOLD BY

Water-Land
INCORPORATED
ATLANTA
PLANT: EAST POINT