

Panel #1, Left

DIRECTIONS FOR USE

GENERAL: Apply with a low pressure sprayer equipped with a coarse spray nozzle. Spray surfaces only until wet. Avoid excessive run-off.

COCKROACHES: For cockroach control, apply to all insect breeding sites such as baseboards; cracks and crevices; around window and door frames; the under surfaces of shelves, drawers and work tables; behind and beneath cabinets, refrigerators, sinks and stoves, in and around waste containers, switch panels, junction boxes, floor drains and other places where cockroaches hide or nest. Spot treat the infested area, as well as cracks and crevices.

ANTS: Apply to trails and places where ants enter premises such as window frames and door sills. Spray hills and runways outdoors.

CADELLS, CIGARETTE BEETLES, DERMESTIDS, DRUGSTORE BEETLES, INDIAN MEAL MOTHS, MEDITERRANEAN FLOUR MOTHS, RED FLOUR BEETLES, SAW-TOOTHED GRAIN BEETLES AND WEEVILS (Exposed Stages): Thoroughly treat floors and the walls of the infested area with attention to cracks, crevices, corners and perimeters. Sanitation and dust control are of primary importance. Fumigate or destroy infested products.

SILVERFISH, FIREBRATS and PAPERLICH: Apply to localized areas infested by these insects.

BROWN DOG TICKS: Spray sleeping quarters of pets, around baseboards, window and door frames, wall cracks, and local areas of floors. Repeat as needed. Fresh bedding should be placed in animal quarters after spraying. Do not spray animals.


ACCEPTED WITH COMMENTS

CLOVER MITES, CRICKETS, LARVAE, MILLIPEDES, CENTIPEDES, SOWBUGS, SCORPIONS AND SPIDERS: Thoroughly spray the interior and exterior perimeter of the infested premises. For clover mites, give special attention to window frames.

FLEAS, FLIES, GNATS, MOSQUITOES, PUZZLES AND SANDFLIES (Outdoors Only): Spray infested areas thoroughly including outside surfaces of screens, doors, window frames, foundation, patios and other areas where insects may enter.

ACCEPTED

Panel #2, Center



LESTER



DEAD END

RESIDUAL SPRAY
contains BAYGON

FOR INDUSTRIAL AND INSTITUTIONAL USE ONLY

DEAD END Residual Spray is a highly effective ready to use formulation for control of the insects listed hereon, including those resistant to certain chlorinated hydrocarbon and phosphate insecticides. DEAD END drives insects from their hiding places, gives rapid kill and long residual activity.

ACTIVE INGREDIENTS:

o-Isopropoxyphenyl Methylcarbamate*	1.00%
2-Butoxyethanol	8.94%
Petroleum Distillate	74.75%

INERT INGREDIENTS	15.31%
-------------------	--------

*U.S. Patent No. 5,111, 79. Material used under license from Chemagro Corporation.


CAUTION: KEEP OUT OF REACH OF CHILDREN.
See back panel for additional cautions.

NET CONTENTS

GALLONS

FORMERLY

1971



FOR USE IN MOTELS, HOTELS, THEATRES, AND OFFICES.

See under DIRECTIONS FOR USE for control of particular insect infestation.

FOR USE IN FOOD PROCESSING PLANTS SUCH AS FLOUR MILLS, BAKERIES, CANNERIES, AND MEAT PROCESSING PLANTS: Do not use in edible products areas of food processing plants, restaurants, or other areas where food is commercially prepared or processed. Do not use in serving areas while food is exposed. Treat food processing plants early in the season before canning or other processing begins. Repeat applications, if necessary, when plant is not in operation. When using the spray in the edible product areas where no exposed food or processing equipment is present, take any needed precautions to avoid any contamination of packaged or stored food commodities. Rinse all equipment with water before resuming operations. Spraying unloading docks and outside areas where insects congregate will aid in preventing re-infestation. Good sanitation and proper disposal of wastes is essential in any effective insect control program.

RESTRICTIONS

Do not use as a space spray. Provide adequate ventilation of area being treated. Do not allow children to contact treated areas until surfaces are thoroughly dry. Do not store near feed or food products. Avoid contamination of feed, food processing equipment and food handling surfaces. Do not apply to animals. Remove pets and cover fish bowls before spraying. Do not spray plants.

Do not reuse empty container. Destroy it by perforating or crushing. Bury or discard in a safe place away from water supplies.

CAUTION

Avoid breathing spray mist. Wash thoroughly after handling. May be harmful if swallowed, inhaled or absorbed through the skin. Do not get in eyes, on skin, or on clothing. In case of contact, wash immediately with soap and warm water. Wash hands, arms and face thoroughly with soap and water before eating or smoking. Wash all contaminated clothing with soap and hot water before re use. Should poisoning occur, get prompt medical aid.

To Physician: At least 10 minutes before treatment.