

NOV 16 1988

Gentlemen:

The submission referred to above, submitted in connection with the Label Improvement Program Notice for Antimicrobial Products Used as Food Contact Surface Sanitizers, is acceptable provided that you:

1. Make the labeling changes listed below before you release the product for shipment bearing the amended labeling.
 - a. Include the two subheadings Pesticide Disposal and Container Disposal in the Storage and Disposal section.
 - b. On the left panel, above the word Danger, add the following statements:

Hazard to Humans and Domestic Animals
Keep Out of Reach of Children

- c. The term "First Aid" must be changed to "Statement of Practical Treatment."
- d. Place the "Directions For Use" heading and the misuse statement directly above the actual Directions For Use section.
- e. Include the 10-minute contact time for the disinfection section.

50318; I:DeLaney; LH-10; RENCO: 10/25/88; 12/7/88; CT: vcr:rw

CONCURRENCES								
SYMBOL								
SURNAME								
DATE								


- f. Revise the referral statement to read "See side panel for additional precautionary statements."
 - g. The statement "Keep Out of Reach of Children" must be placed directly above the signal word "DANGER."
 - h. Delete the word "milk."
2. Submit five (5) copies of your final printed labeling before you release the product for shipment.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with section 6(e) of the Federal Insecticide, Fungicide, and Rodenticide Act. Your release for shipment of the product bearing the amended labeling constitutes acceptance of these conditions.

A stamped copy of the labeling is enclosed for your records.

If you have any questions concerning this letter, please call Martha DeLaney at (703) 557-6982.

Sincerely yours,


John H. Lee
Product Manager (31)
Antimicrobial Program Branch
Registration Division (TS-767C)

Enclosure

ACCEPTED
with COMMENTS
in EPA Letter Dated:

NOV 16 1988

Under the Federal Insecticide,
Fungicide, and Rodenticide Act,
as amended, for the pesticide
registered under EPA Reg. No.

346-38

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

STORAGE AND DISPOSAL

STORAGE. Keep container closed when not in use. Avoid another exposure to extreme temperatures.

DISPOSAL. Do not contaminate water, food, or feed by storage or disposal. Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture, or empty container is a violation of Federal Law. If these wastes cannot be disposed of by use according to label directions, contact your State Pesticide or Environment Agency, or the Hazardous Waste Response Team (nearest EPA Regional Office for guidance). Do not reuse container. Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or incineration, or, if allowed by state and local authority, by burning. If burned, stay out of smoke.

GENERAL FARM USES: Livestock and Poultry Houses, Swine Houses and Equipment

Disinfection Solution: 3 ounces Quat A Mone in 10 gallons of Water (470 ppm active quaternary)

- Remove all poultry and feeds from premises. Wash floors, walls, crates.
- Remove all litter and droppings from floors, walls, surfaces of facilities and fixtures traversed or occupied by poultry.
- Empty all troughs, racks, and other feeding and watering appliances. Wash with good detergent and rinse thoroughly.
- Saturate all surfaces with disinfectant solution using a hot soapy cloth or mop.
- Immerse all forks, shovels, and scrapers.
- Ventilate buildings, cars, boats, camps, and other confined places. Do not house poultry or employ equipment until treatment has been completed, set, or dried.
- As treated feed racks, feeders, troughs, automatic feeders, fountains, and waterers must be thoroughly rinsed with clean water before use.

Hatchery Equipment and Hatching Eggs. To sanitize equipment for hatching, wash to remove gross contamination and immerse briefly in a disinfection solution prepared by adding 1 ounce of Quat A Mone to 10 gallons of water. A contact time of at least 10 minutes is required for disinfection. To sanitize hatching equipment, remove all eggs and embryos. Wash equipment with detergent and rinse with water. Prepare disinfection solution as above. Quat A Mone in 10 gallons of water will sanitize equipment by covering in a brush or with.

DAIRY

CLEANING BULK MILK CANS. As a cleaning agent, Quat A Mone has been used to clean bulk tanks, hose tanks, and other equipment. Wash out with water. Prepare disinfection solution as above. Quat A Mone in 10 gallons of water will sanitize equipment by covering in a brush or with.

QUAT-A
DISINFECTANT
DEO

A Powerful Germicide for use
and for General Farm Use.

ACTIVE INGREDIENTS

n-Alkyl 16thyl (2,4,6-Cl₃)-4-
dimethyl benzyl ammonium
chloride
ammonium chloride

INERT INGREDIENTS

ANGER

KEEP OUT OF REACH

See Right Side Panel
and First

Very Flammable. For use only in
household.

Net Content

EPA Reg. No. 346-38

I. D. RUSSELL COMPANY LABO