



Du Pont

Oct 16, 1961

SUPPLEMENTAL LABELING

"KARMEX"<sup>®</sup>  
DIURON WEED KILLER 352-247  
FOR SELECTIVE CONTROL OF WEEDS IN APPLE AND PEAR ORCHARDS 1cf3

Du Pont "Karmex" Diuron Weed Killer is recommended for use in established apple and pear orchards to control certain seedling annual weeds such as crabgrass, watergrass, foxtail, pigweed, purslane, ragweed, chickweed, wild mustard, and lambs'-quarters.

At the rates recommended, "Karmex" will not control deep-rooted perennial or well-established annual weeds which should first be removed by cultivation or other means. Do not exceed recommended dosage rates; do not use on light, sandy or gravelly soils; and do not apply in areas into which the roots of other valuable plants or trees may extend, as injury may result. Use only under trees which have been established in the orchard for at least one year. Do not use under dwarf or semi-dwarf trees.

HOW TO APPLY: Apply to the area under the trees, or as a band application in the tree row; avoid contact of foliage or fruit. Use 4 lbs. of "Karmex" per acre (of area actually sprayed) in sufficient water (minimum 40 gallons per acre of area actually sprayed) to provide thorough and uniform coverage of the ground. Use a properly calibrated, fixed-boom power sprayer. Spray booms must be shut off while starting, turning, slowing, or stopping, or injury to trees may result. Continuous agitation as directed on container label is required to keep "Karmex" in suspension.

WHEN TO APPLY: Apply in the spring (March <sup>through</sup> to May) just prior to germination and growth of annual weeds. In areas such as the Far West, where there are separate winter and summer annual weed problems, it may be desirable to make a winter treatment (December to February) or to divide the treatment equally between a post-harvest fall application and a spring application (March to May). In any case, do not apply more than 4 lbs. of "Karmex" per <sup>year</sup> per acre (of area actually sprayed).

Best results are obtained if the herbicide is moved into the soil by moisture (rainfall or sprinkler irrigation) shortly after application.

NOTE: Do not replant treated areas to any crop within 2 years after application as injury to subsequent crops may result.

Change "To" to "Through"

OCT 16 1961

N4C T.W.

BEST DOCUMENT AVAILABLE



BETTER THINGS FOR BETTER LIVING... THROUGH CHEMISTRY

Oct 16, 1961  
352-247  
OCT 15 1961

4

IMPORTANT

BEFORE USING "KARMEX", READ AND CAREFULLY OBSERVE THE CAUTIONARY STATEMENTS AND ALL OTHER INFORMATION APPEARING ON THE PRODUCT LABEL.

This bulletin contains new or supplemental instructions for use of this product which may not appear on package label. Follow the instructions carefully.

3/21/61-A  
10/13/61

BEST DOCUMENT AVAILABLE

74