



AGRICULTURAL BULLETIN



LEXONE® 4L WEED KILLER
PLUS KARMEX® WEED KILLER
ASPARAGUS IN THE NORTHWEST

ACCEPTED

APR 26 1984

Under the Federal Insecticide,
Fungicide, and Rodenticide Act,
registered for the pesticide
category under
EPA Reg. No. 352-247

SUPPLEMENTAL LABELING

LEXONE® 4L EPA REG. NO. 352-382

KARMEX® EPA REG. NO. 352-247

LEXONE® 4L WEED KILLER AND COMBINATIONS WITH
KARMEX® FOR USE ON ESTABLISHED ASPARAGUS IN THE NORTHWEST

DIRECTIONS FOR USE

It is a violation of federal law to use this product
in a manner inconsistent with its labeling.

Du Pont Lexone® 4L Weed Killer is recommended alone or in combination with
Karmex® Weed Killer for use on established asparagus in the Northwest. For
weeds controlled, refer to package labels for "Lexone" 4L and "Karmex".

HOW TO USE

Use of the following:

"Lexone" 4L Preemergence Treatment Only: Make a single preemergence
application of 2 to 4 pts. "Lexone" 4L per acre in early spring prior to spear
emergence. Use the lower rate on coarse textured soils (low in clay or
organic matter). If field is to be disked, apply "Lexone" 4L after disking
but before crop emerges.

OR

"Lexone" 4L Preemergence and Post-Harvest Treatment: Make a single
preemergence application of 1 to 2 pts. "Lexone" 4L per acre in early spring
prior to spear emergence. Following harvest but before fern emerges, apply
1 to 3 pts. "Lexone" 4L per acre. Do not exceed 4 pts. "Lexone" 4L per acre
per growing season.

OR

"Lexone" 4L Preemergence and "Lexone" 4L Plus "Karmex" Post Harvest

Treatment: Make a single preemergence application of 1 to 2 pts.

"Lexone" 4L per acre in early spring prior to spear emergence. Following harvest but before fern emerges, apply a tank-mix combination of 1 to 1 1/2 pts. "Lexone" 4L plus 1 to 2 lbs. "Karmex" per acre.

Irrigated Asparagus: After the initial application of "Lexone" 4L, do not exceed more than 1/2" of water for the first irrigation. Additional irrigations may be normal programs.

Application: Apply in 3-10 gallons per acre by air or 10-40 gallons per acre by ground equipment.

NOTE: Do not use on newly seeded nor on young plants during the first growing season after setting crowns. Do not apply within 14 days of harvest. If field is to be disked, apply "Lexone" 4L after disking but before the crop emerges. Do not apply post-harvest application until after the last harvest of spears.

IMPORTANT

BEFORE USING "LEXONE" 4L IN COMBINATIONS, READ AND CAREFULLY OBSERVE THE CAUTIONARY STATEMENTS AND ALL OTHER INFORMATION APPEARING ON THE PRODUCT LABELS.

This bulletin contains new or supplemental instructions for use of these products which may not appear on these package labels. Follow the instructions carefully.

This labeling must be in the possession of the user at the time of pesticide application.



AGRICULTURAL BULLETIN



LEXONE® DF™ WEED KILLER
PLUS KARMEK® WEED KILLER
ASPARAGUS IN THE NORTHWEST

ACCEPTED

APR 26 1984

Under the Federal Insecticide,
Fungicide, and Rodenticide Act,
as amended, for the pesticide
registered under
EPA Reg. No. 352-247

SUPPLEMENTAL LABELING
LEXONE® DF™ EPA REG. NO. 352-390
KARMEK® EPA REG. NO. 352-247

LEXONE® DF™ WEED KILLER AND COMBINATIONS WITH
KARMEK® FOR USE ON ESTABLISHED ASPARAGUS IN THE NORTHWEST

DIRECTIONS FOR USE

It is a violation of federal law to use this product
in a manner inconsistent with its labeling.

Du Pont Lexone® DF™ Weed Killer is recommended alone or in combination with
Karmex® Weed Killer for use on established asparagus in the Northwest. For
weeds controlled, refer to package labels for "Lexone" DF™ and "Karmex".

HOW TO USE

Use of the following:

"Lexone" DF™ Preemergence Treatment Only: Make a single preemergence
application of 1 1/3 to 2 2/3 lbs. "Lexone" DF™ per acre in early spring
prior to spear emergence. Use the lower rate on coarse textured soils (low in
clay or organic matter). If field is to be disked, apply "Lexone" DF™
after disking but before crop emerges.

OR

"Lexone" DF™ Preemergence and Post-Harvest Treatment: Make a single
preemergence application of 2/3 to 1 1/3 lbs. "Lexone" DF™ per acre in
early spring prior to spear emergence. Following harvest but before fern
emerges, apply 2/3 to 2 lbs. "Lexone" DF™ per acre. Do not exceed 2 2/3
lbs. "Lexone" DF™ per acre per growing season.

OR

"Lexone" DFTM Preemergence and "Lexone" DFTM Plus "Karmex" Post Harvest Treatment: Make a single preemergence application of 2/3 to 1 1/3 lbs. "Lexone" DFTM per acre in early spring prior to spear emergence. Following harvest but before fern emerges, apply a tank-mix combination of 2/3 to 1 lb. "Lexone" DFTM plus 1 to 2 lbs. "Karmex" per acre.

Irrigated Asparagus: After the initial application of "Lexone" DFTM, do not exceed more than 1/2" of water for the first irrigation. Additional irrigations may be normal programs.

Application: Apply in 3-10 gallons per acre by air or 10-40 gallons per acre by ground equipment.

NOTE: Do not use on newly seeded nor on young plants during the first growing season after setting crowns. Do not apply within 14 days of harvest. If field is to be disked, apply "Lexone" DFTM after disking but before the crop emerges. Do not apply post-harvest application until after the last harvest of spears.

IMPORTANT

BEFORE USING "LEXONE" DFTM IN COMBINATIONS, READ AND CAREFULLY OBSERVE THE CAUTIONARY STATEMENTS AND ALL OTHER INFORMATION APPEARING ON THE PRODUCT LABELS.

This bulletin contains new or supplemental instructions for use of these products which may not appear on these package labels. Follow the instructions carefully.

This labeling must be in the possession of the user at the time of pesticide application.

030784