

JUL 30 1984

Under the Federal Insecticide,  
Fungicide, and Rodenticide Act,  
as amended, for the pesticide  
registered under  
EPA Reg. No. 352-247

SEE FOLDER IN BOTTOM FLAP FOR USE INSTRUCTIONS



# KARMEX<sup>®</sup>

## HERBICIDE

### WETTABLE POWDER

#### ACTIVE INGREDIENT:

Diuron [3-(3,4-dichlorophenyl)-1,1-dimethylurea] ..... 80%

INERT INGREDIENTS ..... 20%

EPA Reg. No. 352-247

#### KEEP OUT OF REACH OF CHILDREN. PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS

#### CAUTION! MAY IRRITATE EYES, NOSE, THROAT AND SKIN.

Avoid breathing dust or spray mist. Avoid contact with skin, eyes, and clothing.  
Wash thoroughly with soap and water after handling.

Do not apply this product in such a manner as to directly or through drift expose workers,  
or other persons. The area being treated must be vacated by unprotected persons.

#### STATEMENT OF PRACTICAL TREATMENT

If on skin: Wash with plenty of soap and water; get medical attention if irritation persists.  
If in eyes: Flush with plenty of water; get medical attention if irritation persists.

#### ENVIRONMENTAL HAZARDS

Do not apply directly to water. Do not contaminate water by cleaning of equipment or  
disposal of wastes.

**IMPORTANT:** Injury to or loss of desirable trees or other plants may result from failure  
to observe the following:

Do not apply (except as recommended for crop use), or drain or flush equipment on or near  
desirable trees or other plants, or on areas where their roots may extend, or in locations where  
the chemical may be washed or moved into contact with their roots. Do not use on home  
plantings of trees, shrubs or herbaceous plants, nor on lawns, walks, driveways, tennis courts,  
or similar areas. Prevent drift of dry powder or spray to desirable plants. Do not contaminate any  
body of water. Keep from contact with fertilizers, insecticides, fungicides, and seeds.

Thoroughly clean all traces of "Karmex" from application equipment immediately after use.  
Flush tank, pump, hoses, and boom with several changes of water after removing nozzle tips  
and screens (clean these parts separately).

3/20

(T) (H) (C)

EPA Est. 352-TX-1; 5905-AR-1; 477-TX-1  
SUPERSCRIPT USED IS FIRST LETTER OF LOT NUMBER

**NOTICE OF WARRANTY**

Du Pont warrants that this product conforms to the chemical description on the label thereof and is reasonably fit for the purposes stated on such label only when used in accordance with the directions under normal use conditions. It is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness, or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of Du Pont. In no case shall Du Pont be liable for consequential, special or indirect damages resulting from the use or handling of this product. All such risks shall be assumed by the buyer. DU PONT MAKES NO WARRANTIES OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS STATED ABOVE.

**SELECTIVE USE IN CROPS**

Do not use on any crop in Kern County, California

● **FIELD CROPS**

**Alfalfa:** United States  
**Artichoke:** California  
**Asparagus:** United States  
**Barley, Winter:** Western Oregon and Western Washington  
**Bermudagrass Pastures (Newly Sprigged):** United States  
**Birdsfoot Trefoil (Lotus):** Western Oregon  
**Corn (Field):**  
Postemergence—United States  
Preemergence—Arkansas, Louisiana, Mississippi and Tennessee  
**Cotton:**  
Preplant—Arizona and California

**Preemergence—**United States, except Arizona, California, and certain other areas  
**Postemergence—**United States  
**Grass Seed Crops (Perennial):** Colorado, Kansas, New Mexico, Oklahoma and Oregon  
**Oats:** Idaho, Oregon and Washington  
**Peppermint:** Pacific Northwest  
**Red Clover:** Western Oregon  
**Sorghum (Grain):** Southwestern States  
**Sugarcane:** Florida, Hawaii, Puerto Rico and Louisiana  
**Wheat, Winter:** Idaho, Oregon, Washington and certain other areas

● **FRUIT AND NUT CROPS**

**Apples:** United States  
**Bananas and Plantains:** United States  
**Blueberries, Caneberries and Gooseberries:** Indiana, Michigan and Ohio—Blueberries and Raspberries; Massachusetts and New Jersey—Blueberries; California—Raspberries, Blackberries, Boysenberries, Dewberries, and Loganberries; Western Oregon and Western Washington—Blueberries, Caneberries and Gooseberries

**Citrus:** Arizona (except Yuma area) and California (except Imperial and Coachella Valleys), Florida and Texas.  
**Grapes:** United States  
**Macadamia Nuts:** Hawaii  
**Olives:** California  
**Papayas:** United States  
**Peaches:** United States  
**Pears:** United States  
**Pecans:** United States  
**Pineapple:** Hawaii  
**Walnuts (English):** California

● **ORNAMENTAL CROPS**

**Ornamental Bulb Crops (Bulbous Iris, Narcissus):** Western Washington

**Plumous Fern:** Florida  
**Tree Plantings:** Colorado, Montana, Nebraska, North Dakota, South Dakota and Wyoming

**NON-CROP WEED CONTROL**

UTILITY, HIGHWAY, PIPELINE, RAILROAD RIGHT OF WAYS—INDUSTRIAL PLANT SITES—STORAGE AREAS—LUMBERYARDS—PETROLEUM TANK FARMS—FARM BUILDINGS—IRRIGATION AND DRAINAGE DITCHES

**STORAGE AND DISPOSAL**

**STORAGE:** Store product in original container only, away from other pesticides, fertilizer, food, or feed.  
**DISPOSAL:** Do not contaminate water, food, or feed by storage or disposal. Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.



## Product Labeling

### DIRECTIONS FOR USE

E. I. DU PONT DE NEMOURS & CO. (INC.)  
 AGRICULTURAL CHEMICALS DEPT., WILM., DE  
 AG 242 0684 Made in USA Printed in USA

**BEST AVAILABLE COPY**

#### ACTIVE INGREDIENT:

Diuron [3-(3,4-dichlorophenyl)-1,1-dimethylurea] ..... 80%

#### INERT INGREDIENTS .....

20%

EPA Reg. No. 352-247

#### KEEP OUT OF REACH OF CHILDREN PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS

#### CAUTION! MAY IRRITATE EYES, NOSE, THROAT, AND SKIN.

Avoid breathing dust or spray mist. Avoid contact with skin, eyes, and clothing. Wash thoroughly with soap and water after handling.

Do not apply this product in such a manner as to directly or through drift expose workers or other persons. The area being treated must be vacated by unprotected persons.

#### STATEMENT OF PRACTICAL TREATMENT

If on skin: Wash with plenty of soap and water; get medical attention if irritation persists.

If in eyes: Flush with plenty of water; get medical attention if irritation persists.

#### ENVIRONMENTAL HAZARDS

Do not apply directly to water. Do not contaminate water by cleaning of equipment or disposal of wastes.

**IMPORTANT**—Injury to or loss of desirable trees or other plants may result from failure to observe the following:

Do not apply (except as recommended for crop use), or drain or flush equipment on or near desirable trees or other plants, or on areas where their roots may extend, or in locations where the chemical may be washed or moved into contact with their roots. Do not use on home plantings of trees, shrubs or herbaceous plants, nor on lawns, walks, driveways, tennis courts, or similar areas. Prevent drift of dry powder or spray to desirable plants. Do not contaminate any body of water. Keep from contact with fertilizers, insecticides, fungicides, and seeds.

Thoroughly clean all traces of "Karmex" from application equipment immediately after use. Flush tank, pump, hoses, and boom with several changes of water after removing nozzle tips and screens (clean these parts separately).

#### GENERAL INFORMATION

DuPont "Karmex" Herbicide is a wettable powder to be mixed in water and applied as a spray for selective control of weeds in certain crops and for nonselective weed control on non-cropland areas. It is non-corrosive to equipment, nonflammable and nonvolatile.

"Karmex" may be applied to soil prior to emergence of weeds to control susceptible weed seedlings for an extended period of time; the degree of control and duration of effect will vary with the amount of chemical applied, soil texture, rainfall and other conditions. Soils high in clay or organic matter require higher dosages than soil low in clay or organic matter to obtain equivalent herbicide performance. Moisture is required to activate the chemical; best results are obtained if rainfall (or an other equivalent) occurs within 2 weeks

• **ALFALFA:** Treat only stands established for 1 year or more; do not apply to alfalfa under stress from (such as on shallow hard pans), alkali spots; nor to alfalfa on snow-covered or frozen ground.

Idaho, Oregon, Washington: Use 1½ to 3 lbs. per acre. Apply in fall after alfalfa becomes dormant but not later than January.

California (Dormant and Semi-Dormant Varieties): Use 2 to 4 lbs. per acre. Apply in fall or winter after alfalfa growth begins in the spring. Crop injury may result if applied too late. For best results, apply before weeds have emerged or before they are well established. If control of established weeds is improved by applying "Karmex" after they are well established, such use. Sufficient rainfall for soil activation of "Karmex" is required. Treated areas may be replanted to any crop after 2 weeks.

Arizona, Nevada: Use 1½ to 3 lbs. per acre; apply in January.

Eastern Colorado, Kansas: For control of tansymustard, apply in the fall or winter; use 2 lbs. per acre if weeds are present, apply 2 to 3 lbs. per acre in February.

Other Areas Where Alfalfa Becomes Winter Dormant: (Appalachian Mountains). Apply in March or early April.

• **ARTICHOKE**—California: Apply 2 to 4 lbs. per acre. Apply before weeds germinate or to emerging seedlings at the base of artichoke plants, keeping contact with the soil.

• **ASPARAGUS:** Apply as a band or broadcast treatment during the growing season (except as noted below), nor to newly established plants as severe injury may result. Preemergence weed control (greater than 5%).

**Established Plantings:** On light sandy soils and other soils high in clay or organic matter, use 2 to 4 lbs. per acre. Application should be made before weeds become established and no later than the early cutting period. For best cultural practices, application may be delayed until after the first cutting. Application may be made immediately following completion of the growing season. Applications are used in one season, do not exceed (irrigated crop), apply a single treatment of 4 lbs. per acre in early spring, incorporation of the chemical in the soil by rain to activate the herbicide.

**Newly Planted Crowns**—California (San Joaquin Delta): Use 2 to 4 lbs. per acre on soils high in clay or organic matter; use the lower rate on soils containing less than 2% organic matter. Do not treat crowns planted to a depth of 2 to 3 inches.

• **BARLEY, WINTER (Drill-Planted)**—Western Oregon: Apply 1½ to 2 lbs. per acre as soon as possible after planting. Do not treat areas to any crop within 1 year after last application.

• **BERMUDAGRASS PASTURES (Newly Sprigged)**—Apply 1½ to 2 lbs. per acre; add 1 pt. Surfactant WK per acre at time of treatment, temporary burn of exposed plants may result. Plant sprigs (stolons) 2" deep in a well-prepared seed bed; do not graze or mow for more than 2" deep as crop injury may result. Do not graze or mow for 30 days after application.

• **BIRDSFOOT TREFOIL (Lotus)**—Western Oregon: Do not apply to seedling trefoil as injury may result. May be applied to dormant (October 15 to December 15). Do not reseed or apply as injury to subsequent crops may result.

• **CORN (FIELD):** Apply 1½ to 2 lbs. per acre. Apply in fall or winter after alfalfa becomes dormant but not later than January.

• **POSTEMERGENCE**—Make a single application of ¼ lb. per acre. If nitrogen solution is not used, apply 1 lb. per acre. Apply as a directed spray when corn is at least 20" tall.

• **OVER TOP OF CORN.** Do not replant to any crop within 2 weeks. Corn may be planted the spring following treatment.

• **PREEMERGENCE**—Arkansas, Louisiana, Mississippi and Texas: Apply 1½ to 2 lbs. per acre as a broadcast or band treatment after planting.

of application.

"Karmex" may also be used to control emerged weeds. Results vary with rate applied and environmental conditions; best results are obtained on succulent weeds growing under conditions of high humidity and temperatures of 70°F or higher. Addition of a surfactant such as Du Pont Surfactant WK to the spray (where recommended) increases contact effects of "Karmex."

Since the effect of "Karmex" varies with soils, uniformity of application, and environmental conditions, it is suggested that growers limit their first use to small areas. Observe all cautions and limitations on labeling of all products used in mixtures.

**NOTICE OF WARRANTY**

Du Pont warrants that this product conforms to the chemical description on the label thereof and is reasonably fit for purposes stated on such label only when used in accordance with the directions under normal use conditions. It is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of Du Pont. In no case shall Du Pont be liable for consequential, special or indirect damages resulting from the use or handling of this product. All such risks shall be assumed by the Buyer. DU PONT MAKES NO WARRANTIES OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS STATED ABOVE.

**DIRECTIONS FOR USE**

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

DuPont "Karmex" Herbicide should be used only in accordance with recommendations on this label, or in separate published DuPont recommendations available through local dealers

Du Pont will not be responsible for losses or damages resulting from use of this product in any manner not specifically recommended by Du Pont. User assumes all risk associated with such non-recommended use.

Do not enter treated areas without protective clothing until sprays have dried. Because certain states may require more restrictive reentry intervals for various crops treated with this product, consult your State Department of Agriculture for further information

Written or oral warnings must be given to workers who are expected to be in a treated area or in an area about to be treated with this product. When oral warnings are given, warnings shall be given in a language customarily understood by workers. Oral warnings must be given if there is reason to believe that written warnings cannot be understood by workers. Warnings must include the following information. "CAUTION. Area treated with 'Karmex' on (date of application). Do not enter without appropriate protective clothing until sprays have dried. In case of contact, wash skin with plenty of soap and water; flush eyes with plenty of water. Get medical attention if irritation to skin or eyes persist."

Do not use in Kern County, California.

**SELECTIVE USE IN CROPS**

**PREEMERGENCE USE (Germinating Weeds):** "Karmex", at recommended rates, controls annual weeds such as:

Broadleaves			Grasses		
½ to 1 Lb./Acre	1½ to 2 Lbs./Acre	2 to 6 Lbs./Acre	½ to 1 Lb./Acre	1½ to 2 Lbs./Acre	2 to 6 Lbs./Acre
Lambsquarters	Annual groundcherry	Ageratum	Barnyardgrass	Annual bluegrass	Annual lovegrass
Pigweed	Annual morningglory	Annual smartweed	(watergrass)	Annual sweet	Annual ryegrass
Purslane	Chickweed	Annual sowthistle	Crabgrass	vernalgrass	Kyllinga
Ragweed	Corn spurry	Corn speedwell		Foxtail	Orchardgrass
	Dogfennel	Dayflower		Rabbit fescue	Peppergrass
	Fiddleneck	Flora's paintbrush		Red sprangletop	Ricegrass
	(amsackia)	Hawkbeard		Velvetgrass	Sandbur
	Gromwell	Horseweed			Seeding johnson-grass
	Knawel	Kochia			
	Pennycress	Marigold			
	Shepherdspurse	Mexican clover			
	Tansy-mustard	Pineappleweed			
	Wild buckwheat	Pokeweed			
	Wild lettuce	Rabbit tobacco			
	Wild mustard	Spanishneedles			
		Velvetleaf			
		(buttonweed)			
		Wild radish			

**BEST AVAILABLE COPY**

Partial control of the following weeds usually occurs at rates stated:

**Broadleaves**

**Grasses**

Deep. Do not repeat treated areas to crops other than corn or cotton within 18 months following broadcast treatment and 6 months following broadcast treatment as crop injury may result.

● **COTTON:** During a single crop season, do not exceed the following amounts of "Karmex" per acre injury to subsequent crops may result: 1 lb. on loamy sand; 1½ lbs. on sandy loam; 2 lbs. on clay loam; 2½ lbs. on clay. Injury may occur if "Karmex" is used in conjunction with soil-applied organic phosphorus pesticides. Do not allow livestock to graze treated cotton.

**Preplant—Arizona and California:** Use "Karmex" alone, or apply as a separate operation following preplant broadcast treatment with "Treflan" (incorporated according to directions on "Treflan" label). Apply "Karmex" as a broadcast spray after beds are formed, preirrigated, and final seedbeds prepared. Prior to planting, drag-off the tops of the beds and plant in moist soil not treated with "Karmex". Treated soil returned to the bed after planting when irrigation furrows are reformed after cotton has emerged. If more than two furrowing operations are made prior to lay-by, or deep furrows are made early, weed control may be reduced in furrow bottoms. Use at the following rates:

"Karmex" Alone: 1 to 2½ lbs. per acre.

"Karmex" Following "Treflan":

Soil Texture	Product Per Acre—Preplant	
	"Treflan"	"Karmex"
Sandy loam, loam, silt loam, silt	1 pt.	¾ to 1 lb.
Sandy clay loam, clay loam, silty clay loam, sandy clay, clay	1½ pts.	1 to 1½ lbs.

**Note:** Seeding disease may weaken plants and increase the possibility of injury from the use of "Treflan" followed by "Karmex". These treatments should be used only in conjunction with a standard fungicide seed treatment plus a good supplemental soil fungicide program such as captan-PCNB mixture.

\*Reg. trademark of Elanco Products Co

**Preemergence—U.S., except Arizona, California, and areas west of Interstate 35 or 35W in Texas and Oklahoma:** Use "Karmex" alone or apply as a separate operation following preplant treatment with "Treflan". Apply "Karmex" after planting but before cotton emerges. Do not treat cotton in deep furrows as crop injury may result; use only where cotton is planted on flat or raised seedbeds. Shallow incorporation (no deeper than ¼") with a rotary hoe or similar equipment following planting usually improves results, especially during dry weather. A wide press wheel should be used on the planter to provide a level seedbed for subsequent early season postemergence treatments. If moisture is insufficient to activate "Karmex" or if soil becomes crusted before crop emerges, a shallow rotary hoeing (no deeper than ¼") should be made before weeds become established.

"Karmex" Alone: Make a single application as a broadcast or band spray, using the following broadcast rates; for band treatment, use proportionately less.

Soil Texture**	Lbs. "Karmex" Per Acre
Loamy sand	¾
Sandy loam, loam, silt loam, silt	1
Sandy clay loam, clay loam, silty clay loam, sandy clay	1½
Silty clay, clay	2

\*\*Do not use on soils with less than 1% organic matter as crop injury may result

"Karmex" Following "Treflan" Preplant: Apply "Treflan" prior to planting as a broadcast or band treatment; incorporate according to directions on "Treflan" label. As a separate operation, apply "Karmex" band treatment (14" to 20" wide) after planting but before cotton emerges. Use at the following broadcast rates; for band treatment, use proportionately less. See "Note" under Preplant above

Soil Texture**	Product Per Acre	
	Preplant "Treflan"	Preemergence "Karmex"
Loamy sand	½ pt.	¾ lb.
Sandy loam, loam, silt loam, silt	1 pt.	1 lb.
Sandy clay loam, clay loam, silty clay loam, sandy clay, silty clay, clay	1½ pts.	1½ to 2 lbs.

\*\*Do not use on soils with less than 1% organic matter as crop injury may result.

**Postemergence—U.S.:** Apply only as a directed spray to cover weed foliage; adjust nozzles to minimize contact of cotton leaves with spray or drift or crop injury may result. DO NOT SPRAY OVER TOP OF COTTON

**Early Season—**Apply when cotton is at least 6" tall (at least 12" tall for Western irrigated cotton), when weeds are actively growing and do not exceed 2" in height. Apply as a band treatment at follow-up rates; for each 25 gals. of spray, add 1 pt. Surfactant WK. Two applications may be made if need

Tansy-mustard  
 Wild buckwheat  
 Wild lettuce  
 Wild radish  
 Rabbit tobacco  
 (barnyardgrass)  
 Wild radish

Partial control of the following weeds usually occurs at rates stated:

Broadleaves		Grasses	
1 Lb./Acre	4 Lbs./Acre	4 Lbs./Acre	8 to 10 Lbs./Acre
Annual morningglory	Horsenettle	Quackgrass	Guineagrass
Cocklebur			Maidencane
Prickly sida (teaweed)			Pangolagrass
Sesbania			
Sicklepod			

**POSTEMERGENCE USE (Emerging Seedling Weeds):** "Karmex" at recommended rates, controls annual weeds such as annual morningglory, barnyardgrass (watergrass), crabgrass, crowfoot, goosegrass, pigweed and purslane. Addition of Surfactant WK to the spray (where recommended) increases contact effects of "Karmex". Best results are obtained on succulent weeds growing under conditions of high humidity and temperatures of 70°F or higher.

**EQUIPMENT—SPRAY VOLUMES AND PRESSURES:** Use a tractor-mounted fixed-boom power sprayer properly calibrated to a constant speed and rate of delivery. Openings in screens should be equal to or larger than 50 mesh. Continuous agitation in the spray tank is required to keep the material in suspension. Agitate by mechanical or hydraulic means; if by-pass or return line is used, it should terminate at bottom of tank to minimize foaming. Avoid overlapping, and shut off spray booms while starting, turning, slowing or stopping, or injury to the crop may result.

For preemergence application, use 25 to 40 gals. per acre and spray pressure of 30 to 40 psi. For post-emergence application, use sufficient volume (min. 25 gals. per acre) for thorough coverage of weed foliage; use spray pressure of 20 to 25 psi to keep spray drift to a minimum.

**Aerial:** For alfalfa, asparagus, barley (winter), cotton (preplant or preemergence only), grass seed crops, pineapple, sugarcane and wheat (winter), application may be made by aircraft (5 to 10 gals. per acre); avoid overlapping of spray swath and avoid application under conditions where excessive drift may occur. Where land is bedded, make application parallel to rows.

**SPRAY PREPARATION:** Mix proper amount of "Karmex" into necessary volume of water; where use of Du Pont Surfactant WK is recommended, dilute with 10 parts of water and add as last ingredient to nearly full tank.

**USE RATES:** All dosages of "Karmex" are expressed as broadcast rates; for band treatment, use proportionately less. For example, use 1/2 of the broadcast rate when treating a 14" band where row spacing is 42". Where a range of dosages is given, use the lower rate on coarse textured soils (low in clay or organic matter) and the higher rate on fine textured soils (high in clay or organic matter); for postemergence application, use the lower rate on smaller weeds and the higher rate on larger weeds.

**SOIL LIMITATIONS:** Crop injury may result from failure to observe the following:

Unless otherwise directed, do not use on sand, loamy sand, gravelly soils or exposed subsoils; nor on pecans where organic matter is less than 1/2%; nor on alfalfa, apples, artichoke, barley (winter), bermudagrass pasture, citrus, cotton, grapes, oats, olives, papayas, peaches, pears, plumosus fern, sorghum, sugarcane, walnuts and wheat (winter) where organic matter is less than 1%; nor on blueberries, birdfoot trefoil, caneberrries, gooseberries, macadamia nuts and peppermint where organic matter is less than 2%.

Preemergence weed control will be reduced on high organic matter soils (greater than 5%, such as peat or muck).

**REPLANTING:** Unless otherwise directed, do not replant treated areas to any crop within 2 years after last application as injury to subsequent crops may result.

**FIELD CROPS**  
(See Soil Limitations)

A good seedbed must be prepared before preemergence use of "Karmex" as crop injury may result if application is made to ground which is cloddy or compacted resulting in improperly planted seed. Plant seed to depth specified. Unless otherwise directed, surface of the soil should not be cultivated or disturbed after application of "Karmex" and before emergence of the crop as weed control may be reduced and crop injury may result. However, if moisture is insufficient to activate the herbicide, a shallow cultivation (rotary hoe preferred) should be made after emergence of crops while weeds are small enough to be controlled by mechanical means.

silty clay, clay 1 1/2 pts. 2 pts.

\*\*Do not use on soils with less than 1% organic matter as crop injury may result.

**Postemergence—U.S.:** Apply only as a directed spray to cover weed foliage; adjust nozzles to prevent contact of cotton leaves with spray or drift or crop injury may result. DO NOT SPRAY OVER TOP OF COTTON.

**Early Season—**Apply when cotton is at least 6" tall (at least 12" tall for Western irrigated) when weeds are actively growing and do not exceed 2" in height. Apply as a band treatment; rates; for each 25 gals. of spray, add 1 pt. Surfactant WK. Two applications may be made.

Weed Problem (Up to 2' Tall)	Lbs. "Karmex" Per Acre (Broadcast Basis)
Annual grasses	1/2
Pigweed	1/4

For control of seedling perennial grasses such as johnsongrass and partial control of nutsedge, weed growth is under drought stress or as high as 4", add 2 to 3 1/2 lbs. disodium methylarsenate (63% anhydrous or equivalent) to above spray mixture. If DSMA is used, do not apply after 10:00 AM.

**Late Season (Lay-By)—**Apply 1 to 1 1/2 lbs. per acre (1 to 2 lbs. in Arizona and California) when cotton is at least 12" tall (at least 20" tall for Pima S-2). For control of germinating weed seedlings, apply beneath cotton plants and between rows immediately after last cultivation. In irrigated cotton, control is obtained if the field is irrigated within 3 to 4 days after application; thoroughly wet the ground over the row to carry the herbicide into the root zone of germinating weeds. After control of emerged annual weeds (up to 4" in height) at lay-by time, make a single application with Surfactant WK (1 pt. per 25 gals. spray), or use 1/2 to 3/4 lb. "Karmex" (plus surfactant) repeat later if needed.

**Replanting:** If initial seeding fails to produce a stand, cotton may be replanted in soil treated with "Karmex", alone or following "Trellan". Wherever possible, avoid disturbing soil. If necessary to rework soil before replanting, use shallow cultivation such as discing; do not move soil into the original drill area. Plant seed at least 1" deep. Do not retreat field with a second preemergence application during the same crop year as injury to the crop may result.

Subsequent Crops:

"Karmex"—Type of Application	Crops That May Follow Treated Cotton
Band preemergence or postemergence	Any crop 4 months after last application.
Band preemergence plus postemergence -or- Broadcast preemergence (and preplant) -or- Broadcast preemergence plus band postemergence	Cotton, soybeans, corn or grain sorghums or forage sorghums nor grass sorghums in spring. Do not replant treated areas to any crop within one year after last application as injury to subsequent crops may result.
Broadcast postemergence (lay-by)	Cotton, corn, grain sorghums (not sorghum) nor grass sorghums) the next spring. Do not replant treated areas to any other crop within one year after last application as injury to subsequent crops may result.

For subsequent crops in fields where "Trellan" is used, follow instructions on "Trellan".

Continued on

**ACCEPTED**  
  
 JUL 30 1984  
  
 Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 552-241

**BEST AVAILABLE COPY**

Continued from front side

**GRASS SEED CROPS (Perennial):** Except as noted, apply only to established plantings at least 1 year old.

**Colorado, Kansas, New Mexico and Oklahoma:** On sand bluestem, side oats grama and switchgrass, apply 2 to 3 lbs. per acre during the dormant period shortly before weed seedlings emerge. Do not apply after crop begins growth in the spring as crop injury may result. In fields where ash residues have accumulated from burning straw, use 3 lbs. per acre; spread unburned chaff or straw with a harrow or chopper before application.

**Western Oregon:** On alta fescue, Astoria bentgrass, Highland bentgrass, Kentucky bluegrass (Merion bluegrass) and orchardgrass, apply 2 to 4 lbs. per acre between October 1 and November 15. In fields where ash residues have accumulated from burning straw, use 3 to 4 lbs. per acre; spread unburned chaff or straw with a harrow or chopper before application. If perennial velvetgrass (*Holcus lanatus*) is a problem, use 4 lbs. per acre. For best results, apply as soon as possible after fall rains start. Established weeds (beyond 2 to 4 leaf stage) should be removed prior to treatment.

Well established vigorous stands of spring-planted alta fescue, Kentucky bluegrass and orchardgrass may be treated the following fall provided the crop is planted before April 1 and treatment is not applied before October 15; use 2 lbs. per acre.

**Oregon:** For use in newly planted bentgrass, Chewing fescue, Kentucky bluegrass, perennial ryegrass, orchardgrass and tall fescue. During planting operation, spray Aqua Nu-Char† or Gro-Safe†† or other suitable brands of activated charcoal as a 1" band on soil surface at rate of 300 lbs. per acre (broadcast basis; equivalent to 15 lbs. per acre of crop where row spacing is 20"). Mount nozzles to apply directly over seed rows to prevent crop injury. Follow with "Karmex" as a single broadcast spray at rate of 2½ to 3 lbs. per acre; apply as soon as possible after planting but before crops or weeds emerge and before rains or sprinkler irrigation. Fall or spring plantings may be treated; best results usually occur with early fall plantings. Treatment will not control downy brome or wild oats.

† Reg. trademark of Westvaco Corp.  
†† Reg. trademark of C.I. United States Inc.

**OATS (Drill-Planted):** Do not replant treated areas to any crop within one year after last application as injury to subsequent crops may result.

**Spring Oats—Idaho, Eastern Oregon, Eastern Washington:** Use in areas where average annual rainfall exceeds 16". Make a single application of 1 to 1½ lbs. per acre after planting, either before or after oats emerge but within 6 weeks of planting. Best results are usually obtained when application is made 3 to 4 weeks after planting. Apply before weeds are 3" to 4" tall.

**Winter Oats and Mixtures with Peas or Vetch—Western Oregon and Western Washington:** Make a single application of 1½ to 2 lbs. per acre as soon as possible after planting but before emergence of the crop.

**PEPPERMINT—Pacific Northwest:** Apply 3 lbs. per acre just after the last cultivation in the spring prior to emergence of peppermint. Do not apply to newly planted (less than 1 year) nor to emerged peppermint as injury may result.

**RED CLOVER—Western Oregon:** Make a single application of two pounds per acre on established red clover stands (at least 9 months). Apply "Karmex" when red clover is dormant (October 15 to December 15). Do not apply to seedling red clover, and do not replant treated area to any crop within one year after last application.

Treatment will control annual weeds such as bluegrass, chickweed, hawksbeard, rattail fescue, rye grass, and velvet grass.

**SORGHUM (GRAIN)—Southwestern States:** Apply ¼ to ½ lb. per acre; add 1 pt. Surfactant WK per 25 gals. of spray. Apply as a directed postemergence broadcast or band spray after sorghum is 15" tall to control weeds 2" to 4" in height. DO NOT SPRAY OVER TOP OF SORGHUM. Use the lower rate on broad-leaved weeds up to 2" tall; use the higher rate on grasses up to 2" and broadleaved weeds up to 4" tall. When the lower rate is used, a second application may be made if needed provided the amount applied in one crop year does not exceed ½ lb. per acre. Treatment of weeds under drought stress is usually ineffective.

Do not replant treated areas to crops other than cotton or corn within 4 months following band treatment and 6 months following broadcast treatment as crop injury may result.

**SUGARCANE:** To prevent possible crop injury on new cane varieties, tolerance to "Karmex" should be determined prior to adoption as field practice. Do not treat sugarcane growing on thinly covered subsoils or rocky areas as crop injury may result. Temporary chlorosis of the crop may result from application over emerged cane; to minimize chlorosis, use directed postemergence sprays.

**Florida: Preemergence—**For high organic soils, apply 2 to 4 lbs. per acre as a broadcast or band spray prior to weed emergence after planting or after harvesting plant crop (for ratoon crop). **Postemergence—**Make 1 or 2 applications of 2 lbs. per acre as needed by directed spray inter-row. Alternatively, for panicum control, make up to 3 applications of ½ to 1 lb. per acre as a directed spray after cane has emerged but before panicum exceeds 2" in height; add 1 qt. Surfactant WK per 100 gals. of spray. Adjust nozzles to spray beneath cane plants and between rows to cover weed foliage and to minimize contact of cane leaves with spray or drift. Do not apply more than 6 lbs. total per acre between planting (or ratooning) and harvest.

**Hawaii and Puerto Rico:** Apply 4 to 8 lbs. per acre as a broadcast spray prior to weed emergence after planting or after harvesting plant crop (for ratoon crop). A second and third application of 2 to 4 lbs. per acre may

be timed to activate the herbicide. Established perennial weeds require other special control procedure not apply under citrus trees that have been subjected to freezing within 6 months.

**Arizona (except Yuma area) and California (except Imperial and Coachella Valleys):** Apply 3 to 4 lb acre shortly after grove has been laid-up in final form (nontillage program) in late fall or early winter. Alternatively, apply 2 lbs. per acre in October or November and repeat at the same rate in March or April. Subsequent annual applications of 2 to 3 lbs. per acre will usually give adequate weed control.

**Florida:** Make a single application of 4 to 8 lbs. per acre, or apply 3 to 4 lbs. per acre followed by the rate 4 to 6 months later. On bearing citrus, apply any time when seasonal rains are expected; on nonbearing trees, apply when winter banks are pulled down.

For control of guineagrass, loosestrife, maidencane, paragrass, primrose willow and seamyrtle in adjacent citrus groves, use 1 lb. per 1000 sq. ft. (40 lbs. per acre) in sufficient water (min. 4 gals. per sq. ft.) to provide thorough and uniform coverage. Apply in the spring before weed growth starts or removal of vegetation. Repeat treatment on a spot basis to control hard-to-kill species such as guinea. In bedded groves, do not treat water furrows between the beds as injury to the trees may result.

**Texas:** Apply 2 to 4 lbs. per acre for annual weeds; use 4 to 6 lbs. per acre for control of johnson seedlings. Best results accompany application in the spring; well established weeds should be eliminated by cultivation prior to treatment.

**GRAPES:** Apply only to established vineyards (at least 3 years old) as a band treatment to grape on soils low in clay or organic matter (1 to 2%), severe plant injury may result if heavy rainfall or than one inch of irrigation occurs soon after treatment. This risk must be assumed by the user.

**East of the Rocky Mountains:** On soils low in clay or organic matter (1 to 2%), apply 2 to 3 lbs. per acre on soils high in clay or organic matter, apply 3 to 6 lbs. per acre. Apply in the spring just prior to germination and growth of annual weeds.

**West of the Rocky Mountains:** Apply in November, December or January. For initial treatment, apply 4 lbs. per acre; subsequent annual applications of 2 lbs. per acre will usually give adequate weed control. Do not apply to vines with trunks less than 1½" in diameter as injury may result.

**New York and Pennsylvania—Perennial Grasses:** Use only in established vineyards (at least 4 years old) for spot control of perennial grasses such as orchardgrass, quackgrass and ryegrass. Apply in the spring as a band treatment to ridged soil (2" to 4" high) under the trellis at the rate of 8 to 12 lbs. per acre. Band should not exceed 30". Do not apply more than once every 4 years. Use only on heavy soils such as silt loams, clay loams. Do not use in areas where grape roots are shallow or exposed because of high rock, poor drainage, or erosion as injury to grapes may result.

**MACADAMIA NUTS—Hawaii:** Use only under trees established in the orchard for at least 1 year. Apply 2 to 6 lbs. per acre immediately after harvest, preferably before weeds emerge; if weeds have emerged, add 1 pt. Surfactant WK per 25 gals. of spray. Retreat as needed but do not exceed 10 lbs. per acre per year.

**OLIVES—California:** Use only under trees established in the grove for at least 1 year. Apply 2 lbs. per acre after grove has been laid-up in final form in late October or November; repeat at same rate in March or April. Remove weed growth prior to treatment.

**PAPAYAS:** Use only under trees established in the orchard for at least 1 year. Apply 2½ to 5 lbs. per acre preferably before weeds emerge; if weeds have emerged, add 1 pt. Surfactant WK per 25 gals. of spray.

**PEACHES—**Use "Karmex" alone, or apply as a tank mixture with "Sinbar".

**"Karmex" Alone—**Use only under trees established in the orchard for at least 3 years. Apply 2 to 5 lb acre in the early spring before weeds emerge or during the early seedling stage of weed growth. Do not apply within 3 months of harvest; in the Far West, do not apply within 8 months of harvest.

**"Karmex" + "Sinbar"—**Use only under trees established in the orchard for at least 2 years. Apply in the spring or after harvest in the fall before weeds emerge or during early seedling stage of weed growth.

Soil Texture	Lbs. Product Per Acre	
	1 to 2% Organic Matter "Karmex" + "Sinbar"	More Than 2% Organic Matter "Karmex" + "Sinbar"
Sandy loam . . . . .	1 + 1	1½ + 1½
Loam, silt loam, silt . . . . .	1½ + 1½	2 + 2
Clay loam, clay . . . . .	2 + 2	2 + 2

Where crop is grown under furrow irrigation or under raised-basin flood irrigation (trees 4" to 6" waterline), apply only as a band treatment. Do not treat trees planted in the bottom of irrigation furrows or trees grown under flat flood or basin irrigation, as injury to trees may result. Where complete weed control to harvest is desired, additional weed control measures may be required during the growing season. Use only under trees established in the orchard for at least 2 years.

If weeds are present, add a surfactant (such as "Surfactant WK," "Osamul" 95 or "Starox" SK) to the spray at the rate of 1 to 2 qts. per 100 gals. and apply as a directed spray. DO NOT SPRAY OVER TOP OF CANE.

Do not apply more than 3 treatments nor more than 10 lbs. (Puerto Rico) or 12 lbs. (Hawaii) total per acre between planting (or ratooning) and harvest. Treated areas may be planted to sugarcane or pineapple one year after last application.

**Louisiana:** Use on plant cane seeded on fallowed ground. Make a single application of 3 to 3½ lbs. per acre at either of the following times. **Fall Treatment** (August through October)—Treat a 2 ft. band over the row after planting of cane, but before weeds or cane emerge. **Spring Treatment** (January through April)—if shading and off-barring are practiced, treat a 2 ft. band over the row before weeds or cane emerge.

**WHEAT, WINTER (Drill-Planted):** Crop injury may result where severe winter stress, disease or insect damage follows application; winter-sensitive varieties such as McDermid and Hyslop may be less tolerant of "Karmex" than winter-hardy varieties such as Gaines and Nugaines. Crop injury may also result from failure to observe the following: Do not use on sand or loamy sand soils, nor on gravelly or sandy loams low in organic matter (less than 1%), nor on thinly covered or exposed subsoil areas (clay knobs); do not treat wheat planted less than 1" deep; do not treat wheat where winter climatic conditions have caused "heaving" of plants; do not treat wheat plants lacking in vigor due to poor emergence, insect damage, disease, high alkalinity or other causes; do not apply after wheat has reached the "boot" stage of maturity; do not use with surfactants, or nitrogen solution. Do not replant treated areas to any other crop within 1 year after last treatment (except as noted) as injury to subsequent crops may result.

**Idaho, Oregon and Washington—East of Cascade Range:**

**Areas Where Average Annual Rainfall Exceeds 16 Inches:** Make a single application of 1 to 1½ lbs. per acre. **Fall Treatment:** For early fall-planted wheat (seeded before September 10), apply 3 to 6 weeks after planting but before weeds are 3" to 4" tall. Treatment after October 1 has generally given best results. Application should not be made after soil freezes in the fall. Wheat planted in late October should not be treated until the following spring. **Spring Treatment:** Apply as soon as wheat starts to grow in the spring. Treatment made prior to April 10 will usually give good results provided weed growth is less than 4" tall. Application later than May 1 may give poor results.

Alternatively, make a single application of ½ to 1 lb. "Karmex" plus ¼ lb. bromoxynil per acre as a tank mixture, either in the fall after wheat has emerged but before soil freezes or in the spring as soon as soil thaws; apply before weeds are 2" tall or across.

**Areas Where Average Annual Rainfall is 10 to 16 Inches:** After wheat is planted in the fall, make a single application of 1 to 1½ lbs. per acre when sufficient moisture is available to germinate wheat seed. Apply before soil freezes and before weeds are 2" tall. Application later than March 1 may give poor results.

**Note—**If fall-planted wheat fails to grow due to winter kill or adverse growing conditions after fall treatment, only fields treated before November 1 may be replanted to spring wheat. Spring wheat should not be planted before April 1, and only after deep disking and plowing to a depth of 4" to 6" prior to planting. Do not retreat field with a second application during the same crop year as injury to the crop may result.

**Oregon and Washington—West of Cascade Range:** Make a single application of 1½ to 2 lbs. per acre as soon as possible after planting; if wheat and weeds have emerged, apply before weeds are 3" to 4" tall. Alternatively, apply a tank mixture of "Karmex" plus bromoxynil as detailed above for "East of Cascade Range".

**Other Areas:** Make a single application in the spring as soon as wheat (fall-planted) starts to grow and before weeds are 2" tall. Application later than May 1 may give poor results.

**Central Plains and Midwest:** Use 1 to 2 lbs. per acre.

**Kansas, Oklahoma and Texas:** Do not use on sand or sandy loam soils. Use 1 lb. per acre on silt and silt loam soils and 1½-2 lbs. per acre on clay, clay loam, and silty clay loam soils.

**Northeast:** Use 1 to 1½ lbs. per acre.

### ● FRUIT AND NUT CROPS

(See Soil Limitations)

Unless otherwise directed, make a single application per year as a directed spray, avoiding contact of foliage and fruit with spray or drift. Do not graze livestock in treated orchards or groves.

**APPLES:** Use "Karmex" alone, or apply as a tank mixture with DuPont "Sinbar" Herbicide.

**"Karmex" Alone—**Use only under trees established in the orchard for at least 1 year; do not treat varieties grafted on full-dwarf root stocks. Apply 4 lbs. per acre in the spring (March through May). In the Far West, treatment may be made in winter (December through February), or apply 2 lbs. per acre as a postharvest treatment followed by 2 lbs. in the spring.

**"Karmex" + "Sinbar"—**Use only under trees established in the orchard for at least 2 years. Apply either in the spring or after harvest in the fall before weeds emerge or during early seedling stage of weed growth.

Soil Texture	Lbs. Product Per Acre	
	1 to 2% Organic Matter "Karmex" + "Sinbar"	More Than 2% Organic Matter "Karmex" + "Sinbar"
Sandy loam	2	1½ + 1½
Loam, silt loam, silt	3	1¾ + 1¾
Clay loam, clay	4	2 + 2

**FULL-DWARF ROOT STOCKS.** Apply 4 lbs. per acre in the spring (March through May), or 2 lbs. per acre, which may be made in winter (December through February), or apply 2 lbs. per acre as a postharvest treatment followed by 2 lbs. in the spring.

- **PECANS:** Use "Karmex" alone or apply as a tank mixture with "Sinbar." Make a single band or broadcast application as a directed spray using a minimum of 30 gals. of water per acre. Apply in the spring before weeds emerge or during the early seedling stage of growth.

Soil Texture	Lbs. Product Per Acre		
	"Karmex" Alone*	-OR-	Tank Mixture "Karmex" + "Sinbar" **
Sandy loam	2		1½ + 1½
Loam, silt loam, silt	3		1¾ + 1¾
Clay loam, clay	4		2 + 2

\*Use only on trees established in the grove for at least 3 yrs. and on soils with at least ½% organic matter

\*\*Use on trees established in the grove for at least 1 yr. and on soils with at least 1% organic matter

**Note:** Do not use on eroded areas where subsoil or roots are exposed, nor on trees that are diseased or lacking in vigor or on trees planted in irrigation furrows as injury to the trees may result

- **PINEAPPLE—Hawaii:** Apply 4 to 8 lbs. per acre as a broadcast spray just before or immediately after planting but prior to weed emergence. Use 4 lbs. per acre after harvesting plant crop (for ratoon crop). For plant crop only, a second and third broadcast or interspace application may be made prior to differentiation at the rate of 2 lbs. per acre at intervals of not less than 2 months. Additional applications to plant crop may be made as needed to interspace only using 2 lbs. per acre. Do not apply more than 3 broadcast sprays (maximum 12 lbs. per acre) prior to differentiation nor more than 16 lbs. total per acre per plant crop. Treated areas may be planted to pineapple or sugarcane 1 year after last application.
- **WALNUTS (English)—California:** Use only under trees established in the orchard for at least 1 year. At initial treatment, apply 3 to 5 lbs. per acre after the orchard has been laid up in final form (nontillage system) in late fall or early winter; retreat annually with 2 to 3 lbs. per acre. Alternatively, apply 2 lbs. per acre in October or November and repeat at same rate in March or April.

### ● ORNAMENTAL CROPS

(See Soil Limitations)

- **ORNAMENTAL BULB CROPS (Bulbous Iris, Narcissus)—Western Washington:** Make a single application of 4 lbs. per acre. Apply after planting but no later than 4 weeks prior to bulb emergence (usually late September or October). Do not replant treated areas to any crop within 1 year after last application as injury to subsequent crops may result.
- **PLUMOSUS FERN—Florida:** Hand weed and mow fern; then make a single application of 3 lbs. per acre within 3 to 5 days. Do not cultivate or disturb soil after application as crop injury may result. Treat established stands at least 1 year old.
- **TREE PLANTINGS—Colorado, Montana, Nebraska, North Dakota, South Dakota, Wyoming:** Use only under established plantings (1 year or older) of American elm, caragana, cottonwood, Douglas fir, green ash, honeysuckle, Ponderosa pine, redcedar, Russian olive and Siberian elm. Use 2½ to 5 lbs. per acre, apply as a band 4 ft. wide in the tree row (2 ft. on each side of row). For example, 1 oz. "Karmex" (4 level tablespoonfuls) treats 135 ft. of tree row (2 ft. on each side of row) at the rate of 5 lbs. per acre. Apply as a directed spray in early spring before weeds emerge and before trees leaf out. Do not apply to foliage of trees, nor under trees growing in low areas as injury to the trees may result.

**BEST AVAILABLE COPY**

### ● NON-CROP WEED CONTROL

"Karmex" is an effective herbicide for the control of many annual and perennial grasses and herbaceous weeds on non-cropland areas where bare ground is desired. The degree of control and duration of effect vary with the amount of chemical applied, soil texture, rainfall and other conditions.

"Karmex" may be used as a preemergence treatment at any time of the year except when ground is frozen, provided adequate moisture is supplied by rainfall or artificial means to activate the herbicide. Best results are obtained if application is made to the soil shortly before weed growth begins. If dense growth is present, remove tops and spray the ground.

Increased contact activity on established weeds may be obtained by the addition of Du Pont Surfactant at the rate of 2 qts. per 100 gals. of spray mixture. Apply as a drenching spray to actively growing weeds.



Soil Texture	1 to 2% Organic Matter "Karmex" + "Sinbar"	More Than 2% Organic Matter "Karmex" + "Sinbar"
Sandy loam	1 + 1	1½ + 1½
Loam, silt loam, silt	1½ + 1½	2 + 2
Clay loam, clay	2 + 2	2 + 2

Where crop is grown under furrow irrigation or under raised-berm flood irrigation (trees 4" to 6" above waterline), apply only as a band treatment. Do not treat trees planted in the bottom of irrigation furrows, nor trees grown under flat flood or basin irrigation, as injury to trees may result. Where complete weed control to harvest is desired, additional weed control measures may be required during the growing season.

- **BANANAS AND PLANTAINS—New Plantings:** To control annual weeds, apply 1½ to 3 lbs. per acre after planting but before weeds emerge. Do not apply to loose soil directly over the planting material.

**Established Plantings:** For control of annuals and for top-kill of perennials such as bermudagrass, birdseed grass and guineagrass, apply 3 to 6 lbs. per acre plus 1 pt. Surfactant WK (or suitable equivalent) per 25 gals. of spray; avoid contact of plants with spray or drift as injury may result. When tall, dense weed growth is present, remove weed growth before application. If application is made to soil free of weeds, omit the surfactant from the spray. Repeat treatment as needed, but do not apply more often than 6-week intervals nor more than a total of 2 lbs. per acre (broadcast basis) in a 12-month period.

**Note:** Do not replant treated areas to any crop within 2 years after last application as injury to subsequent crops may result, except that sugarcane or pineapple may be planted one year after last application.

- **BLUEBERRIES, CANEBERRIES AND GOOSEBERRIES:** Use only in fields which have been established for at least 1 year. Do not apply to berries interplanted with fruit trees; do not apply to plants whose roots are exposed as injury may result. Apply as a band treatment at base of canes or bushes; for spring application, apply before germination and growth of annual weeds.

**Indiana, Michigan and Ohio—Blueberries:** Apply 2 to 4 lbs. per acre in late spring; alternatively, apply 2 lbs. per acre in the fall and repeat at same rate in the spring. **Raspberries:** Apply 3 lbs. per acre in the spring. **Massachusetts—Blueberries:** Apply 2 lbs. per acre in late spring.

**New Jersey—Blueberries:** For control of winter annuals, apply 2 lbs. per acre in October, November or December, or a single application of 2½ lbs. per acre may be applied in early to mid spring.

**California—Raspberries, Blackberries, Boysenberries, Dewberries and Loganberries:** For control of winter annuals, apply 2 lbs. per acre in October or November; repeat at same rate in late spring to control summer annuals. A single application of 3 lbs. per acre in January or February will control both winter and summer annuals in some areas, but the separate fall and spring schedule is preferred.

**Western Oregon and Western Washington—Blueberries, Caneberries and Gooseberries:** Use same schedule as recommended for California.

- **CITRUS:** Use only under trees established in the grove for at least 1 year. Time application as indicated for specific areas, except application may be made any time of the year where sprinkler or flood irrigation can

remove tops and spray the ground.

Increased contact activity on established weeds may be obtained by the addition of Du Pont Surfact at the rate of 2 qts. per 100 gals. of spray mixture. Apply as a drenching spray to actively growing during warm weather when daily temperature will exceed 70°F.

Except for small areas, use a fixed-boom power sprayer properly calibrated to insure a constant application. Mix proper amount of "Karmex" into volume of water necessary to obtain uniform coverage. Surfactant WK is used, dilute with 10 parts of water and add as last ingredient to nearly full tank. must be kept in suspension at all times. Agitate by mechanical or hydraulic means in the spray tank or return line is used, it should terminate at bottom of tank to minimize foaming. Openings in should be equal to or larger than 50 mesh.

**General Weed Control:** To control most weeds for an extended period of time on non-cropland such as utility, highway, pipeline and railroad right of ways, petroleum tank farms, lumberyards, storage area, trial plant sites, and around farm buildings—apply 5 to 20 lbs. per acre to control most annual weeds; 20 to 60 lbs. per acre for perennial weeds; additional treatment may be required where a longer period of control is desired or when hard-to-kill, deep-rooted perennial weeds such as johnsongrass are present. In low rainfall areas, "Karmex" may not provide satisfactory control of deep-rooted perennial weeds. For weed control on small areas, use one-half cupful of "Karmex" per 100 sq. ft. for a dosage of approximately 50 lbs. per acre.

**Irrigation and Drainage Ditches:** Apply 5 to 20 lbs. per acre to control most annual weeds; use 20 lbs. per acre to control both annual and perennial weeds. Apply only when water is not in the ditch. For irrigation ditches, apply during the non-crop season, and when ditch is not in use. To minimize movement of "Karmex" with irrigation water (to avoid possible crop injury), it is essential that the herbicide be in the soil by moisture. Apply before expected seasonal rainfall (if possible when soil in the ditch is moist). Following treatment, if rainfall has not totaled at least 4 inches, fill ditch with water and stand for 72 hours; drain off and waste remaining water before using ditch. Do not treat any ditch in which roots of trees or other desirable plants may extend as injury may result.

## STORAGE AND DISPOSAL

**STORAGE:** Store product in original container only, away from other pesticides, fertilizer, food or feed.

**DISPOSAL:** Do not contaminate water, food, or feed by storage or disposal. Wastes resulting from use of this product may be disposed of on site or at an approved waste disposal facility.

Completely empty bag into application equipment. Then dispose of empty bag in a sanitary land fill by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

BEST AVAILABLE COPY



DU PONT  
WETTABLE POWDER

**KARMEX**<sup>®</sup>

WETTABLE POWDER

## Product Labeling

### DIRECTIONS FOR USE

E. I. DU PONT DE NEMOURS & CO. (INC.)  
BIOCHEMICALS DEPT., WILMINGTON, DEL.  
B 20160 283    Made in USA    Printed in USA

BEST AVAILABLE COPY

### ACTIVE INGREDIENT:

Diuron [3-(3,4-dichlorophenyl)-1,1-dimethylurea] ..... 80%

### INERT INGREDIENTS .....

20%

EPA Reg. No. 352-247

### KEEP OUT OF REACH OF CHILDREN. PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS

#### CAUTION! MAY IRRITATE EYES, NOSE, THROAT, AND SKIN.

Avoid breathing dust or spray mist. Avoid contact with skin, eyes, and clothing. Wash thoroughly with soap and water after handling.

#### STATEMENT OF PRACTICAL TREATMENT

If on skin: Wash with plenty of soap and water; get medical attention if irritation persists.  
If in eyes: Flush with plenty of water; get medical attention if irritation persists.

#### ENVIRONMENTAL HAZARDS

Do not apply directly to water. Do not contaminate water by cleaning of equipment or disposal of wastes.

**IMPORTANT**—Injury to or loss of desirable trees or other plants may result from failure to observe the following:

Do not apply (except as recommended for crop use), or drain or flush equipment on or near desirable trees or other plants, or on areas where their roots may extend, or in locations where the chemical may be washed or moved into contact with their roots. Do not use on home plantings of trees, shrubs or herbaceous plants, nor on lawns, walks, driveways, tennis courts, or similar areas. Prevent drift of dry powder or spray to desirable plants. Do not contaminate any body of water. Keep from contact with fertilizers, insecticides, fungicides, and seeds.

Thoroughly clean all traces of "Karmex" from application equipment immediately after use. Flush tank, pump, hoses and boom with several changes of water after removing nozzle tips and screens (clean these parts separately).

### GENERAL INFORMATION

Du Pont "Karmex" Weed Killer is a wettable powder to be mixed in water and applied as a spray for selective control of weeds in certain crops and for nonselective weed control on non-cropland areas. It is non-corrosive to equipment, nonflammable and nonvolatile.

"Karmex" may be applied to soil prior to emergence of weeds to control susceptible weed seedlings for an extended period of time; the degree of control and duration of effect will vary with the amount of chemical applied, soil texture, rainfall and other conditions. Soils high in clay or organic matter require higher dosages than soil low in clay or organic matter to obtain equivalent herbicide performance. Moisture is required to activate the chemical; best results occur if rainfall (or sprinkler irrigation) occurs within 2 weeks of application.

"Karmex" may also be used to control emerged weeds. Results vary with rate applied and environmental conditions; best results are obtained on succulent weeds growing under conditions of high humidity and

grass mixture. Do not apply to alfalfa under stress from disease, insect damage, shallow root penetration (such as on shallow hard pans), alkaline spots; nor to flooded fields as crop injury may result. Do not use on snow-covered or frozen ground.

**Idaho, Oregon, Washington:** Use 1½ to 2 lbs. per acre for control of volunteer alfalfa. Use 4 lbs. per acre. Apply in fall after alfalfa becomes dormant but no later than mid December.

**California (Dormant and Semi-Dormant Varieties):** Use 1½ to 3 lbs. per acre for control of volunteer alfalfa. Use 4 lbs. per acre. Apply in fall or winter after alfalfa becomes dormant or semi-dormant, but before growth begins in the spring. Crop injury may result if application is made to actively growing alfalfa. Best results, apply before weeds have emerged or become established (2" in height or diameter). Control of established weeds is improved by applying "Karmex" with a suitable contact herbicide registered for such use. Sufficient rainfall for soil activation of "Karmex" is unlikely in California after February. Treated areas may be replanted to any crop after one year from last application if rate does not exceed 2 lbs. per acre.

**Arizona, Nevada:** Use 1½ to 3 lbs. per acre. Apply in fall after alfalfa becomes dormant but no later than January.

**Eastern Colorado, Kansas:** For control of pansy mustard, apply 1 lb. per acre shortly after emergence of alfalfa in the fall or winter; use 2 lbs. per acre if weeds are 2" to 4" in height. Alternatively, if other weeds are present, apply 2 to 3 lbs. per acre in February or March.

**Other Areas Where Alfalfa Becomes Winter Dormant:** Use 1½ to 3 lbs. per acre (1½ to 2 lbs. per acre in the Appalachian Mountains). Apply in March or early April, but before spring growth begins.

- **ARTICHOKE**—California: Apply 2 to 4 lbs. per acre in late fall or early winter after the last cultivation. Apply before weeds germinate or to emerging seedlings. Direct spray to cover the area between the rows at the base of artichoke plants, keeping contact with crop plants at a minimum.
- **ASPARAGUS:** Apply as a band or broadcast treatment. Do not apply to young plants during the growing season (except as noted below), nor to newly seeded asparagus, nor on plants with exposed roots. Severe injury may result. Preemergence weed control will be reduced on high organic matter (greater than 5%).

**Established Plantings:** On light sandy soils and other soils low in clay or organic matter, apply 1 to 2 lbs. per acre. On soils high in clay or organic matter, use 2 to 4 lbs. per acre. Two applications may be used; the first application should be made before weeds become established but no earlier than 4 weeks before emergence and no later than the early cutting period (if weeds are controlled until the cutting period, cultural practices, application may be delayed until immediately after the last cultivation), a second application may be made immediately following completion of harvest provided rainfall is expected. When two applications are used in one season, do not exceed 3 lbs. per acre per application. In Washington (irrigated crop), apply a single treatment of 4 lbs. per acre. If treatment is delayed until late fall or early spring, incorporation of the chemical in the top 1" to 2" of soil may substitute for irrigation of rain to activate the herbicide.

**Newly Planted Crowns**—California (San Joaquin Delta): Make a single application of 2 to 4 lbs. per acre on soils high in clay or organic matter, use the lower rate on clay loams and the higher rate on peaty soils. Do not use on soils containing less than 2% organic matter. Soil must be settled by rainfall or irrigation prior to treatment. Do not treat crowns planted to a depth of less than 2".

- **BARLEY, WINTER (Drill-Planted)**—Western Oregon and Western Washington: Make a single application of 1½ to 2 lbs. per acre as soon as possible after planting but before emergence of barley. Do not apply to treated areas to any crop within 1 year after last application as injury to subsequent crops may result.
- **BERMUDAGRASS PASTURES (Newly Sprigged):** Apply 1 to 3 lbs. after planting and before emergence of bermudagrass or weeds. Alternatively, for control of emerged annual weeds up to 4" in height, apply ½ to 1 lb. per acre; add 1 pt. Surfactant WK per 25 gals. of spray. If bermudagrass has emerged, temporary burn of exposed plant parts may occur. Plant sprigs (stolons) 2" deep in a well-prepared seedbed; do not treat areas where sprigs are planted less than 2" deep as crop injury may result. Do not graze or feed foliage from treated areas to livestock within 7 days after application.

- **BIRDSFOOT TREFOIL (Lotus)**—Western Oregon: Treat only stands established for at least 1 year. Do not apply to seedling trefoil as injury may result. Make a single application of 2 lbs. per acre when dormant (October 15 to December 15). Do not replant treated areas to any crop within 1 year after application as injury to subsequent crops may result.

- **CORN (FIELD):**
  - **Postemergence**—Make a single application of ¾ lb. per acre in combination with non-pressure nitrogen solution. If nitrogen solution is not used, apply 1 lb. per acre; add 1 pt. Surfactant WK per 25 gals. of spray. Apply as a directed spray when corn is at least 20" high and weeds are no taller than 3". DO NOT OVER TOP OF CORN. Do not replant to any crop within 1 year, except that cotton, corn and grain sorghum may be planted the spring following treatment.
  - **Preemergence**—Arkansas, Louisiana, Mississippi and Tennessee: Make a single application of ½ to 1 lb. per acre as a broadcast or band treatment after planting but before corn emerges. Plant corn at least 2" deep. Do not replant treated areas to crops other than corn or cotton within 4 months following broadcast treatment and 6 months following broadcast treatment as crop injury may result.
- **COTTON:** During a single crop season, do not exceed the following amounts of "Karmex" per

of 70°F or higher. Addition of a surfactant such as Du Pont Surfactant WK to the spray (where recommended) increases contact effects of "Karmex".

Since the effect of "Karmex" varies with soils, uniformity of application, and environmental conditions, it is suggested that growers limit their first use to small areas. Observe all cautions and limitations on labeling of all products used in mixtures.

### NOTICE OF WARRANTY

Du Pont warrants that this product conforms to the chemical description on the label thereof and is reasonably fit for purposes stated on such label only when used in accordance with the directions under normal use conditions. It is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of Du Pont. In no case shall Du Pont be liable for consequential, special or indirect damages resulting from the use or handling of this product. All such risks shall be assumed by the Buyer. DU PONT MAKES NO WARRANTIES OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS STATED ABOVE.

### DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

Du Pont "Karmex" Weed Killer should be used only in accordance with recommendations on this label, or in separate published Du Pont recommendations available through local dealers.

Du Pont will not be responsible for losses or damages resulting from use of this product in any manner not specifically recommended by Du Pont. User assumes all risk associated with such non-recommended use.

Do not use in Kern County, California.

### SELECTIVE USE IN CROPS

**PREEMERGENCE USE (Germinating Weeds):** "Karmex", at recommended rates, controls annual weeds such as:

Broadleaves			Grasses		
% to 1 Lb./Acre	1½ to 2 Lbs./Acre	2 to 6 Lbs./Acre	% to 1 Lb./Acre	1½ to 2 Lbs./Acre	2 to 6 Lbs./Acre
Lambquarters	Annual groundcherry	Ageratum	Barnyardgrass	Annual bluegrass	Annual ryegrass
Pigweed	Annual morningglory	Annual smartweed	(watergrass)	Annual sweet	Annual lovegrass
Purslane	Chickweed	Annual sowthistle	Crabgrass	vernalgrass	Kyllinga
Ragweed	Corn spurry	Corn speedwell		foxtail	Orchardgrass
	Dogfennel	Dayflower		Rattail fescue	Peppergrass
	Fiddleneck	Flora's paintbrush		Red sprangletop	Ricegrass
	(amsinckia)	Hawksbeard		Velvetgrass	Sandbur
	Gromwell	Horseweed			Seedling johnson
	Knawel	Kochia			grass
	Pennycress	Marrigold			
	Shepherdspurse	Mexican clover			
	Tansy-mustard	Pineappleweed			
	Wild buckwheat	Pokeweed			
	Wild lettuce	Rabbit tobacco			
	Wild mustard	Spanishneedles			
		Velvetleaf			
		(bultonweed)			
		Wild radish			

Partial control of the following weeds usually occurs at rates stated:

Broadleaves		Grasses	
1 Lb./Acre	4 Lbs./Acre	4 Lbs./Acre	8 to 10 Lbs./Acre
Annual morningglory	Horsecelle	Quackgrass	Guineagrass
Cocklebur			Maidencane
Prickly sida (leaved)			Pangolagrass
Sesbania			
Sicklepod			

**BEST AVAILABLE COPY**

**POSTEMERGENCE USE (Emerging Seedling Weeds):** "Karmex" at recommended rates, controls annual weeds such as annual morningglory, barnyardgrass (watergrass), crabgrass, crowfoot, goosegrass, pigweed and purslane. Addition of Surfactant WK to the spray (where recommended) increases contact effects of "Karmex". Best results are obtained on succulent weeds growing under conditions of high humidity and temperatures of 70°F or higher.

**EQUIPMENT—SPRAY VOLUMES AND PRESSURES:** Use a tractor-mounted fixed boom power sprayer properly calibrated to a constant speed and rate of delivery. Openings in screens should be equal to or larger than 50 mesh. Continuous agitation in the spray tank is required to keep the material in suspension. Apply by mechanical or hydraulic means if the use of cotton lines is desired.

injury to subsequent crops may result: 1 lb. on loamy sand; 1½ lbs. on silty loam; 2 lbs. on clay; 2½ lbs. on clay. Injury may occur if "Karmex" is used in conjunction with soil-applied organic phosphorus pesticides. Do not allow livestock to graze treated cotton.

**Preplant—Arizona and California:** Use "Karmex" alone, or apply as a separate operation following preplant broadcast treatment with "Treflan" (incorporated according to directions on "Treflan" label). "Karmex" as a broadcast spray after beds are formed, preirrigated, and final seedbeds prepared. Planting, drag off the tops of the beds and plant in moist soil not treated with "Karmex". Treated beds returned to the bed after planting when irrigation furrows are reformed after cotton has emerged. If more than two furrowing out operations are made prior to lay by, or deep furrows are made early, weed control may be reduced in furrow bottoms. Use at the following rate:

"Karmex" Alone: 1 to 2½ lbs. per acre  
"Karmex" Following "Treflan":

Soil Texture	Product Per Acre—Preplant	
	"Treflan"	"Karmex"
Sandy loam, loam, silt loam, silt	1 pt.	1 to 1½ lbs.
Sandy clay loam, clay loam, silty clay loam, sandy clay, clay	1½ pts.	1 to 1½ lbs.

**Note:** Seeding disease may weaken plants and increase the possibility of injury from the use of "Karmex". These treatments should be used only in conjunction with a standard fungicide treatment plus a good supplemental soil fungicide program such as Du Pont "Derosan" 65W Fungicide or captan-PCNB mixture.

\*R.g. trademark of Elanco Products Co.

**Preemergence—U.S., except Arizona, California, and areas west of Interstate 35 or 35W in Texas:** Use "Karmex" alone or apply as a separate operation following preplant broadcast treatment with "Treflan". Apply "Karmex" after planting but before cotton emerges. Do not treat cotton in deep furrows as crop injury may result; use only where cotton is planted on flat or raised seedbeds. Shallow incorporation (no deeper than ¼") with a rotary hoe or similar equipment following planting usually improves results, especially during dry weather. A wide press wheel should be used on the planter to provide a level seedbed for subsequent early season postemergence treatments. If moisture is insufficient to activate "Karmex" or if soil becomes crusted before crop emerges, a shallow rotary hoeing (no deeper than ¼") should be made before weeds become established.

"Karmex" Alone: Make a single application as a broadcast or band spray, using the following broadcast rates. For band treatment, use proportionately less.

Soil Texture**	Lbs. "Karmex" Per Acre
Loamy sand	2
Sandy loam, loam, silt loam, silt	1
Sandy clay loam, clay loam, silty clay loam, sandy clay	1½
Silty clay, clay	2

\*\*Do not use on soils with less than 1% organic matter as crop injury may result.

"Karmex" Following "Treflan" Preplant: Apply "Treflan" prior to planting as a broadcast or band treatment; incorporate according to directions on "Treflan" label. As a separate operation, apply "Karmex" as a band treatment (14" to 20" wide) after planting but before cotton emerges. Use at the following broadcast rates. For band treatment, use proportionately less. See "Note" under Preplant above.

Soil Texture**	Product Per Acre	
	Preplant "Treflan"	Preemergence "Karmex"
Loamy sand	1 pt.	1 lb.
Sandy loam, loam, silt loam, silt	1 pt.	1 lb.
Sandy clay loam, clay loam, silty clay loam, sandy clay, silty clay, clay	1½ pts.	1½ to 2 lbs.

\*\*Do not use on soils with less than 1% organic matter as crop injury may result.

**Postemergence—U.S.:** Apply only as a directed spray to cover weed foliage; adjust nozzles to minimize contact of cotton leaves with spray or drift or crop injury may result. DO NOT SPRAY OVER TOP OF COTTON.

**Early Season—Apply** when cotton is at least 6" tall (at least 12" tall for Western irrigated cotton) when weeds are actively growing and do not exceed 2" in height. Apply as a band treatment at the following rates. For each 25 gals. of spray, add 1 pt. Surfactant WK. Two applications may be made if necessary.

Weed Problem (Up to 2" Tall)	Lbs. "Karmex" Per Acre (Broadcast Basis)
Annual Weeds	1 to 2 lbs.

Velvetleaf  
(buttonweed)  
Wild radish

Fungicide, and Rodenticide Act,  
as amended, for the pesticide  
registered under  
EPA Reg. No. 354-247

Partial control of the following weeds usually occurs at rates stated:

1 Lb./Acre	4 Lbs./Acre	4 Lbs./Acre	8 to 10 Lbs./Acre
Annual morningglory	Horsenettle	Quackgrass	Guineagrass
Cocklebur			Maidencane
Prickly sida (teaweed)			Pangolagrass
Sesbania			
Sicklepod			

**POSTEMERGENCE USE (Emerging Seedling Weeds):** "Karmex" at recommended rates, controls annual weeds such as annual morningglory, barnyardgrass (watergrass), crabgrass, crowfoot, goosegrass, bigweed and purslane. Addition of Surfactant WK to the spray (where recommended) increases contact effects of "Karmex". Best results are obtained on succulent weeds growing under conditions of high humidity and temperatures of 70°F or higher.

**EQUIPMENT—SPRAY VOLUMES AND PRESSURES:** Use a tractor-mounted fixed boom power sprayer properly calibrated to a constant speed and rate of delivery. Openings in screens should be equal to or larger than 50 mesh. Continuous agitation in the spray tank is required to keep the material in suspension. Agitate by mechanical or hydraulic means; if by-pass or return line is used, it should terminate at bottom of tank to minimize foaming. Avoid overlapping, and shut off spray booms while starting, turning, slowing or stopping, or injury to the crop may result.

For preemergence application, use 25 to 40 gals. per acre and spray pressure of 30 to 40 psi. For post emergence application, use sufficient volume (min. 25 gals. per acre) for thorough coverage of weed foliage; use spray pressure of 20 to 25 psi to keep spray drift to a minimum.

**Aerial:** For alfalfa, asparagus, barley (winter), cotton (preplant or preemergence only), grass seed crops, pineapple, sugarcane and wheat (winter), application may be made by aircraft (5 to 10 gals. per acre); avoid overlapping of spray swath and avoid application under conditions where excessive drift may occur. Where land is bedded, make application parallel to rows.

**SPRAY PREPARATION:** Mix proper amount of "Karmex" into necessary volume of water, where use of Du Pont Surfactant WK is recommended, dilute with 10 parts of water and add as last ingredient to nearly full tank.

**USE RATES:** All dosages of "Karmex" are expressed as broadcast rates; for band treatment, use proportionately less. For example, use 1/2 of the broadcast rate when treating a 14" band where row spacing is 42". Where a range of dosages is given, use the lower rate on coarse textured soils (low in clay or organic matter) and the higher rate on fine textured soils (high in clay or organic matter); for postemergence application, use the lower rate on smaller weeds and the higher rate on larger weeds.

**SOIL LIMITATIONS:** Crop injury may result from failure to observe the following.

Unless otherwise directed, do not use on sand, loamy sand, gravelly soils or exposed subsoils nor on pecans where organic matter is less than 1/2%; nor on alfalfa, apples, artichoke, barley (winter), bermudagrass pasture, citrus, cotton, grapes, oats, olives, papayas, peaches, pears, plumosus fern, sorghum, sugarcane, walnuts and wheat (winter) where organic matter is less than 1%; nor on blueberries, birdsfoot trefoil, caneberries, gooseberries, macadamia nuts and peppermint where organic matter is less than 2%.

Preemergence weed control will be reduced on high organic matter soils (greater than 5%, such as peat or muck).

**REPLANTING:** Unless otherwise directed, do not replant treated areas to any crop within 2 years after last application as injury to subsequent crops may result.

● **FIELD CROPS**  
(See Soil Limitations)

A good seedbed must be prepared before preemergence use of "Karmex" as crop injury may result if application is made to ground which is cloddy or compacted resulting in improperly planted seed. Plant seed to depth specified. Unless otherwise directed, surface of the soil should not be cultivated or disturbed after application of "Karmex" and before emergence of the crop as weed control may be reduced and crop injury may result. However, if moisture is insufficient to activate the herbicide, a shallow cultivation (rotary hoe preferred) should be made after emergence of crops while weeds are small enough to be controlled by mechanical means.

BEST AVAILABLE COPY

rates for band treatment, use proportionately less. See "Note" under Preplant above.

Soil Texture**	Product Per Acre	
	Preplant "Treflan"	Preemergence "Karmex"
Loamy sand	1/2 pt.	1/3 lb.
Sandy loam, loam, silt loam, silt	1 pt.	1 lb.
Sandy clay loam, clay loam, silty clay loam, sandy clay, silty clay, clay	1 1/2 pts.	1 1/4 to 2 lbs.

\*\*Do not use on soils with less than 1% organic matter as crop injury may result.

**Postemergence—U.S.:** Apply only as a directed spray to cover weed foliage; adjust nozzles to minimize contact of cotton leaves with spray or drift or crop injury may result. DO NOT SPRAY OVER TOP OF COTTON.

**Early Season—**Apply when cotton is at least 6" tall (at least 12" tall for Western irrigated cotton) when weeds are actively growing and do not exceed 2" in height. Apply as a band treatment at following rates for each 25 gals. of spray, add 1 pt. Surfactant WK. Two applications may be made if needed.

Weed Problem (Up to 2" Tall)	Lbs. "Karmex" Per Acre (Broadcast Basis)
Annual grasses	1 1/2
Pigweed	1 1/4

For control of seedling perennial grasses such as johnsongrass and partial control of nutsedge or weed growth is under drought stress or as high as 4", add 2 to 3 1/2 lbs. disodium methylarsenate (16% anhydrous or equivalent) to above spray mixture. If DSMA is used, do not apply after first cotton harvest.

**Late Season (Lay-By)—**Apply 1 to 1 1/2 lbs. per acre (1 to 2 lbs. in Arizona and California) when cotton is at least 12" tall (at least 20" tall for Pima S-2). For control of germinating weed seedlings, apply beneath cotton plants and between rows immediately after last cultivation. In irrigated cotton, best control is obtained if the field is irrigated within 3 to 4 days after application; thoroughly wet the surface of the ground over the row to carry the herbicide into the root zone of germinating weeds. Alternative control of emerged annual weeds (up to 4" in height) at lay-by time, make a single application in combination with Surfactant WK (1 pt. per 25 gals. spray), or use 1/2 to 3/4 lb. "Karmex" (plus surfactant) per acre, repeat later if needed.

**Replanting:** If initial seeding fails to produce a stand, cotton may be replanted in soil treated preemergence with "Karmex", alone or following "Treflan". Wherever possible, avoid disturbing the seedbed. If necessary to rework soil before replanting, use shallow cultivation such as disking; do not remove soil into the original drill area. Plant seed at least 1" deep. Do not retreat field with a second preplant or preemergence application during the same crop year as injury to the crop may result.

**Subsequent Crops:**

"Karmex"—Type of Application	Crops That May Follow Treated Cotton
Band preemergence or postemergence	Any crop 4 months after last application.
Band preemergence plus postemergence -or- Broadcast preemergence (and preplant) -or- Broadcast preemergence plus band postemergence	Cotton, soybeans, corn or grain sorghums (not sorghums or forage sorghums nor grass sorghums) the next spring. Do not replant treated areas to any other crop within one year after last application as injury to subsequent crops may result.
Broadcast postemergence (lay-by)	Cotton, corn, grain sorghums (not sorghums or forage sorghums nor grass sorghums) the next spring. Do not replant treated areas to any other crop within one year after last application as injury to subsequent crops may result.

For subsequent crops in fields where "Treflan" is used, follow instructions on "Treflan" label.

Continued on reverse

Continued from front side

**GRASS SEED CROPS (Perennial):** Except as noted, apply only to established plantings at least 1 year old.

**Colorado, Kansas, New Mexico and Oklahoma:** On sand bluestem, side oats grama and switchgrass, apply 2 to 3 lbs. per acre during the dormant period shortly before weed seedlings emerge. Do not apply after crop begins growth in the spring as crop injury may result. In fields where ash residues have accumulated from burning straw, use 3 lbs. per acre; spread unburned chaff or straw with a harrow or chopper before application.

**Western Oregon:** On alta fescue, Astoria bentgrass, Highland bentgrass, Kentucky bluegrass (Merion bluegrass) and orchardgrass, apply 2 to 4 lbs. per acre between October 1 and November 15. In fields where ash residues have accumulated from burning straw, use 3 to 4 lbs. per acre; spread unburned chaff or straw with a harrow or chopper before application. If perennial velvetgrass (*Holcus lanatus*) is a problem, use 4 lbs. per acre. For best results, apply as soon as possible after fall rains start. Established weeds (beyond 2 to 4 leaf stage) should be removed prior to treatment.

Well established vigorous stands of spring-planted alta fescue, Kentucky bluegrass and orchardgrass may be treated the following fall provided the crop is planted before April 1 and treatment is not applied before October 15; use 2 lbs. per acre.

**Oregon:** For use in newly planted bentgrass, Chewing fescue, Kentucky bluegrass, perennial ryegrass, orchardgrass and tall fescue. During planting operation, spray Aqua Nu-Char† or Gro-Safe†† or other suitable brands of activated charcoal as a 1" band on soil surface at rate of 300 lbs. per acre (broadcast basis; equivalent to 15 lbs. per acre of crop where row spacing is 20"). Mount nozzles to apply directly over seed rows to prevent crop injury. Follow with "Karmex" as a single broadcast spray at rate of 2 1/2 to 3 lbs. per acre; apply as soon as possible after planting but before crops or weeds emerge and before rains or sprinkler irrigation. Fall or spring plantings may be treated; best results usually occur with early fall plantings. Treatment will not control downy brome or wild oats.

† Reg. trademark of Westvaco Corp  
 †† Reg. trademark of ICI United States Inc

**OATS (Drill-Planted):** Do not replant treated areas to any crop within one year after last application as injury to subsequent crops may result.

**Spring Oats—Idaho, Eastern Oregon, Eastern Washington:** Use in areas where average annual rainfall exceeds 16". Make a single application of 1 to 1 1/2 lbs. per acre after planting, either before or after oats emerge but within 6 weeks of planting. Best results are usually obtained when application is made 3 to 4 weeks after planting. Apply before weeds are 3" to 4" tall.

**Winter Oats and Mixtures with Peas or Vetch—Western Oregon and Western Washington:** Make a single application of 1 1/2 to 2 lbs. per acre as soon as possible after planting but before emergence of the crop.

**PEPPERMINT—Pacific Northwest:** Apply 3 lbs. per acre just after the last cultivation in the spring prior to emergence of peppermint. Do not apply to newly planted (less than 1 year) nor to emerged peppermint as injury may result.

**RED CLOVER—Western Oregon:** Make a single application of two pounds per acre on established red clover stands (at least 9 months). Apply Karmex when red clover is dormant (October 15 to December 15). Do not apply to seedling red clover, and do not replant treated area to any crop within one year after last application.

Treatment will control annual weeds such as bluegrass, chickweed, hawksbeard, rattail fescue, rye grass, and velvet grass.

**SORGHUM (GRAIN)—Southwestern States:** Apply 1 1/4 to 1/2 lb. per acre; add 1 pt. Surfactant WK per 25 gals. of spray. Apply as a directed postemergence broadcast or band spray after sorghum is 15" tall to control weeds 2" to 4" in height. DO NOT SPRAY OVER TOP OF SORGHUM. Use the lower rate on broad leaved weeds up to 2" tall; use the higher rate on grasses up to 2" and broadleaved weeds up to 4" tall. When the lower rate is used, a second application may be made if needed provided the amount applied in one crop year does not exceed 1/2 lb. per acre. Treatment of weeds under drought stress is usually ineffective.

Do not replant treated areas to crops other than cotton or corn within 4 months following band treatment and 6 months following broadcast treatment as crop injury may result.

**SUGARCANE:** To prevent possible crop injury on new cane varieties, tolerance to "Karmex" should be determined prior to adoption as field practice. Do not treat sugarcane growing on thinly covered subsoils or rocky areas as crop injury may result. Temporary chlorosis of the crop may result from application over emerged cane; to minimize chlorosis, use directed postemergence sprays.

**Florida: Preemergence—**For high organic soils, apply 2 to 4 lbs. per acre as a broadcast or band spray prior to weed emergence after planting or after harvesting plant crop (for ratoon crop). **Postemergence—**Make 1 or 2 applications of 2 lbs. per acre as needed by directed spray inter-row. Alternatively, for panicum control, make up to 3 applications of 1/2 to 1 lb. per acre as a directed spray after cane has emerged but before panicum exceeds 2" in height; add 1 qt. Surfactant WK per 100 gals. of spray. Adjust nozzles to spray beneath cane plants and between rows to cover weed foliage and to minimize contact of cane leaves with spray or drift. Do not apply more than 6 lbs. total per acre between planting (or ratooning) and harvest.

**Hawaii and Puerto Rico:** Apply 4 to 8 lbs. per acre as a broadcast spray prior to weed emergence after planting or after harvesting plant crop (for ratoon crop). A second and third application of 2 to 4 lbs. per acre may

**Arizona (except Yuma) and California (except Imperial and Coachella Valleys):** Apply 3 to 4 lbs. per acre shortly after grove has been laid-up in final form (nontillage program) in late fall or early winter. Alternatively, apply 2 lbs. per acre in October or November and repeat at the same rate in March or April. Subsequent annual applications of 2 to 3 lbs. per acre will usually give adequate weed control.

**Florida and Puerto Rico:** Make a single application of 4 to 8 lbs. per acre, or apply 3 to 4 lbs. per acre followed by the same rate 4 to 6 months later. On bearing citrus, apply any time when seasonal rains are expected; on nonbearing trees, apply when winter banks are pulled down.

For control of guineagrass, loosestrife, maidencane, paragrass, primrose willow and seamyrtle in ditch adjacent to citrus groves, use 1 lb. per 1000 sq. ft. (40 lbs. per acre) in sufficient water (min. 4 gals. per 10 sq. ft.) to provide thorough and uniform coverage. Apply in the spring before weed growth starts or after removal of vegetation. Repeat treatment on a spot basis to control hard-to-kill species such as guineagrass in bedded groves, do not treat water furrows between the beds as injury to the trees may result.

**Texas:** Apply 2 to 4 lbs. per acre for annual weeds; use 4 to 6 lbs. per acre for control of johnsongrass seedlings. Best results accompany application in the spring; well established weeds should be eliminated by cultivation prior to treatment.

**GRAPES:** Apply only to established vineyards (at least 3 years old) as a band treatment to grape rows. On soils low in clay or organic matter (1 to 2%), severe plant injury may result if heavy rainfall or more than one inch of irrigation occurs soon after treatment. This risk must be assumed by the user.

**East of the Rocky Mountains:** On soils low in clay or organic matter (1 to 2%), apply 2 to 3 lbs. per acre; on soils high in clay or organic matter, apply 3 to 6 lbs. per acre. Apply in the spring just prior to germination and growth of annual weeds.

**West of the Rocky Mountains:** Apply in November, December or January. For initial treatment, apply 3 to 4 lbs. per acre; subsequent annual applications of 2 lbs. per acre will usually give adequate weed control. Do not apply to vines with trunks less than 1 1/2" in diameter as injury may result.

**New York and Pennsylvania—Perennial Grasses:** Use only in established vineyards (at least 4 years old) for spot control of perennial grasses such as orchardgrass, quackgrass and ryegrass. Apply in the spring as a band treatment to ridged soil (2" to 4" high) under the trellis at the rate of 8 to 12 lbs. per acre. Band width should not exceed 30". Do not apply more than once every 4 years. Use only on heavy soils such as loam or silt loams, clay loams. Do not use in areas where grape roots are shallow or exposed because of high bank rock, poor drainage, or erosion as injury to grapes may result.

**MACADAMIA NUTS—Hawaii:** Use only under trees established in the orchard for at least 1 year. Apply 2 to 6 lbs. per acre immediately after harvest, preferably before weeds emerge; if weeds have emerged, add 1 pt. Surfactant WK per 25 gals. of spray. Retreat as needed but do not exceed 10 lbs. per acre per year.

**OLIVES—California:** Use only under trees established in the grove for at least 1 year. Apply 2 lbs. per acre after grove has been laid-up in final form in late October or November, repeat at same rate in March or April. Remove weed growth prior to treatment.

**PAPAYAS:** Use only under trees established in the orchard for at least 1 year. Apply 2 1/2 to 5 lbs. per acre, preferably before weeds emerge; if weeds have emerged, add 1 pt. Surfactant WK per 25 gals. of spray.

**PEACHES—**Use "Karmex" alone, or apply as a tank mixture with "Sinbar".

**"Karmex" Alone—**Use only under trees established in the orchard for at least 3 years. Apply 2 to 5 lbs. per acre in the early spring before weeds emerge or during the early seedling stage of weed growth. Do not apply within 3 months of harvest; in the Far West, do not apply within 8 months of harvest.

**"Karmex" + "Sinbar"—**Use only under trees established in the orchard for at least 2 years. Apply either in the spring or after harvest in the fall before weeds emerge or during early seedling stage of weed growth.

Soil Texture	Lbs. Product Per Acre	
	1 to 2% Organic Matter "Karmex" + "Sinbar"	More Than 2% Organic Matter "Karmex" + "Sinbar"
Sandy loam	1 + 1	1 1/2 + 1 1/2
Loam, silt loam, silt	1 1/2 + 1 1/2	2 + 2
Clay loam, clay	2 + 2	2 + 2

Where crop is grown under furrow irrigation or under raised berm flood irrigation (trees 4" to 6" at waterline), apply only as a band treatment. Do not treat trees planted in the bottom of irrigation furrow nor trees grown under flat flood or basin irrigation, as injury to trees may result. Where complete weed control to harvest is desired, additional weed control measures may be required during the growing season.

If weeds are emerged, add a surfactant (such as Surfactant WK, "Osamul" 95 or "Sterox" SK) to the spray at the rate of 1 to 2 qts. per 100 gals. and apply as a directed spray. DO NOT SPRAY OVER TOP OF CANE.

Do not apply more than 3 treatments nor more than 10 lbs. (Puerto Rico) or 12 lbs. (Hawaii) total per acre between planting (or ratooning) and harvest. Treated areas may be planted to sugarcane or pineapple one year after last application.

**Louisiana:** Use on plant cane seeded on fallowed ground. Make a single application of 3 to 3½ lbs. per acre at either of the following times. **Fall Treatment** (August through October)—Treat a 2 ft. band over the row after planting of cane, but before weeds or cane emerge. **Spring Treatment** (January through April)—if shaving and off-barring are practiced, treat a 2 ft. band over the row before weeds or cane emerge.

**WHEAT, WINTER (Drill-Planted):** Crop injury may result where severe winter stress, disease or insect damage follows application; winter-sensitive varieties such as McDermid and Hyslop may be less tolerant of "Karmex" than winter-hardy varieties such as Gaines and Nugaines. Crop injury may also result from failure to observe the following: Do not use on sand or loamy sand soils, nor on gravelly or sandy loams low in organic matter (less than 1%), nor on thinly covered or exposed subsoil areas (clay knobs); do not treat wheat planted less than 1½" deep; do not treat wheat where winter climatic conditions have caused "heaving" of plants; do not treat wheat plants lacking in vigor due to poor emergence, insect damage, disease, high alkalinity or other causes; do not apply after wheat has reached the "boot" stage of maturity; do not use with surfactants, or nitrogen solution. Do not replant treated areas to any other crop within 1 year after last treatment (except as noted) as injury to subsequent crops may result.

**Idaho, Oregon and Washington—East of Cascade Range:**

**Areas Where Average Annual Rainfall Exceeds 16 Inches:** Make a single application of 1 to 1½ lbs. per acre. **Fall Treatment:** For early fall-planted wheat (seeded before September 10), apply 3 to 6 weeks after planting but before weeds are 3" to 4" tall. Treatment after October 1 has generally given best results. Application should not be made after soil freezes in the fall. Wheat planted in late October should not be treated until the following spring. **Spring Treatment:** Apply as soon as wheat starts to grow in the spring. Treatment made prior to April 10 will usually give good results provided weed growth is less than 4" tall. Application later than May 1 may give poor results.

Alternatively, make a single application of ½ to 1 lb. "Karmex" plus ¼ lb. bromoxynil per acre as a tank mixture in the fall after wheat has emerged but before soil freezes or in the spring as soon as soil thaws; apply before weeds are 2" tall or across.

**Areas Where Average Annual Rainfall is 10 to 16 Inches:** After wheat is planted in the fall, make a single application of 1 to 1½ lbs. per acre when sufficient moisture is available to germinate wheat seed. Apply before soil freezes and before weeds are 2" tall. Application later than March 1 may give poor results.

**Note—**If fall-planted wheat fails to grow due to winter kill or adverse growing conditions after fall treatment, only fields treated before November 1 may be replanted to spring wheat. Spring wheat should not be planted before April 1, and only after deep discing and plowing to a depth of 4" to 6" prior to planting. Do not retreat field with a second application during the same crop year as injury to the crop may result.

**Oregon and Washington—West of Cascade Range:** Make a single application of 1½ to 2 lbs. per acre as soon as possible after planting, if wheat and weeds have emerged, apply before weeds are 3" to 4" tall. Alternatively, apply a tank mixture of "Karmex" plus bromoxynil as detailed above for "East of Cascade Range".

**Other Areas:** Make a single application in the spring as soon as wheat (fall-planted) starts to grow and before weeds are 2" tall. Application later than May 1 may give poor results.

**Central Plains and Midwest:** Use 1 to 2 lbs. per acre.

**Kansas, Oklahoma and Texas:** Use 1½ to 2 lbs. per acre.

**Northeast:** Use 1 to 1½ lbs. per acre.

## ● FRUIT AND NUT CROPS

(See Soil Limitations)

Unless otherwise directed, make a single application per year as a directed spray, avoiding contact of foliage and fruit with spray or drift. Do not graze livestock in treated orchards or groves.

● **APPLES:** Use "Karmex" alone, or apply as a tank mixture with Du Pont "Sinbar" Weed Killer.

**"Karmex" Alone—**Use only under trees established in the orchard for at least 1 year, do not treat varieties grafted on full-dwarf root stocks. Apply 4 lbs. per acre in the spring (March through May). In the Far West, treatment may be made in winter (December through February), or apply 2 lbs. per acre as a postharvest treatment followed by 2 lbs. in the spring.

**"Karmex" + "Sinbar"—**Use only under trees established in the orchard for at least 2 years. Apply either in the spring or after harvest in the fall before weeds emerge or during early seedling stage of weed growth.

Soil Texture	Lbs. Product Per Acre					
	1 to 2% Organic Matter "Karmex" + "Sinbar"		More Than 2% Organic Matter "Karmex" + "Sinbar"			
Sandy loam	1	+	1	1½	+	1½
Loam, silt loam, silt	1	+	1	1½	+	1½
Clay loam, clay	1	+	1	2	+	2

full-dwarf root stocks. Apply 4 lbs. per acre in the spring (March through May). In the Far West, treatment may be made in winter (December through February), or apply 2 lbs. per acre as a postharvest treatment followed by 2 lbs. in the spring.

● **PECANS:** Use "Karmex" alone or apply as a tank mixture with "Sinbar." Make a single band or broadcast application as a directed spray using a minimum of 30 gals. of water per acre. Apply in the spring before weeds emerge or during the early seedling stage of growth.

Soil Texture	Lbs. Product Per Acre				
	"Karmex" Alone*	-OR-	Tank Mixture "Karmex" + "Sinbar" **		
Sandy loam	2		1½	+	1½
Loam, silt loam, silt	3		1½	+	1½
Clay loam, clay	4		2	+	2

\* Use only on trees established in the grove for at least 3 yrs. and on soils with at least 1% organic matter.  
\*\* Use on trees established in the grove for at least 1 yr. and on soils with at least 1% organic matter.

**Note:** Do not use on eroded areas where subsoil or roots are exposed, nor on trees that are diseased or lacking in vigor or on trees planted in irrigation furrows as injury to the trees may result.

● **PINEAPPLE—Hawaii and Florida:** Apply 4 to 8 lbs. per acre as a broadcast spray just before or immediately after planting but prior to weed emergence. Use 4 lbs. per acre after harvesting plant crop (for ratooned crop). For plant crop only, a second and third broadcast or interspace application may be made prior to differentiation at the rate of 2 lbs. per acre at intervals of not less than 2 months. Additional application to plant crop may be made as needed to interspace only using 2 lbs. per acre. Do not apply more than 3 broadcast sprays (maximum 12 lbs. per acre) prior to differentiation nor more than 16 lbs. total per plant crop. Treated areas may be planted to pineapple or sugarcane 1 year after last application.

● **WALNUTS (English)—California:** Use only under trees established in the orchard for at least 1 year. A preharvest treatment, apply 3 to 5 lbs. per acre after the orchard has been laid up in final form (nontillage program) in late fall or early winter; retreat annually with 2 to 3 lbs. per acre. Alternatively, apply 2 lbs. per acre in October or November and repeat at same rate in March or April.

## ● ORNAMENTAL CROPS

(See Soil Limitations)

● **ORNAMENTAL BULB CROPS (Bulbous Iris, Narcissus)—Western Washington:** Make a single application of 4 lbs. per acre. Apply after planting but no later than 4 weeks prior to bulb emergence (usually late September or October). Do not replant treated areas to any crop within 1 year after last application as injury to subsequent crops may result.

● **PLUMOSUS FERN—Florida:** Hand weed and mow fern, then make a single application of 3 lbs. per acre within 3 to 5 days. Do not cultivate or disturb soil after application as crop injury may result. Treat established stands at least 1 year old.

● **TREE PLANTINGS—Colorado, Montana, Nebraska, North Dakota, South Dakota, Wyoming:** Use only under established plantings (1 year or older) of American elm, caragana, cottonwood, Douglas fir, green ash, honeysuckle, Ponderosa pine, redcedar, Russian olive and Siberian elm. Use 2½ to 5 lbs. per acre, apply as a band 4 ft. wide in the tree row (2 ft. on each side of row). For example, 1 oz. "Karmex" (4 level tablespoons) treats 135 ft. of tree row (2 ft. on each side of row) at the rate of 5 lbs. per acre. Apply as a directed spray in early spring before weeds emerge and before trees leaf out. Do not apply to foliage of trees, nor under trees growing in low areas as injury to the trees may result.

## ● NON-CROP WEED CONTROL

"Karmex" is an effective herbicide for the control of many annual and perennial grasses and herbaceous weeds on non-cropland areas where bare ground is desired. The degree of control and duration of effect vary with the amount of chemical applied, soil texture, rainfall and other conditions.

"Karmex" may be used as a preemergence treatment at any time of the year except when ground is frozen. provided adequate moisture is supplied by rainfall or artificial means to activate the herbicide. Best results are obtained if application is made to the soil shortly before weed growth begins. If dense growth is present, remove tops and spray the ground.

Increased contact activity on established weeds may be obtained by the addition of Du Pont Surfactant at the rate of 2 qts. per 100 gals. of spray mixture. Apply as a drenching spray to actively growing weeds during warm weather when daily temperature will exceed 70° F.

**APPLES:** Use "Karmex" alone, or apply as a tank mixture with Du Pont "Sinbar" Weed Killer.

**"Karmex" Alone:** Use only under trees established in the orchard for at least 1 year, do not treat varieties grafted on full-dwarf root stocks. Apply 4 lbs. per acre in the spring (March through May). In the Far West, treatment may be made in winter (December through February), or apply 2 lbs. per acre as a postharvest treatment followed by 2 lbs. in the spring.

**"Karmex" + "Sinbar"**—Use only under trees established in the orchard for at least 2 years. Apply either in the spring or after harvest in the fall before weeds emerge or during early seedling stage of weed growth.

Soil Texture	Lbs. Product Per Acre			
	1 to 2% Organic Matter "Karmex" + "Sinbar"		More Than 2% Organic Matter "Karmex" + "Sinbar"	
Sandy loam	1	+ 1	1½	+ 1½
Loam, silt loam, silt.	1½	+ 1½	2	+ 2
Clay loam, clay	2	+ 2	2	+ 2

Where crop is grown under furrow irrigation or under raised-berm flood irrigation (trees 4" to 6" above waterline), apply only as a band treatment. Do not treat trees planted in the bottom of irrigation furrows, nor trees grown under flat flood or basin irrigation, as injury to trees may result. Where complete weed control to harvest is desired, additional weed control measures may be required during the growing season.

**BANANAS AND PLANTAINS—New Plantings:** To control annual weeds, apply 1½ to 3 lbs. per acre after planting but before weeds emerge. Do not apply to loose soil directly over the planting material.

**Established Plantings:** For control of annuals and for top-kill of perennials such as bermudagrass, bird seed grass and guinea grass, apply 3 to 6 lbs. per acre plus 1 pt. Surfactant WK (or suitable equivalent) per 25 gals. of spray; avoid contact of plants with spray or drift as injury may result. When tall, dense weed growth is present, remove weed growth before application. If application is made to soil free of weeds, omit the surfactant from the spray. Repeat treatment as needed, but do not apply more often than 6-week intervals nor more than a total of 12 lbs. per acre (broadcast basis) in a 12-month period.

**Note:** Do not replant treated areas to any crop within 2 years after last application as injury to subsequent crops may result, except that sugarcane or pineapple may be planted one year after last application.

**BLUEBERRIES, CANEBERRIES AND GOOSEBERRIES:** Use only in fields which have been established for at least 1 year. Do not apply to berries interplanted with fruit trees; do not apply to plants with roots exposed as injury may result. Apply as a band treatment at base of canes or bushes; for spring application, apply before germination and growth of annual weeds.

**Indiana, Michigan and Ohio—Blueberries:** Apply 2 to 4 lbs. per acre in late spring; alternatively, apply 2 lbs. per acre in the fall and repeat at same rate in the spring. **Raspberries:** Apply 3 lbs. per acre in the spring. **Massachusetts—Blueberries:** Apply 2 lbs. per acre in late spring.

**New Jersey—Blueberries:** For control of winter annuals, apply 2 lbs. per acre in October, November or December, or a single application of 2½ lbs. per acre may be applied in early to mid spring.

**California—Raspberries, Blackberries, Boysenberries, Dewberries and Loganberries:** For control of winter annuals, apply 2 lbs. per acre in October or November; repeat at same rate in late spring to control summer annuals. A single application of 3 lbs. per acre in January or February will control both winter and summer annuals in some areas, but the separate fall and spring schedule is preferred.

**Western Oregon and Western Washington—Blueberries, Caneberries and Gooseberries:** Use same schedule as recommended for California.

**CITRUS:** Use only under trees established in the grove for at least 1 year. Time application as indicated for specific areas, except application may be made any time of the year where sprinkler or flood irrigation can be timed to activate the herbicide. Established perennial weeds require other special control procedures. Do not apply under citrus trees that have been subjected to freezing within 6 months.

**NON-CROP WEED CONTROL**

"Karmex" is an effective herbicide for the control of many annual and perennial grasses and herbaceous weeds on non-cropland areas where bare ground is desired. The degree of control and duration of effect vary with the amount of chemical applied, soil texture, rainfall and other conditions.

"Karmex" may be used as a preemergence treatment at any time of the year except when ground is frozen, provided adequate moisture is supplied by rainfall or artificial means to activate the herbicide. Best results are obtained if application is made to the soil shortly before weed growth begins. If dense growth is present, remove tops and spray the ground.

Increased contact activity on established weeds may be obtained by the addition of Du Pont Surfactant WK at the rate of 2 qts. per 100 gals. of spray mixture. Apply as a drenching spray to actively growing weeds during warm weather when daily temperature will exceed 70 F.

Except for small areas, use a fixed boom power sprayer properly calibrated to insure a constant rate application. Mix proper amount of "Karmex" into volume of water necessary to obtain uniform coverage. Surfactant WK is used, dilute with 10 parts of water and add as last ingredient to nearly full tank. Material must be kept in suspension at all times. Agitate by mechanical or hydraulic means in the spray tank; if it passes or return line is used, it should terminate at bottom of tank to minimize foaming. Openings in screen should be equal to or larger than 50 mesh.

**General Weed Control:** To control most weeds for an extended period of time on non-cropland such as utility, highway, pipeline and railroad right of ways, petroleum tank farms, lumberyards, storage areas, industrial plant sites, and around farm buildings—apply 5 to 20 lbs. per acre to control most annual weeds. Use 20 to 60 lbs. per acre for perennial weeds; additional treatment may be required where a longer period of control is desired or when hard-to-kill, deep-rooted perennial weeds such as johnsongrass are present. In low rainfall areas, "Karmex" may not provide satisfactory control of deep-rooted perennial weeds.

For weed control on small areas, use one-half cupful of "Karmex" per 100 sq. ft. for a dosage of approximately 50 lbs. per acre.

**Irrigation and Drainage Ditches:** Apply 5 to 20 lbs. per acre to control most annual weeds; use 20 to 60 lbs. per acre to control both annual and perennial weeds. Apply only when water is not in the ditch. For irrigation ditches, apply during the non-crop season, and when ditch is not in use. To minimize movement of "Karmex" with irrigation water (to avoid possible crop injury), it is essential that the herbicide be fixed in the soil by moisture. Apply before expected seasonal rainfall (if possible when soil in the ditch is still moist). Following treatment, if rainfall has not totaled at least 4 inches, fill ditch with water and allow stand for 72 hours; drain off and waste remaining water before using ditch. Do not treat any ditch in which roots of trees or other desirable plants may extend as injury may result.

**STORAGE AND DISPOSAL**

- 1. Prohibitions:** Do not contaminate water, food, or feed by storage, disposal, or cleaning of equipment. Open dumping is prohibited.
- 2. Pesticide Disposal:** Pesticide or rinse water that cannot be used according to label instructions must be disposed of according to applicable Federal, state, or local procedures.
- 3. Container Disposal:** Completely empty bag by shaking and tapping sides and bottom to loosen clinging particles. Empty residue into application equipment. Then dispose of bag in a sanitary landfill or by incineration if allowed by state and local authorities.

**BEST AVAILABLE COPY**



DU PONT

**KARMEX**

WETTABLE POWDER

# Product Labeling

## DIRECTIONS FOR USE

E. I. DU PONT DE NEMOURS & CO. (INC.)  
BIOCHEMICALS DEPT., WILMINGTON, DEL.  
B 20160 2 83 Made in USA Product in USA

### ACTIVE INGREDIENT:

Diuron [3-(3,4-dichlorophenyl)-1,1-dimethylurea] ..... 80%

INERT INGREDIENTS ..... 20%

EPA Reg. No. 352-247

BEST AVAILABLE COPY

KEEP OUT OF REACH OF CHILDREN.  
PRECAUTIONARY STATEMENTS  
HAZARDS TO HUMANS

### CAUTION! MAY IRRITATE EYES, NOSE, THROAT, AND SKIN.

Avoid breathing dust or spray mist. Avoid contact with skin, eyes, and clothing. Wash thoroughly with soap and water after handling.

#### STATEMENT OF PRACTICAL TREATMENT

If on skin: Wash with plenty of soap and water; get medical attention if irritation persists.  
If in eyes: Flush with plenty of water; get medical attention if irritation persists.

#### ENVIRONMENTAL HAZARDS

Do not apply directly to water. Do not contaminate water by cleaning of equipment or disposal of wastes.

**IMPORTANT**—Injury to or loss of desirable trees or other plants may result from failure to observe the following:

Do not apply (except as recommended for crop use), or drain or flush equipment on or near desirable trees or other plants, or on areas where their roots may extend, or in locations where the chemical may be washed or moved into contact with their roots. Do not use on home plantings of trees, shrubs or herbaceous plants, nor on lawns, walks, driveways, tennis courts, or similar areas. Prevent drift of dry powder or spray to desirable plants. Do not contaminate any body of water. Keep from contact with fertilizers, insecticides, fungicides, and seeds.

Thoroughly clean all traces of "Karmex" from application equipment immediately after use. Flush tank, pump, hoses, and boom with several changes of water after removing nozzle tips and screens (clean these parts separately).

### GENERAL INFORMATION

Du Pont "Karmex" Weed Killer is a wettable powder to be mixed in water and applied as a spray for selective control of weeds in certain crops and for nonselective weed control on non-cropland areas. It is non-corrosive to equipment, nonflammable and nonvolatile.

"Karmex" may be applied to soil prior to emergence of weeds to control susceptible weed seedlings for an extended period of time; the degree of control and duration of effect will vary with the amount of chemical applied, soil texture, rainfall and other conditions. Soils high in clay or organic matter require higher dosages than soil low in clay or organic matter to obtain equivalent herbicide performance. Moisture is required to activate the chemical; best results occur if rainfall (or sprinkler irrigation) occurs within 2 weeks of application.

● **ALFALFA:** Treat only stands established for 1 year or more. Do not apply to seedling alfalfa nor to grass mixture. Do not apply to alfalfa under stress from disease, insect damage, shallow root penetration (such as on shallow hard pans), alkali spots; nor to flooded fields as crop injury may result. Do not apply on snow-covered or frozen ground.

**Idaho, Oregon, Washington:** Use 1½ to 3 lbs. per acre for control of volunteer alfalfa, use 4 lbs. per acre. Apply in fall after alfalfa becomes dormant but no later than mid December.

**California (Dormant and Semi-Dormant Varieties):** Use 1½ to 3 lbs. per acre for control of volunteer alfalfa. Use 4 lbs. per acre. Apply in fall or winter after alfalfa becomes dormant or semi-dormant, but before growth begins in the spring. Crop injury may result if application is made to actively growing alfalfa. For best results, apply before weeds have emerged or become established (2" in height or diameter). For control of established weeds is improved by applying "Karmex" with a suitable contact herbicide registered for such use. Sufficient rainfall for soil activation of "Karmex" is unlikely in California after February. Treated areas may be replanted to any crop after one year from last application if rate does not exceed 2 lbs. per acre.

**Arizona, Nevada:** Use 1½ to 3 lbs. per acre; apply in fall after alfalfa becomes dormant but no later than January.

**Eastern Colorado, Kansas:** For control of tansymustard, apply 1 lb. per acre shortly after emergence of tansymustard in the fall or winter; use 2 lbs. per acre if weeds are 2" to 4" in height. Alternatively, if other weeds are present, apply 2 to 3 lbs. per acre in February or March.

**Other Areas Where Alfalfa Becomes Winter Dormant:** Use 1½ to 3 lbs. per acre (1½ to 2 lbs. per acre in the Appalachian Mountains). Apply in March or early April, but before spring growth begins.

● **ARTICHOKE**—California: Apply 2 to 4 lbs. per acre in late fall or early winter after the last cutting. Apply before weeds germinate or to emerging seedlings. Direct spray to cover the area between the rows at the base of artichoke plants, keeping contact with crop plants at a minimum.

● **ASPARAGUS:** Apply as a band or broadcast treatment. Do not apply to young plants during the growing season (except as noted below), nor to newly seeded asparagus, nor on plants with exposed roots. As severe injury may result. Preemergence weed control will be reduced on high organic matter soils (greater than 5%).

**Established Plantings:** On light sandy soils and other soils low in clay or organic matter, apply 1 to 2 lbs. per acre. On soils high in clay or organic matter, use 2 to 4 lbs. per acre. Two applications may be used; the first application should be made before weeds become established but no earlier than 4 weeks before emergence and no later than the early cutting period (if weeds are controlled into the cutting period). For cultural practices, application may be delayed until immediately after the last cultivation, a second application may be made immediately following completion of harvest provided rainfall is expected. When multiple applications are used in one season, do not exceed 3 lbs. per acre per application. In Washington (irrigated crop), apply a single treatment of 4 lbs. per acre. If treatment is delayed until late fall or early spring, incorporation of the chemical in the top 1" to 2" of soil may substitute for irrigation or rain to activate the herbicide.

**Newly Planted Crowns**—California (San Joaquin Delta): Make a single application of 2 to 4 lbs. per acre on soils high in clay or organic matter, use the lower rate on clay loams and the higher rate on peat soils. Do not use on soils containing less than 2% organic matter. Soil must be settled by rainfall or irrigation prior to treatment. Do not treat crowns planted to a depth of less than 2".

● **BARLEY, WINTER (Drill-Planted)**—Western Oregon and Western Washington: Make a single application of 1½ to 2 lbs. per acre as soon as possible after planting but before emergence of barley. Do not apply to treated areas to any crop within 1 year after last application as injury to subsequent crops may result.

● **BERMUDAGRASS PASTURES (Newly Sprigged):** Apply 1 to 3 lbs. after planting and before emergence of bermudagrass or weeds. Alternatively, for control of emerged annual weeds up to 4" in height, apply ½ to 1 lb. per acre; add 1 pt. Surfactant WK per 25 gals. of spray. If bermudagrass has emerged at the time of treatment, temporary burn of exposed plant parts may occur.

Plant sprigs (stolons) 2" deep in a well-prepared seedbed; do not treat areas where sprigs are planted less than 2" deep as crop injury may result. Do not graze or feed foliage from treated areas to livestock within 30 days after application.

● **BIRDSFOOT TREFLOIL (Lotus)**—Western Oregon: Treat only stands established for at least 1 year. Do not apply to seedling trefoil as injury may result. Make a single application of 2 lbs. per acre when dormant (October 15 to December 15). Do not replant treated areas to any crop within 1 year after application as injury to subsequent crops may result.

● **CORN (FIELD):**

**Postemergence**—Make a single application of ¾ lb. per acre in combination with non-pressure nitrogen solution, if nitrogen solution is not used, apply 1 lb. per acre, add 1 pt. Surfactant WK per 25 gals. of spray. Apply as a directed spray when corn is at least 20" high and weeds are no taller than 3". DO NOT APPLY TO OVER TOP OF CORN. Do not replant to any crop within 1 year, except that cotton, corn and grain sorghum may be planted the spring following treatment.

**Preemergence**—Arkansas, Louisiana, Mississippi and Tennessee: Make a single application of 1½ to 2 lbs. per acre as a broadcast or band treatment after planting but before corn emerges. Plant corn 1 to 2 weeks deep. Do not replant treated areas to crops other than corn on cotton with 4 months following treatment and 6 months following broadcast treatment.



Best results are obtained on succulent weeds growing under conditions of high humidity and temperatures of 70°F or higher. Addition of a surfactant such as Du Pont Surfactant WK to the spray (where recommended) increases contact effects of "Karmex."

Since the effect of "Karmex" varies with soils, uniformity of application, and environmental conditions, it is suggested that growers limit their first use to small areas. Observe all cautions and limitations on labeling of all products used in mixtures.

### NOTICE OF WARRANTY

Du Pont warrants that this product conforms to the chemical description on the label thereof and is reasonably fit for purposes stated on such label only when used in accordance with the directions under normal use conditions. It is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of Du Pont. In no case shall Du Pont be liable for consequential, special or indirect damages resulting from the use or handling of this product. All such risks shall be assumed by the Buyer. DU PONT MAKES NO WARRANTIES OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS STATED ABOVE.

### DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

Du Pont "Karmex" Weed Killer should be used only in accordance with recommendations on this label, or in separate published Du Pont recommendations available through local dealers.


Du Pont will not be responsible for losses or damages resulting from use of this product in any manner not specifically recommended by Du Pont. User assumes all risk associated with such non-recommended use.

Do not use in Kern County, California.

### SELECTIVE USE IN CROPS

**PREEMERGENCE USE (Germinating Weeds):** "Karmex", at recommended rates, controls annual weeds such as:

Broadleaves			Grasses		
1/2 to 1 Lb./Acre	1 1/2 to 2 Lbs./Acre	2 to 6 Lbs./Acre	1/2 to 1 Lb./Acre	1 1/2 to 2 Lbs./Acre	2 to 6 Lbs./Acre
Lambquarters	Annual groundcherry	Ageratum	Barnyardgrass	Annual bluegrass	Annual lovegrass
Pigweed	Annual morningglory	Annual smartweed	(watergrass)	Annual sweet	Annual ryegrass
Purslane	Chickweed	Annual sowthistle	Crabgrass	vernalgrass	Kyllinga
Ragweed	Corn spurry	Corn speedwell		foxtail	Orchardgrass
	Dogfennel	Dayflower		Rattail fescue	Peppergrass
	Fiddleneck	Flora's paintbrush		Red sprangletop	Ricegrass
	(amsinckia)	Hawksbeard		Velvetgrass	Sandbur
	Gromwell	Horseweed			Seeding johnson grass
	Knawel	Kochia			
	Pennycress	Marigold			
	Shepherdspurse	Mexican clover			
	Tansy-mustard	Pineappleweed			
	Wild buckwheat	Pokeweed			
	Wild lettuce	Rabbit tobacco			
	Wild mustard	Spanishneedles			
		Velvetleaf			
		(buttonweed)			
		Wild radish			



Partial control of the following weeds usually occurs at rates stated:

Broadleaves		Grasses	
1 Lb./Acre	4 Lbs./Acre	4 Lbs./Acre	8 to 10 Lbs./Acre
Annual morningglory	Horsenettle	Quackgrass	Guineagrass
Cocklebur			Maidencane
Prickly sida (teaweed)			Pangolagrass
Sesbania			
Sicklepod			

**BEST AVAILABLE COPY**

**POSTEMERGENCE USE (Emerging Seeding Weeds):** "Karmex" at recommended rates, controls annual weeds such as annual morningglory, barnyardgrass (watergrass), crabgrass, crowfoot, goosegrass, pigweed and purslane. Addition of Surfactant WK to the spray (where recommended) increases contact effects of "Karmex". Best results are obtained on succulent weeds growing under conditions of high humidity and temperatures of 70°F or higher.

**EQUIPMENT—SPRAY VOLUMES AND PRESSURES:** Use a tractor mounted fixed-boom power sprayer properly calibrated to a constant speed and rate of delivery. Openings in screens should be equal to or larger than the nozzle. Continuous application of the surfactant is required to provide the best results.

Injury to subsequent crops may result: 1 lb. on loamy sand; 1 1/2 lbs. on sandy loam, 2 lbs. on clay; 2 1/4 lbs. on clay. Injury may occur if "Karmex" is used in conjunction with soil-applied organic phosphorus pesticides. Do not allow livestock to graze treated cotton.

**Preplant—Arizona and California:** Use "Karmex" alone, or apply as a separate operation following preplant broadcast treatment with "Treflan" (incorporated according to directions on "Treflan" label). "Karmex" as a broadcast spray after beds are formed, preirrigated, and final seedbeds prepared. Prior to planting, drag off the tops of the beds and plant in moist soil not treated with "Karmex". Treated beds returned to the bed after planting when irrigation furrows are reformed after cotton has emerged. If more than two furrowing out operations are made prior to lay by, or deep furrows are made early, weed control may be reduced in furrow bottoms. Use at the following rates:

"Karmex" Alone: 1 to 2 1/2 lbs. per acre  
 "Karmex" Following "Treflan":

Soil Texture	Product Per Acre—Preplant	
	"Treflan"	"Karmex"
Sandy loam, loam, silt loam, silt	1 pt.	2 1/2 to 1 lb.
Sandy clay loam, clay loam, silty clay loam, sandy clay, clay	1 1/2 pts.	1 to 1 1/2 lbs.

Note: Seeding disease may weaken plants and increase the possibility of injury from the use of "Karmex". These treatments should be used only in conjunction with a standard fungicide treatment plus a good supplemental soil fungicide program such as Du Pont, Demosan® 65W Fungicide or captan PCNB mixture.

Reg. trademark of Elanco Products Co.  
**Preemergence—U.S., except Arizona, California, and areas west of Interstate 35 or 35W in Texas and Oklahoma:** Use "Karmex" alone or apply as a separate operation following preplant treatment with "Treflan". Apply "Karmex" after planting but before cotton emerges. Do not treat cotton in deep furrows as crop injury may result, use only where cotton is planted on flat or raised seedbeds. Shallow incorporation (no deeper than 1") with a rotary hoe or similar equipment following planting usually improves results, especially during dry weather. A wide press wheel should be used on the planter to provide a level seedbed for subsequent early season postemergence treatments. If moisture is insufficient to activate "Karmex" or if soil becomes crusted before crop emerges, a shallow rotary hoeing (no deeper than 1") should be made before weeds become established.

"Karmex" Alone: Make a single application as a broadcast or band spray, using the following broadcast rates; for band treatment, use proportionately less.

Soil Texture**	Lbs. "Karmex" Per Acre
Loamy sand	2 1/2
Sandy loam, loam, silt loam, silt	1
Sandy clay loam, clay loam, silty clay loam, sandy clay	1 1/4
Silty clay, clay	2

Do not use on soils with less than 1% organic matter as crop injury may result.  
**"Karmex" Following "Treflan" Preplant:** Apply "Treflan" prior to planting as a broadcast or band treatment; incorporate according to directions on "Treflan" label. As a separate operation, apply "Karmex" as a band treatment (14" to 20" wide) after planting but before cotton emerges. Use at the following broadcast rates; for band treatment, use proportionately less. See "Note" under Preplant above.

Soil Texture**	Product Per Acre	
	Preplant "Treflan"	Postemergence "Karmex"
Loamy sand	2 1/2 pt.	2 1/2 lb.
Sandy loam, loam, silt loam, silt	1 pt.	1 lb.
Sandy clay loam, clay loam, silty clay loam, sandy clay, silty clay, clay	1 1/2 pts.	1 1/4 to 2 lbs.

\*Do not use on soils with less than 1% organic matter as crop injury may result.  
**Postemergence—U.S.:** Apply only as a directed spray to cover weed foliage; adjust nozzles to minimize contact of cotton leaves with spray or drift or crop injury may result. DO NOT SPRAY OVER TOP OF COTTON.  
**Early Season—**Apply when cotton is at least 6" tall (at least 12" tall for Western irrigated cotton) when weeds are actively growing and do not exceed 2" in height. Apply as a band treatment at following rates, for each 25 gals. of spray, add 1 pt. Surfactant WK. Two applications may be made if necessary.

Weed Problem (Up to 2" Tall)	Lbs. "Karmex" Per Acre (Broadcast Basis)

Wild radish

registered under  
EPA Reg. No. 352-247

Partial control of the following weeds usually occurs at rates stated:

1 Lb./Acre	4 Lbs./Acre	4 Lbs./Acre	8 to 10 Lbs./Acre
Annual morningglory	Horsenettle	Quackgrass	Guinea grass
Cocklebur			Maidencane
Prickly sida (leaweed)			Pangolagrass
Sesbania			
Sicklepod			

**POSTEMERGENCE USE (Emerging Seedling Weeds):** "Karmex" at recommended rates, controls annual weeds such as annual morningglory, barnyardgrass (watergrass), crabgrass, crowfoot, goosegrass, pigweed and purslane. Addition of Surfactant WK to the spray (where recommended) increases contact effects of "Karmex". Best results are obtained on succulent weeds growing under conditions of high humidity and temperatures of 70°F or higher.

**EQUIPMENT—SPRAY VOLUMES AND PRESSURES:** Use a tractor-mounted fixed-boom power sprayer properly calibrated to a constant speed and rate of delivery. Openings in screens should be equal to or larger than 50 mesh. Continuous agitation in the spray tank is required to keep the material in suspension. Agitate by mechanical or hydraulic means; if by-pass or return line is used, it should terminate at bottom of tank to minimize foaming. Avoid overlapping, and shut off spray booms while starting, turning, slowing or stopping, or injury to the crop may result.

For preemergence application, use 25 to 40 gals. per acre and spray pressure of 30 to 40 psi. For post-emergence application, use sufficient volume (min. 25 gals. per acre) for thorough coverage of weed foliage; use spray pressure of 20 to 25 psi to keep spray drift to a minimum.

**Aerial:** For alfalfa, asparagus, barley (winter), cotton (preplant or preemergence only), grass seed crops, pineapple, sugarcane and wheat (winter), application may be made by aircraft (5 to 10 gals. per acre); avoid overlapping of spray swath and avoid application under conditions where excessive drift may occur. Where land is bedded, make application parallel to rows.

**SPRAY PREPARATION:** Mix proper amount of "Karmex" into necessary volume of water; where use of Du Pont Surfactant WK is recommended, dilute with 10 parts of water and add as last ingredient to nearly full tank.

**USE RATES:** All dosages of "Karmex" are expressed as broadcast rates; for band treatment, use proportionately less. For example, use 1/3 of the broadcast rate when treating a 14" band where row spacing is 42". Where a range of dosages is given, use the lower rate on coarse textured soils (low in clay or organic matter) and the higher rate on fine textured soils (high in clay or organic matter); for postemergence application, use the lower rate on smaller weeds and the higher rate on larger weeds.

**SOIL LIMITATIONS:** Crop injury may result from failure to observe the following:

Unless otherwise directed, do not use on sand, loamy sand, gravelly soils or exposed subsoils; nor on pecans where organic matter is less than 1/2%; nor on alfalfa, apples, artichoke, barley (winter), bermudagrass pasture, citrus, cotton, grapes, oats, olives, papayas, peaches, pears, plumous fern, sorghum, sugarcane, walnuts and wheat (winter) where organic matter is less than 1%; nor on blueberries, birdsfoot trefoil, caneberries, gooseberries, macadamia nuts and peppermint where organic matter is less than 2%.

Preemergence weed control will be reduced on high organic matter soils (greater than 5%, such as peat or muck).

**REPLANTING:** Unless otherwise directed, do not replant treated areas to any crop within 2 years after last application as injury to subsequent crops may result.

**FIELD CROPS**  
(See Soil Limitations)

A good seedbed must be prepared before preemergence use of "Karmex" as crop injury may result if application is made to ground which is cloddy or compacted resulting in improperly planted seed. Plant seed to depth specified. Unless otherwise directed, surface of the soil should not be cultivated or disturbed after application of "Karmex" and before emergence of the crop as weed control may be reduced and crop injury may result. However, if moisture is sufficient to activate the herbicide, a shallow cultivation (rotary hoe) should be made after emergence of crops while weeds are small enough to be controlled by mechanical means.

BEST AVAILABLE COPY

Soil Texture**	Product Per Acre	
	Preplant "Trellan"	Preemergence "Karmex"
Loamy sand	1 pt	1 lb
Sandy loam, loam, silt loam, silt	1 pt	1 lb
Sandy clay loam, clay loam, silty clay loam, sandy clay, silty clay, clay	1 1/2 pts	1 1/4 to 2 lbs

\*\*Do not use on soils with less than 1% organic matter as crop injury may result.

**Postemergence—U.S.:** Apply only as a directed spray to cover weed foliage; adjust nozzles to minimize contact of cotton leaves with spray or drift or crop injury may result. **DO NOT SPRAY OVER TOP OF COTTON.**

**Early Season—**Apply when cotton is at least 6" tall (at least 12" tall for Western irrigated cotton), a when weeds are actively growing and do not exceed 2" in height. Apply as a band treatment at follow rates, for each 25 gals. of spray, add 1 pt Surfactant WK. Two applications may be made if needed.

Weed Problem (Up to 2" Tall)	Lbs. "Karmex" Per Acre (Broadcast Basis)
Annual grasses	1
Pigweed	1

For control of seedling perennial grasses such as johnsongrass and partial control of nutsedge or wh weed growth is under drought stress or as high as 4", add 2 to 3 1/2 lbs. disodium methylarsonate (DSM 63% anhydrous or equivalent) to above spray mixture. If DSMA is used, do not apply after first bloom.

**Late Season (Lay-By)—**Apply 1 to 1 1/2 lbs. per acre (1 to 2 lbs. in Arizona and California) when cotton is least 12" tall (at least 20" tall for Pima S-2). For control of germinating weed seedlings, apply to beneath cotton plants and between rows immediately after last cultivation. In irrigated cotton, best weed control is obtained if the field is irrigated within 3 to 4 days after application, thoroughly wet the surface of the ground over the row to carry the herbicide into the root zone of germinating weeds. Alternatively, control of emerged annual weeds (up to 4" in height) at lay-by time, make a single application in combination with Surfactant WK (1 pt. per 25 gals. spray), or use 1/2 to 3/4 lb. "Karmex" (plus surfactant) per acre and repeat later if needed.

**Replanting:** If initial seeding fails to produce a stand, cotton may be replanted in soil treated preplant preemergence with "Karmex", alone or following "Trellan". Wherever possible, avoid disturbing original bed. If necessary to rework soil before replanting, use shallow cultivation such as disking, do not plow or move soil into the original drill area. Plant seed at least 1" deep. Do not retreat field with a second preplant or preemergence application during the same crop year as injury to the crop may result.

**Subsequent Crops:**

"Karmex"—Type of Application	Crops That May Follow Treated Cotton
Band preemergence or postemergence	Any crop 4 months after last application.
Band preemergence plus postemergence -or- Broadcast preemergence (and preplant) -or- Broadcast preemergence plus band postemergence	Cotton, soybeans, corn or grain sorghums (not sorgo or forage sorghums nor grass sorghums) the next spring. Do not replant treated areas to any other crop within one year after last application as injury to subsequent crops may result.
Broadcast postemergence (lay-by)	Cotton, corn, grain sorghums (not sorgos or forag sorghums nor grass sorghums) the next spring. Do not replant treated areas to any other crop within one year after last application as injury to subsequent crop may result.

For subsequent crops in fields where "Trellan" is used, follow instructions on "Trellan" label.

Continued on reverse

KARMEX® WEED KILLER  
 FOR USE IN WALNUTS (ENGLISH)  
 IN THE STATES OF OREGON AND WASHINGTON

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

Du Pont Karmex® Weed Killer is recommended for weed control in English Walnuts in Oregon and Washington.

HOW TO USE

Use only under trees established in the orchard for at least one year. As an initial treatment, apply 3 to 5 lbs. per acre after the orchard has been laid up in final form (non-tillage program) in late fall or early winter; retreat annually with 2 to 3 lbs. per acre. Alternatively, apply 2 lbs. per acre in October or November and repeat at the same rate in March or April.

NOTES: Do not use on sand, loamy sand, gravelly soils or exposed subsoils, nor where organic matter is less than 1%. Do not graze livestock in treated orchards and groves.


IMPORTANT

BEFORE USING "KARMEX", READ AND CAREFULLY OBSERVE THE CAUTIONARY STATEMENTS, AND ALL OTHER INFORMATION APPEARING ON THE PRODUCT LABEL.

This bulletin contains new or supplemental instructions for use of this product which may not appear on the package label. Follow the instructions carefully.

This labeling must be in the possession of the user at the time of pesticide application.

081183  
 042484





# AGRICULTURAL BULLETIN

19/ 6.0




KARMEX® WEED KILLER  
FILBERTS

SUPPLEMENTAL LABELING  
EPA REG. NO. 352-247

KARMEX® WEED KILLER  
FOR USE IN FILBERTS

### DIRECTIONS FOR USE



It is a violation of federal law to use this product in a manner inconsistent with its labeling.

Du Pont Karmex® Weed Killer is recommended for control of certain weeds in filbert orchards established for at least one year.

### HOW TO USE

Apply "Karmex" as a directed spray, avoiding contact on the foliage and fruit with spray or drift. Make an initial treatment of 4 to 5 lbs per acre in the late fall or early winter after harvest. Repeat annually with 3 to 4 lbs per acre, or, apply 2 lbs per acre in October or November after harvest and repeat at the same rate in March or April.

### NOTES:

Do not apply when nuts are on the ground.

Do not graze livestock in treated orchards.

Do not use on light sandy soils.

If trees are planted on hillsides, the elimination of weeds and ground cover may cause excessive soil erosion. Under these conditions strip applications of Karmex® (at proportionately lower rates) may be made near the trees or to the tree rows perpendicular to the slope.

### IMPORTANT

BEFORE USING KARMEX®, READ AND CAREFULLY OBSERVE THE CAUTIONARY STATEMENTS AND ALL OTHER INFORMATION APPEARING ON THE PRODUCT LABEL.

This bulletin contains new or supplemental instructions for use of this product which may not appear on the package label. Follow the instructions carefully.

This labeling must be in the possession of the user at the time of pesticide application.

081583



# AGRICULTURAL BULLETIN

20/20



KARMEX® WEED KILLER  
WALNUTS  
OREGON AND WASHINGTON

SUPPLEMENTAL LABELING  
EPA REG. NO. 352-247

KARMEX® WEED KILLER  
FOR USE IN WALNUTS (ENGLISH)  
IN THE STATES OF OREGON AND WASHINGTON

### DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

Du Pont Karmex® Weed Killer is recommended for weed control in English Walnuts in Oregon and Washington.

### HOW TO USE

Use only under trees established in the orchard for at least one year. As an initial treatment, apply 3 to 5 lbs. per acre after the orchard has been laid up in final form (non-tillage program) in late fall or early winter; retreat annually with 2 to 3 lbs. per acre. Alternatively, apply 2 lbs. per acre in October or November and repeat at the same rate in March or April.

NOTES: Do not use on sand, loamy sand, gravelly soils or exposed subsoils, nor where organic matter is less than 1%. Do not graze livestock in treated orchards and groves.

### IMPORTANT

BEFORE USING "KARMEX", READ AND CAREFULLY OBSERVE THE CAUTIONARY STATEMENTS, AND ALL OTHER INFORMATION APPEARING ON THE PRODUCT LABEL.

This bulletin contains new or supplemental instructions for use of this product which may not appear on the package label. Follow the instructions carefully.

This labeling must be in the possession of the user at the time of pesticide application.

081183  
042484

